BURNSIDE'S THEOREM FOR HOPF ALGEBRAS

D. S. PASSMAN AND DECLAN QUINN

(Communicated by Ken Goodearl)

ABSTRACT. A classical theorem of Burnside asserts that if χ is a faithful complex character for the finite group G , then every irreducible character of G is a constituent of some power χ^n of χ . Fifty years after this appeared, Steinberg generalized it to a result on semigroup algebras $K[G]$ with K an arbitrary field and with G a semigroup, finite or infinite. Five years later, Rieffel showed that the theorem really concerns bialgebras and Hopf algebras. In this note, we simplify and amplify the latter work.

Let K be a field and let A be a K -algebra. A map $\Delta: A \rightarrow A \otimes A$ is said to be a comultiplication on A if Δ is a coassociative K -algebra homomorphism. For convenience, we call such a pair (A, Δ) a b -algebra. Admittedly, this is rather nonstandard notation. One is usually concerned with bialgebras, that is, algebras which are endowed with both a comultiplication Δ and a counit $\epsilon: A \rightarrow K$. However, semigroup algebras are not bialgebras in general, and the counit rarely comes into play here. Thus it is useful to have a name for this simpler object.

Now a b -algebra homomorphism $\theta: A \rightarrow B$ is an algebra homomorphism which is compatible with the corresponding comultiplications, and the kernel of such a homomorphism is called a b -ideal. It is easy to see that I is a b -ideal of A if and only if $I \triangleleft A$ with $\Delta(I) \subseteq I \otimes A + A \otimes I$. Of course, the b -algebra structure can be used to define the tensor product of A -modules. Specifically, if V and W are left A -modules, then A acts on $V \otimes W$ via $a(v \otimes w) = \Delta(a)(v \otimes w)$ for all $a \in A$, $v \in V$, $w \in W$. Notice that if I is a b -ideal of A , then the set of all A -modules V with $\text{ann}_A V \supseteq I$ is closed under tensor product. Conversely, we have

Proposition 1. *Let A be a b -algebra and let $\mathcal{F}$ be a family of A -modules closed under tensor product. Then*

$$I = \bigcap_{V \in \mathcal{F}} \text{ann}_A V$$

is a b -ideal of A .

Proof. Certainly I is an ideal of A . Now let $X = \bigoplus_{V \in \mathcal{F}} V$ be the direct sum of the modules in $\mathcal{F}$. Then X is an A -module and $\text{ann}_A X = \bigcap_{V \in \mathcal{F}} \text{ann}_A V =$

Received by the editors February 17, 1993 and, in revised form, April 27, 1993.
1991 *Mathematics Subject Classification.* Primary 16W30, 16S34, 16S30.

© 1994 American Mathematical Society
0002-9939/94 \$1.00 + \$.25 per page

I . Furthermore, since $X \otimes X = \oplus_{V, W \in \mathcal{F}} V \otimes W$ and since each $V \otimes W \in \mathcal{F}$, it follows that I annihilates $X \otimes X$. In other words,

$$\Delta(I) \subseteq \text{ann}_{A \otimes A} X \otimes X = I \otimes A + A \otimes I$$

and I is a b-ideal of A . $\square$

The assumption that $\mathcal{F}$ is closed under tensor product can be weakened somewhat in the above. Indeed, suppose that for each $V, W \in \mathcal{F}$ there exists $U \in \mathcal{F}$ with $\text{ann}_A U \subseteq \text{ann}_A V \otimes W$. Then certainly $I \subseteq \text{ann}_A U$ annihilates $V \otimes W$, so I annihilates $X \otimes X$ and hence I is a b-ideal of A .

Now if (A, Δ, ϵ) is a bialgebra with counit ϵ , then I is a bi-ideal of A if and only if it is a b-ideal with $\epsilon(I) = 0$. Furthermore, we can trivially guarantee that the ideal I of the previous proposition satisfies $\epsilon(I) = 0$ by including the principal module K_ϵ in the set $\mathcal{F}$. Thus we have

Proposition 1*. *Let A be a bialgebra and let $\mathcal{F}$ be a family of A -modules closed under tensor product. If $K_\epsilon \in \mathcal{F}$, then*

$$I = \bigcap_{V \in \mathcal{F}} \text{ann}_A V$$

is a bi-ideal of A .

Since the coassociativity of Δ guarantees that the tensor product of A -modules is associative, it makes sense to define the n th tensor power of V by

$$V^{\otimes n} = V \otimes V \otimes \cdots \otimes V \quad (n \text{ times})$$

for all $n \geq 1$. Here, $V^{\otimes 1} = V$ and $V^{\otimes m} \otimes V^{\otimes n} = V^{\otimes(m+n)}$ for all $m, n \geq 1$. It is now a simple matter to prove the following result of [Ri].

Corollary 2. *Let A be a b-algebra and let V be an A -module. If $\text{ann}_A V$ contains no nonzero b-ideal, then $\mathcal{F}(V) = \oplus_{n=1}^{\infty} V^{\otimes n}$ is a faithful A -module.*

Proof. $\mathcal{F} = \{V^{\otimes n} \mid n = 1, 2, \dots\}$ is a set of A -modules which is clearly closed under tensor product. Thus, by Proposition 1,

$$I = \bigcap_{n=1}^{\infty} \text{ann}_A V^{\otimes n} = \text{ann}_A \mathcal{F}(V)$$

is a b-ideal of A . But $I \subseteq \text{ann}_A V^{\otimes 1} = \text{ann}_A V$, so the hypothesis implies that $I = 0$ and hence that $\mathcal{F}(V)$ is faithful. $\square$

If A is a bialgebra, then one usually defines $V^{\otimes 0}$ to equal K_ϵ , since the latter module behaves like the identity element under tensor product. Thus we have

Corollary 2*. *Let A be a bialgebra and let V be an A -module. If $\text{ann}_A V$ contains no nonzero bi-ideal, then $\mathcal{F}^*(V) = \oplus_{n=0}^{\infty} V^{\otimes n}$ is a faithful A -module.*

Let V be an A -module. If J is an ideal of A contained in $\text{ann}_A V$, then we can think of V as having been lifted from an A/J -module. In particular, V is faithful if and only if it is not lifted from any proper homomorphic image of A . Similarly, if A is a b-algebra, we might say that V is b-faithful if it is

not lifted from any proper b -algebra homomorphic image of A . In other words, V is b -faithful if and only if $\text{ann}_A V$ contains no nonzero b -ideal of A . Thus Corollary 2 asserts that any b -faithful module V gives rise to the faithful tensor module $\mathcal{F}(V)$. This is essentially Burnside's Theorem.

Let us look at some examples. To start with, recall that a multiplicative semigroup G is a set having an associative multiplication and an identity element 1. Semigroups may contain a zero element $0 \neq 1$ satisfying $0g = g0 = 0$ for all $g \in G$, and as usual we let $G^\# = G \setminus \{0\}$ denote the set of nonzero elements of G . The semigroup algebra $K[G]$ is then a K -vector space with basis $G^\#$ and with multiplication inherited from that of G . Notice that the zero element of G , if it exists, is identified with the zero element of $K[G]$. Furthermore, $K[G]$ is a b -algebra with Δ defined by $\Delta(g) = g \otimes g$ for all $g \in G^\#$. Given this comultiplication, it is easy to see that the only possible bialgebra structure on $K[G]$ would have counit ϵ given by $\epsilon(g) = 1$ for all $g \in G^\#$. But then, ϵ is an algebra homomorphism if and only if $G^\#$ is multiplicatively closed, or equivalently if and only if there are no zero divisors in G . In other words, most semigroup algebras are just not bialgebras in this way.

If H is also a semigroup, then a semigroup homomorphism $\theta: G \rightarrow H$ preserves the multiplication and, by definition, it satisfies $\theta(1) = 1$ and $\theta(0) = 0$ if G has a zero element. In particular, it follows that θ extends to a K -algebra homomorphism $\bar{\theta}: K[G] \rightarrow K[H]$ which is clearly a b -algebra map. Hence $\ker \bar{\theta}$ is a b -ideal of $K[G]$. As is well known, these are the only possible b -ideals. Since the argument is so simple, we briefly sketch it here.

Let I be a b -ideal of $K[G]$ and let $\bar{\phi}$ be the b -algebra epimorphism defined by $\bar{\phi}: K[G] \rightarrow K[G]/I = C$. Then $H = \bar{\phi}(G)$ is a multiplicative subsemigroup of C and, since $\bar{\phi}$ is a b -algebra homomorphism, it is easy to see that H consists of group-like elements. In particular, it follows from [Sw, Proposition 3.2.1(b)] that $H^\#$ is a linearly independent subset of C . Furthermore, since $G^\#$ spans $K[G]$, we know that $H^\#$ spans C . Thus it is clear that $C = K[H]$ and that the map $\bar{\phi}: K[G] \rightarrow K[H]$ is the natural extension of the semigroup epimorphism $\phi: G \rightarrow H$, namely, the restriction of $\bar{\phi}$ to G . Since $I = \ker \bar{\phi}$, this fact is proved.

By combining the above with Corollary 2, we can quickly obtain Steinberg's generalization of the classical result of Burnside [B, §226]. The original Burnside theorem concerned modules for the complex group algebra $\mathbb{C}[G]$ with $|G| < \infty$, and the proof used the character theory of finite groups. The argument in [St] is more transparent and, of course, it is more general. But the following proof, due to Rieffel in [Ri], shows precisely why the G -faithfulness assumption on the $K[G]$ -module V is both natural and relevant.

Let G be a semigroup and let V be a $K[G]$ -module. We say that G acts faithfully on V if for all distinct $g_1, g_2 \in G$ we have $(g_1 - g_2)V \neq 0$. Of course, if G is a group, then this condition is equivalent to $(g - 1)V \neq 0$ for all $1 \neq g \in G$.

Theorem 3. *Let G be a semigroup and let G act faithfully on the $K[G]$ -module V . Then $K[G]$ acts faithfully on the tensor module $\mathcal{F}(V) = \bigoplus_{n=1}^{\infty} V^{\otimes n}$.*

Proof. Let I be a b -ideal of $K[G]$ contained in $\text{ann}_{K[G]} V$. As we observed, there exists a semigroup epimorphism $\phi: G \rightarrow H$ such that I is the kernel of the corresponding algebra map $\bar{\phi}: K[G] \rightarrow K[H]$. If $I \neq 0$, then ϕ cannot be

one-to-one on G and hence there exist distinct $g_1, g_2 \in G$ with $\bar{\phi}(g_1 - g_2) = 0$. In particular, this implies that $g_1 - g_2 \in I$, so $(g_1 - g_2)V = 0$, contradicting the fact that G is faithful on V . In other words, the G -faithfulness assumption implies that $\text{ann}_{K[G]} V$ contains no nonzero b-ideal. Corollary 2 now yields the result. $\square$

An analogous result holds for enveloping algebras. For simplicity of notation, let us assume that either

- (1) K is a field of characteristic 0, L is a Lie algebra over K , and $U(L)$ is its enveloping algebra, or
- (2) K has characteristic $p > 0$, L is a restricted Lie algebra over K , and $U(L)$ is its restricted enveloping algebra.

In either case, $U(L)$ is a b-algebra, and in fact a Hopf algebra, with comultiplication determined by $\Delta(\ell) = \ell \otimes 1 + 1 \otimes \ell$ for all $\ell \in L$. Furthermore, if H is a second (restricted) Lie algebra and if $\theta: L \rightarrow H$ is a (restricted) Lie algebra homomorphism, then θ extends uniquely to a b-algebra homomorphism $\bar{\theta}: U(L) \rightarrow U(H)$. In particular, $\ker \bar{\theta}$ is a b-ideal of $U(L)$. As is well known, the converse is also true, namely, every b-ideal of $U(L)$ arises in this manner. The argument for this is elementary and similar to the one for semigroup rings. A sketch of the proof is as follows.

Let I be a b-ideal of $U(L)$ and let $\bar{\phi}$ be the b-algebra epimorphism defined by $\bar{\phi}: U(L) \rightarrow U(L)/I = C$. Then $H = \bar{\phi}(L)$ is a (restricted) Lie subalgebra of C and H generates C as a K -algebra. In particular, if $\{h_i \mid i \in \mathcal{J}\}$ is a basis for H , indexed by the ordered set $(\mathcal{J}, <)$, then C is spanned by monomials of the form $h_{i_1}^{e_1} h_{i_2}^{e_2} \cdots h_{i_n}^{e_n}$ with $i_1 < i_2 < \cdots < i_n$ and with integers $e_j \geq 0$. Furthermore, when $\text{char } K = p > 0$ and L is restricted, then $e_j < p$ for all j . Since $\bar{\phi}$ is a b-algebra epimorphism, it follows that the elements of H are primitive. Thus, by the work of [Sw, Chapter 13], these straightened monomials are K -linearly independent and therefore $C = U(H)$. In other words, the map $\bar{\phi}: U(L) \rightarrow U(H)$ is the natural extension of the (restricted) Lie algebra epimorphism $\phi: L \rightarrow H$ where, of course, ϕ is the restriction of $\bar{\phi}$ to L . Since $I = \ker \bar{\phi}$, this fact is proved.

Now let V be a $U(L)$ -module. We say that L acts faithfully on V if, for all $0 \neq \ell \in L$, we have $\ell V \neq 0$. The Lie algebra analog of the preceding result is then

Theorem 4. *Let $U(L)$ be a (restricted) enveloping algebra satisfying (1) or (2) above. If L acts faithfully on the $U(L)$ -module V , then $U(L)$ acts faithfully on the tensor module $\mathcal{T}(V) = \bigoplus_{n=1}^{\infty} V^{\otimes n}$.*

As indicated in [M], a theorem of this nature can be used to prove the following interesting result of Harish-Chandra [H, Theorem 1]. Recall that a K -algebra A is residually finite if the collection of its ideals I of finite codimension has intersection equal to 0. In other words, these algebras are precisely the subdirect products of finite-dimensional K -algebras.

Corollary 5. *If L is a finite-dimensional Lie algebra over a field K of characteristic 0, then $U(L)$ is residually finite.*

Proof. By Ado's theorem (see [J, §VI.2]), $A = U(L)$ has a finite K -dimensional module V on which L acts faithfully. Thus, the preceding theorem implies

that $0 = \text{ann}_A \mathcal{T}(V) = \bigcap_{n=1}^{\infty} I_n$, where $I_n = \text{ann}_A V^{\otimes n}$. But each $V^{\otimes n}$ is a finite-dimensional A -module, so $I_n = \text{ann}_A V^{\otimes n}$ is an ideal of A of finite codimension, and the result follows. $\square$

If L is a finite-dimensional restricted Lie algebra, then its restricted enveloping algebra $U(L)$ is also finite dimensional. Thus the characteristic $p > 0$ analog of the above is trivial. On the other hand, infinite-dimensional analogs in all characteristics are obtained in [M].

In the remainder of this paper we will restrict our attention to finite-dimensional Hopf algebras. To start with, a Hopf algebra (A, Δ, ϵ, S) is a bialgebra with antipode $S: A \rightarrow A$, and a Hopf ideal is the kernel of a Hopf algebra homomorphism. It is easy to see that $I \triangleleft A$ is a Hopf ideal if and only if it is a b-ideal with $\epsilon(I) = 0$ and $S(I) \subseteq I$. Similarly, a K -subalgebra B of A is a Hopf subalgebra if and only if it is a b-subalgebra which is closed under the antipode S . Of course, the b-subalgebra condition means that $\Delta(B) \subseteq B \otimes B$. The following is a special case of a surprising result due to Nichols [N, Theorem 1]. A simple proof of the subalgebra case can also be found in [Ra, Lemma 1].

Lemma 6. *If A is a finite-dimensional Hopf algebra, then any b-subalgebra of A is a Hopf subalgebra and any b-ideal of A is a Hopf ideal of A .*

Proof. Let B denote either a b-subalgebra of A or a b-ideal of A . Furthermore, let $E = \text{Hom}_K(A, A)$ be the convolution algebra of A and set

$$F = \{ f \in E \mid f(B) \subseteq B \}.$$

Certainly F is a K -subspace of E and, in fact, F is closed under convolution multiplication. To see the latter, let $f, g \in F$. If B is a b-subalgebra of A , then $\Delta(B) \subseteq B \otimes B$ implies that

$$(f * g)(B) \subseteq f(B)g(B) \subseteq B^2 = B.$$

On the other hand, if B is a b-ideal of A , then $\Delta(B) \subseteq A \otimes B + B \otimes A$ implies that

$$(f * g)(B) \subseteq f(A)g(B) + f(B)g(A) \subseteq AB + BA = B$$

since $B \triangleleft A$.

Now observe that the identity map Id is contained in F . Thus, by the above, F contains the convolution powers Id^{*n} of Id for all $n > 0$. Furthermore, since A is finite dimensional, E is also finite dimensional and hence the map Id is algebraic over K . In particular, for some $m \geq 1$, we can write Id^{*m} as a finite K -linear combination of the powers Id^{*i} with $i < m$. But Id has convolution inverse S , so by multiplying the expression for Id^{*m} by S^{*m} and by $S^{*(m+1)}$ in turn, we deduce first that $\epsilon = \text{Id}^{*0} \in F$ and then that $S = \text{Id}^{*(-1)} \in F$. In other words, $\epsilon(B) \subseteq B$ and $S(B) \subseteq B$.

Finally, if B is a b-subalgebra of A , then $S(B) \subseteq B$ implies that B is a Hopf subalgebra. On the other hand, if B is a b-ideal of A , then $\epsilon(B) \subseteq B \cap K = 0$. Thus, since $S(B) \subseteq B$, we conclude that B is a Hopf ideal of A . $\square$

The preceding result is false in general for infinite-dimensional Hopf algebras. Some rather complicated counterexamples appear in [N].

Theorem 7. *Let A be a finite-dimensional Hopf algebra.*

- (i) *If $\mathcal{F}$ is a family of A -modules which is closed under tensor product, then $\bigcap_{V \in \mathcal{F}} \text{ann}_A V$ is a Hopf ideal of A .*
- (ii) *Suppose V is an A -module whose annihilator contains no nonzero Hopf ideal of A . Then $\mathcal{F}(V) = \bigoplus_{n=1}^{\infty} V^{\otimes n}$ is a faithful A -module.*

This follows immediately from Proposition 1, Corollary 2, and Lemma 6. We can now obtain some consequences of interest. First, recall that an A -module V is semisimple if it is a direct sum of simple modules.

Corollary 8. *If A is a finite-dimensional Hopf algebra, then the set of semisimple A -modules is closed under tensor product if and only if the Jacobson radical $J(A)$ is a Hopf ideal of A .*

Proof. Let $\mathcal{F}$ be the set of all semisimple A -modules. If $\mathcal{F}$ is closed under $\otimes$, then Theorem 7(i) implies that $J(A) = \bigcap_{V \in \mathcal{F}} \text{ann}_A V$ is a Hopf ideal of A . Conversely, if $J(A)$ is a Hopf ideal, then $\mathcal{F}$ consists of all the modules for the Hopf algebra $A/J(A)$ and therefore $\mathcal{F}$ is surely closed under tensor product. $\square$

In a similar manner, we prove

Corollary 9. *Let A be a finite-dimensional semisimple Hopf algebra and let $\mathcal{F}$ be a family of simple A -modules. Suppose that, for all $V, W \in \mathcal{F}$, every irreducible submodule of $V \otimes W$ is contained in $\mathcal{F}$. Then $I = \bigcap_{V \in \mathcal{F}} \text{ann}_A V$ is a Hopf ideal of A and $\mathcal{F}$ is the set of all simple A/I -modules.*

Proof. Let $\mathcal{F}$ be the set of all finite direct sums (with multiplicities) of elements of $\mathcal{F}$. Since A is semisimple, the hypothesis implies that $\mathcal{F}$ is closed under tensor product. Hence, by Theorem 7(i), $I = \bigcap_{V \in \mathcal{F}} \text{ann}_A V = \bigcap_{W \in \mathcal{F}} \text{ann}_A W$ is a Hopf ideal of A . Furthermore, since A/I is semisimple, it follows that $\mathcal{F}$ must be the set of all simple A/I -modules. $\square$

Our final consequence uses the fact that any finite-dimensional Hopf algebra A is a Frobenius algebra [LS, §5] and hence that every simple A -module is isomorphic to a minimal left ideal of A .

Corollary 10. *Let A be a finite-dimensional Hopf algebra and let V be an A -module whose annihilator contains no nonzero Hopf ideal of A . Then every simple A -module is isomorphic to a submodule of $V^{\otimes n}$ for some $n \geq 1$.*

Proof. It follows from Theorem 7(ii) that $\mathcal{F}(V) = \bigoplus_{n=1}^{\infty} V^{\otimes n}$ is a faithful A -module. Now let W be a simple A -module, so that W is isomorphic to a minimal left ideal $L \subseteq A$. Since $L \neq 0$, we have $L\mathcal{F}(V) \neq 0$ and hence $LV^{\otimes n} \neq 0$ for some $n \geq 1$. In particular, there exists $u \in V^{\otimes n}$ with $Lu \neq 0$. But then the minimality of L implies that $W \cong L \cong Lu \subseteq V^{\otimes n}$, as required. $\square$

ACKNOWLEDGMENT

In closing, the authors would like to thank S. Montgomery and M. Cohen for informing us about references [M] and [Ri], respectively.

REFERENCES

- [B] W. Burnside, *Theory of groups of finite order* (reprint of 2nd edition, 1911), Dover, New York, 1955.
- [H] Harish-Chandra, *On representations of Lie algebras*, Ann. of Math. (2) **50** (1949), 900–915.
- [J] N. Jacobson, *Lie algebras*, Interscience, New York, 1962.
- [LS] R. G. Larson and M. E. Sweedler, *An associative orthogonal bilinear form for Hopf algebras*, Amer. J. Math. **91** (1969), 75–94.
- [M] W. Michaelis, *Properness of Lie algebras and enveloping algebras I*, Proc. Amer. Math. Soc. **101** (1987), 17–23.
- [N] W. D. Nichols, *Quotients of Hopf algebras*, Comm. Algebra **6** (1978), 1789–1800.
- [Ra] D. E. Radford, *On the quasitriangular structure of a semisimple Hopf algebra*, J. Algebra **141** (1991), 354–358.
- [Ri] M. A. Rieffel, *Burnside's theorem for representations of Hopf algebras*, J. Algebra **6** (1967), 123–130.
- [St] R. Steinberg, *Complete sets of representations of algebras*, Proc. Amer. Math. Soc. **13** (1962), 746–747.
- [Sw] M. E. Sweedler, *Hopf Algebras*, Benjamin, New York, 1969.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN 53706

E-mail address: `passman@math.wisc.edu`

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13244

E-mail address: `dpquinn@mailbox.syr.edu`