ON FREE GROUP ALGEBRAS IN DIVISION RINGS WITH UNCOUNTABLE CENTER

J. GONCALVES AND M. SHIRVANI

(Communicated by Ken Goodearl)

ABSTRACT. Let D be a division algebra with uncountable center k . If D contains a noncommutative free k -algebra, then D also contains the k -group algebra of the free group of rank 2.

1. INTRODUCTION

Let D be a division ring with center k . In [7] Makar-Limanov states the following conjecture:

(ML) *If D is finitely-generated and infinite dimensional over k , then D contains a 2-generator free algebra.*

An earlier claim about the existence of the free subgroups in division rings was stated by Lichtman [2]:

(L) *The multiplicative group of a noncommutative division ring contains a free group of rank 2.*

Evidence for (ML) can be found in [1], [5], [11], [12], and for (L) in [2], [3]. Recently, Makar-Limanov [11] surprisingly proved that the ring of fractions of the first Weyl algebra A_1 , over the complex field $\mathbb{C}$, contains a $\mathbb{C}$ -free group algebra (= the group algebra of a free group over $\mathbb{C}$).

In this note we show that this result is not accidental, but a general phenomenon. We will use a reasoning similar to the one in [9] to prove:

Theorem. *If k is uncountable and D contains a k -free algebra of rank 2, then D contains a k -free group algebra.*

Thus, we obtain new examples of division rings containing free group algebras. In particular, if k is uncountable, then (ML) implies (L).

We observe that A_1 can be regarded as a skew polynomial ring over the field of rational functions $\mathbb{C}(t)$. Therefore the existence of free group algebras in the ring of fractions of A_1 follows from the existence of free algebras in the ring of fractions of skew polynomial rings. (See [1], [5] for more on this.)

2. PROOF OF THE THEOREM

The lemma below is proved in [12], Lemma 1, for algebras inside domains. It is also true for free group algebras.

Received by the editors September 21, 1993 and, in revised form, July 25, 1994.

1991 *Mathematics Subject Classification.* Primary 16A39; Secondary 16A06, 16A48.

Key words and phrases. Division ring, free algebra, free group algebra.

Lemma 2.1. *Let A be a domain with prime subfield k_0 , and let k be any central subfield of A . If $x_1, x_2 \in A$, then x_1 and x_2 are free over k_0 if and only if they are free over k .* $\square$

We also need the following:

Lemma 2.2. *Let $D[x]$ be the polynomial ring in the central indeterminate x , and let $D(x)$ be its ring of fractions. If $f \in D(x)$ and $f(z) = 0$ for infinitely many $z \in k$, then f is identically zero.*

Proof. See [9], Lemma 1. $\square$

Let L be a field, and let us denote by $L\{x_1, x_2\}$ the free L -group algebra on the generators x_1 and x_2 . Let $L\langle y_1, y_2 \rangle$ be the free L -algebra in the generators y_1 and y_2 , and let $L\langle y_1, y_2 \rangle[[\lambda_1, \lambda_2]]$ be the ring of formal power series in the central indeterminates λ_1 and λ_2 over $L\langle y_1, y_2 \rangle$. With this notation we have:

Lemma 2.3. *The homomorphism ψ from $L\{x_1, x_2\}$ to $L\langle y_1, y_2 \rangle[[\lambda_1, \lambda_2]]$ defined by*

$$\psi(x_i) = 1 + \lambda_i y_i, \quad \psi(x_i^{-1}) = \sum_{j=0}^{\infty} (-\lambda_i y_i)^j, \quad i = 1, 2,$$

is injective.

Proof. See [10], Lemma 1. $\square$

Proof of the theorem. Let a and b be elements of D that generate a free k -algebra. We claim that there exist σ and τ in k such that $1 + \sigma a$ and $1 + \tau b$ generate a free k -group algebra. In view of Lemma 2.1 it is enough to show that they generate a free k_0 -group algebra. Fix $\sigma \neq -a^{-1}$. Since $k_0\{x_1, x_2\}$ is countable but k is not, there exists $0 \neq f_\sigma(x_1, x_2) \in k_0\{x_1, x_2\}$ such that for uncountably many $\tau \in k \setminus \{-b^{-1}\}$ we have

$$f_\sigma(1 + \sigma a, 1 + \tau b) = 0.$$

By Lemma 2.2 we may replace τ by the central variable λ_2 , obtaining the identity

$$f_\sigma(1 + \sigma a, 1 + \lambda_2 b) = 0 \quad \text{in } D(\lambda_2).$$

Arguing analogously, we may find a $0 \neq f(x_1, x_2) \in k_0\{x_1, x_2\}$ among all f_σ , such that

$$(*) \quad f(1 + \lambda_1 a, 1 + \lambda_2 b) = 0 \quad \text{in } D(\lambda_2)(\lambda_1).$$

Now we embed $D(\lambda_1)(\lambda_2)$ in the Laurent series division ring $D((\lambda_2))((\lambda_1))$. We obtain

$$0 \neq \psi(f(x_1, x_2)) = \lambda_1^{m_1} \lambda_2^{m_2} f_{m_1, m_2}(y_1, y_2) + R,$$

with $f_{m_1, m_2} \neq 0$, and where all the terms in R possess higher (λ_1, λ_2) -degree (in the lexicographic order, say). But from $(*)$ $f_{m_1, m_2}(a, b) = 0$. This contradicts the hypothesis that a and b generate a free k_0 -algebra. The result follows. $\square$

REFERENCES

1. J. Z. Goncalves and M. Shirvani, *On free subalgebras in division algebras and related rings*, preprint.
2. A. I. Lichtman, *On subgroups of the multiplicative group of skew fields*, Proc. Amer. Math. Soc. **63** (1977), 15–16. MR **56**:5744
3. ———, *Free subgroups of normal subgroups of the multiplicative group of skew fields*, Proc. Amer. Math. Soc. **71** (1978), 174–178. MR **58**:779
4. ———, *On normal subgroups of the multiplicative group of skew fields generated by a polycyclic-by-finite group*, J. Algebra **78** (1982), 548–577. MR **84f**:16016
5. M. Lorenz, *On free subalgebras of certain division algebras*, Proc. Amer. Math. Soc. **98** (1986), 401–405. MR **87m**:16040
6. L. Makar-Limanov, *The skew field of fractions of the Weyl algebra contains a free noncommutative subalgebra*, Comm. Algebra **17** (1983), 2003–2006. MR **84j**:16012
7. ———, *On free subobjects of skew fields*, Methods in Ring Theory (F. van Oystaeyen, ed.), Reidel, Dordrecht, 1984, pp. 281–285. CMP 17:06
8. ———, *On group rings of nilpotent groups*, Israel J. Math. **48** (1984), 245–248. MR **86c**:16006
9. L. Makar-Limanov and P. Malcolmson, *Words periodic over the center of a division ring*, Proc. Amer. Math. Soc. **93** (1985), 590–592. MR **86g**:16026
10. L. Makar-Limanov, *On subnormal subgroups of skew fields*, J. Algebra **114** (1988), 261–267. MR **89d**:16024
11. ———, *On subalgebras of the first Weyl algebra*, Comm. Algebra **19** (1991), 1971–1982. MR **92g**:16049
12. L. Makar-Limanov and P. Malcolmson, *Free subalgebras of enveloping algebras*, Proc. Amer. Math. Soc. **11** (1991), 315–322. MR **91f**:16023

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SAO PAULO, CP 66281, 05389-970 SAO PAULO SP, BRAZIL

E-mail address: jzg@ime.usp.br

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA T6G 2G1

E-mail address: mazi@schur.math.ualberta.ca