A WHITNEY-STRATIFIED CURVE IN $\mathbf{R}^3$ WITH MULTIPLE-POINT PROJECTIONS

MICHAŁ KWIECIŃSKI AND LAURENT NOIREL

(Communicated by James E. West)

ABSTRACT. We construct a compact $\mathcal{C}^\infty$ Whitney-stratified curve in $\mathbf{R}^3$, such that each of its plane projections has an infinite number of multiple points.

1. INTRODUCTION

Almost all hyperplane projections of a stratified set of large codimension in $\mathbf{R}^n$ are stratified embeddings. In [1], the authors showed that in general the projections do not preserve Whitney regularity, answering a question of Thom [2]. The aim of this paper is to show that in small codimensions, the image under a hyperplane projection of a $\mathcal{C}^\infty$ Whitney-stratified set is in general not stratifiable (i.e. cannot be partitioned into a locally finite union of submanifolds).

Let us first specify some terminology and notation. We say that a point M is a *multiple point* of a $\mathcal{C}^\infty$ curve $\psi : I \rightarrow \mathbf{R}^n$ (with I an interval of $\mathbf{R}$) if there exist $t_1, t_2 \in I$, $t_1 \neq t_2$ such that $\psi(t_1) = \psi(t_2) = M$ and $\psi'(t_1), \psi'(t_2)$ are not colinear. For a point x in the two-dimensional sphere $\mathbf{S}^2 \subset \mathbf{R}^3$, we denote by $\pi_x : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ the orthogonal projection of $\mathbf{R}^3$ with kernel $\mathbf{R}x$, followed by the inclusion of the hyperplane $x^\perp$ (orthogonal to x) in $\mathbf{R}^3$. We denote by $\|\cdot\|$ the Euclidean norm on $\mathbf{R}^3$. The main result of the paper is the following

Example. We will construct a $\mathcal{C}^\infty$ curve $f : [0, +\infty[\rightarrow \mathbf{R}^3$, such that

1. $f'(t) \neq 0$ for all $t \in \mathbf{R}$,
2. $\|f(t)\|$ is strictly decreasing and $f(t) \rightarrow 0$, $t \rightarrow +\infty$,
3. $(\frac{f'(t)}{\|f'(t)\|} + \frac{f(t)}{\|f(t)\|}) \rightarrow 0$, $t \rightarrow +\infty$,
4. for *any* $x \in \mathbf{S}^2$, the curve $\pi_x \circ f$ has a sequence of multiple points (different from 0) converging to 0.

The first two conditions imply that $\{0\}, \{f(0)\}, f([0, \infty[)$ is a stratification of the set $S = \overline{f([0, \infty[)}$. The third condition means precisely that the above stratification is Whitney b -regular at 0 (locally, near $f(0)$, the set S is a manifold with boundary). The fourth condition means that no hyperplane projection of the set S is stratifiable. Such a phenomenon does not occur for hyperplane projections of stratified curves in Euclidean n -space for $n \geq 4$ (see [1]), nor does it occur for smooth curves in $\mathbf{R}^3$. Indeed, almost all hyperplane projections of a $\mathcal{C}^\infty$ compact curve in $\mathbf{R}^3$ are injective outside a finite set of points. This is easily proved using

Received by the editors October 1, 1994.

1991 *Mathematics Subject Classification.* Primary 57N80; Secondary 53A04.

Sard's theorem. More generally, if the set S is subanalytic, then all projections are stratifiable.

Our example will be a *rapid spiral*, i.e. a curve of the type $f(t) = e^{-h(t)}\phi(t)$. Here $h : [0, +\infty[\rightarrow \mathbf{R}$ is a $\mathcal{C}^\infty$ function, with $h'(t) > 0$ for all t and $h'(t) \rightarrow +\infty$, $t \rightarrow +\infty$, whereas $\phi : [0, +\infty[\rightarrow \mathbf{S}^2$ is a $\mathcal{C}^\infty$ curve, such that $\|\phi'(t)\| \leq 1$ for all t . We will say that the rapid spiral f has *base* ϕ and *speed* h . It is easily seen that f satisfies the first three conditions required for the example, given any base and speed. A particular choice of ϕ and h will ensure that the fourth condition is also fulfilled.

2. A CURVE WITH A MULTIPLE POINT

Let $N = (0, 0, 1)$ be the “North Pole” of the sphere $\mathbf{S}^2$ in $\mathbf{R}^3$. Choose any $\mathcal{C}^\infty$ map $\psi : [0, 1] \rightarrow \mathbf{S}^2$, such that

1. $\|\psi'(t)\| \leq 1$, for all $t \in [0, 1]$,
2. $\psi(\frac{1}{3}) = \psi(\frac{2}{3}) = N$,
3. $\psi'(\frac{1}{3})$ and $\psi'(\frac{2}{3})$ are not colinear.

We have the following

Property (α). For any $k \in \mathbf{R}$, $k > 0$, there exists a neighbourhood U^k of N in $\mathbf{S}^2$, such that for all $x \in U^k - \{N\}$, the curve $\tilde{\psi}_x : [0, 1] \rightarrow x^\perp \hookrightarrow \mathbf{R}^3$; $\tilde{\psi}_x(t) = \pi_x(e^{-kt}\psi(t))$ has a multiple point different from 0.

Sketch of proof. For any $x \in (\mathbf{S}^2 - \mathbf{R}^2 \times \{0\})$, let $\hat{\pi}_x$ be the projection with kernel $\mathbf{R}x$, on the coordinate hyperplane $\mathbf{R}^2 \times \{0\}$. Let $\hat{\psi}_x : [0, 1] \rightarrow \mathbf{R}^2 \times \{0\}$ be the curve defined by $\hat{\psi}_x(t) = \hat{\pi}_x(e^{-kt}\psi(t))$. Of course it is equivalent to show Property (α) with $\tilde{\psi}_x$ replaced by $\hat{\psi}_x$.

That, in turn, will follow automatically from the existence of local diffeomorphisms δ and μ from a neighbourhood of 0 in $\mathbf{R}^2$, to respectively $\mathbf{S}^2$ and $\mathbf{R}^2 \times \{0\}$, with $\delta(0) = N$ and $\mu(0) = 0$, such that $\mu(s, t)$ is a multiple point of the curve $\hat{\psi}_{\delta(s, t)}$. To construct δ and μ , set

$$P(s) = e^{-k(s+\frac{1}{3})}\psi\left(s + \frac{1}{3}\right) \quad \text{and} \quad Q(t) = e^{-k(t+\frac{2}{3})}\psi\left(t + \frac{2}{3}\right).$$

Then define

$$\delta(s, t) = \frac{P(s) - Q(t)}{\|P(s) - Q(t)\|} \quad \text{and} \quad \mu(s, t) = \frac{Q_3(t)\pi_N(P(s)) - P_3(s)\pi_N(Q(t))}{Q_3(t) - P_3(s)}.$$

Here P_i and Q_i , are the i th coordinates of P and Q . One calculates:

$$\begin{aligned} (1 - e^{-\frac{k}{3}})\frac{\partial \delta}{\partial s}(0, 0) &= (1 - e^{-\frac{k}{3}})\frac{\partial \mu}{\partial s}(0, 0) = \psi'\left(\frac{1}{3}\right), \\ (1 - e^{-\frac{k}{3}})\frac{\partial \delta}{\partial t}(0, 0) &= (1 - e^{-\frac{k}{3}})\frac{\partial \mu}{\partial t}(0, 0) = \psi'\left(\frac{2}{3}\right), \end{aligned}$$

which shows that both mappings are local diffeomorphisms. Furthermore $\mu(s, t) = \hat{\psi}_{\delta(s, t)}(s + \frac{1}{3}) = \hat{\psi}_{\delta(s, t)}(t + \frac{2}{3})$ is a multiple point of $\hat{\psi}_{\delta(s, t)}$ for (s, t) sufficiently close to 0. $\square$

3. THE CONSTRUCTION

We are now ready to construct ϕ and h . For each point $y \in \mathbf{S}^2$, choose a rotation $\mathcal{R}_y$ of $\mathbf{R}^3$ around 0 which sends N to y . For every $k = 1, 2, \dots$, choose $y_1^k, \dots, y_{n_k}^k \in \mathbf{S}^2$ such that the punctured open sets $\mathcal{R}_{y_1^k}(U^k - \{N\}), \dots, \mathcal{R}_{y_{n_k}^k}(U^k - \{N\})$ cover $\mathbf{S}^2$. Now ϕ can be chosen as any curve joining all the curves $\mathcal{R}_{y_i^k} \circ \psi$. To be precise, we take ϕ as any $\mathcal{C}^\infty$ map for which there exist, for each $k = 1, 2, \dots$, sequences of reals $T_1^k, \dots, T_{n_k}^k$, with $T_i^k + 1 < T_{i+1}^k$, for all k, i and $T_{n_k}^k + 1 < T_1^{k+1}$ for all k , and such that $\phi(t) = \mathcal{R}_{y_i^k}(\psi(t - T_i^k))$ for $t \in [T_i^k, T_i^k + 1]$, for all k, i . Furthermore, we require that $\|\phi'(t)\| \leq 1$ for all t . Such a curve is easily obtained by standard smoothing techniques.

As h , we can take any $\mathcal{C}^\infty$ function, with weakly increasing positive derivative, for which there exists a sequence of reals $\{t_k\}_{k=1}^\infty$, such that $h(t) = kt - t_k$ for $t \in [T_1^k, T_{n_k}^k]$, for all k . Such a function is also easily constructed.

We now show that the rapid spiral f with base ϕ and speed h satisfies property 4 of the example. Indeed, let $x \in \mathbf{S}^2$. For each k , choose $i_k \in \{1, \dots, n_k\}$, such that $x \in \mathcal{R}_{y_{i_k}^k}(U^k - \{N\})$. Denote $\mathcal{R} = \mathcal{R}_{y_{i_k}^k}$. On $[T_{i_k}^k, T_{i_k}^k + 1]$ we have

$$\pi_x(f(t)) = \pi_x(e^{-kt+t_k}\mathcal{R}(\psi(t - T_{i_k}^k))) = e^{(t_k - kT_{i_k}^k)}\mathcal{R}(\tilde{\psi}_{\mathcal{R}^{-1}(x)}(t - T_{i_k}^k)).$$

Since $\mathcal{R}^{-1}(x) \in U^k - \{N\}$, by Property (α), the curve $\tilde{\psi}_{\mathcal{R}^{-1}(x)}$ has a multiple point different from 0 and thus, by the last equality, so has $\pi_x \circ f|_{[T_{i_k}^k, T_{i_k}^k + 1]}$. Since the above holds for any k , this means that for any $T > 0$, the curve $\pi_x \circ f$ has a multiple point $\pi_x \circ f(t)$, with $t > T$. Hence, $\pi_x \circ f$ has multiple points different from 0, arbitrarily close to 0. This finishes the proof that our rapid spiral satisfies the fourth condition of the example.

REFERENCES

1. M. Kwieciński and L. Noirel, *Sur une question de René Thom à propos des projections d'une stratification de Whitney*, C. R. Acad. Sci. Paris Sér. I **318** (1994), 149–152. MR **95c**:58010
2. R. Thom, *Quid des stratifications canoniques*, in *Singularities*, Proc. Lille 1991 (J. P. Brasselet, ed.), London Math. Soc. Lecture Note Series, No. 201, 1994. CMP 95:01

(on leave from) UNIwersytet Jagielloński, Instytut Matematyki, ul. Reymonta 4, 30-059 KRAKÓW, POLAND

E-mail address: kwiecins@im.uj.edu.pl

Current address: Equipe CNRS SiGmA, CIRM Luminy Case 916, 13288 Marseille Cedex 9, France

E-mail address: michal@cirm5.univ-mrs.fr

URA 225 du CNRS, UFR MIM, Université de Provence, 3 Place Victor Hugo, 13331 MARSEILLE CEDEX 3, FRANCE

E-mail address: noirel@gyptis.univ-mrs.fr