COMPOSITION OF BLOCHS WITH BOUNDED ANALYTIC FUNCTIONS

E. G. KWON

(Communicated by Theodore Gamelin)

ABSTRACT. If f is a holomorphic self-map of the open unit disc and $1 \leq p < \infty$, then the following are equivalent.

(1) $h \circ f \in H^{2p}$ for all Bloch functions h .

$$(2) \quad \sup_r \int_0^{2\pi} \left(\log \frac{1}{1 - |f(re^{i\theta})|^2} \right)^p d\theta < \infty.$$

$$(3) \quad \int_0^{2\pi} \left(\int_0^1 (f^\#)^2(re^{i\theta})(1-r)dr \right)^p d\theta < \infty,$$

where $f^\#$ is the hyperbolic derivative of f : $f^\# = |f'|/(1 - |f|^2)$.

1. INTRODUCTION

Initiated by works of P. Ahern and W. Rudin ([A], [AR]), there is extensive research on Bloch-to-BMOA pullbacks, that is, research on those holomorphic maps f of the unit ball of $\mathbf{C}^n$ into the unit disc of $\mathbf{C}$ for which the composition operator defined by

$$\mathcal{C}_f(h) = h \circ f$$

takes Bloch functions to functions of BMOA. We refer to [RU] for recent research on Bloch to BMOA pullbacks.

It is known (see [RU], Section 5), when $n = 1$, that one of the necessary and sufficient conditions for $\mathcal{C}_f$ to take all Blochs to BMOA is that f be a function of BMOA_σ , the Yamashita hyperbolic BMOA class (see [G] and [Y3] for BMOA and BMOA_σ). This note is to resolve a natural parallelism to the phenomenon.

Theorem 1 (Main Result). *If f is a holomorphic self-map of the open unit disc and $1 \leq p < \infty$, then the following are equivalent.*

(1) $\mathcal{C}_f$ takes Blochs to H^{2p} , that is,

$$h \circ f \in H^{2p} \quad \text{for all Bloch functions } h.$$

Received by the editors January 31, 1994 and, in revised form, October 19, 1994.

1991 *Mathematics Subject Classification*. Primary 30D55, 30D45.

Key words and phrases. H^p space, Bloch space, hyperbolic Hardy class, pullbacks.

This paper was supported by NON DIRECTED RESEARCH FUND, Korea Research Foundation, 1993.

(2) f belongs to Yamashita's hyperbolic Hardy class H_σ^p , that is,

$$\sup_r \int_0^{2\pi} \left(\log \frac{1}{1 - |f(re^{i\theta})|^2} \right)^p d\theta < \infty.$$

$$(3) \quad \int_0^{2\pi} \left(\int_0^1 (f^\#)^2(re^{i\theta})(1-r)dr \right)^p d\theta < \infty,$$

where $f^\#$ is the hyperbolic derivative of f : $f^\# = |f'|/(1 - |f|^2)$.

2. PRELIMINARIES

The Bloch space $\mathcal{B}$ consists of those f holomorphic in the open unit disc D of the complex plane for which

$$\|f\|_{\mathcal{B}} := \sup_{z \in D} |f'(z)|(1 - |z|^2) < \infty.$$

We let $1 \leq p < \infty$ and set for f subharmonic in D

$$\|f\|_p := \sup_r \left(\int_0^{2\pi} |f(re^{i\theta})|^p \frac{d\theta}{2\pi} \right)^{1/p}.$$

Then $H^p = H^p(D)$ consists of those f holomorphic in D for which $\|f\|_p < \infty$. See [D] and [G] for Bloch and H^p spaces.

The Yamashita hyperbolic Hardy class H_σ^p is defined as the set of those holomorphic self-maps f of D for which

$$\|\sigma(f)\|_p < \infty,$$

where $\sigma(z)$ denotes the hyperbolic distance of z and 0 in D , namely,

$$\sigma(z) = \frac{1}{2} \log \frac{1 + |z|}{1 - |z|}.$$

Though H_σ^p is not a linear space, it has, as hyperbolic counterparts, many properties analogous to those of H^p . We let T be the boundary of D and set, following Yamashita,

$$\lambda(f) = \log \frac{1}{1 - |f|^2} \quad \text{and} \quad f^\# = \frac{|f'|}{1 - |f|^2}$$

for the holomorphic self-map f of D . Then $\sigma(f)^p$, $\lambda(f)^p$, and $(f^\#)^p$ are subharmonic functions, so that their integral means over rT are nondecreasing functions of r : for example,

$$\int_0^{2\pi} \lambda(f)^p(re^{i\theta}) \frac{d\theta}{2\pi} \nearrow \|\lambda(f)\|_p^p \quad \text{as } r \nearrow 1.$$

Also, there are corresponding maximal theorems for these functions: Set

$$M_\lambda(f, \theta) = \sup \{ \lambda(f)(re^{i\theta}) : 0 \leq r < 1 \};$$

then

$$(2.1) \quad \|M_\lambda(f, \cdot)\|_{L^p} \leq C_p \|\lambda(f)\|_p$$

for $f \in H_\sigma^p$ ([Y1]). The left side of (2.1) means usual $L^p(T)$ norm. The function $f^\#$ is the hyperbolic counterpart of f' and it easily follows that

$$(2.2) \quad \frac{1}{2}\lambda(f) \leq \sigma(f) < \frac{1}{2}\lambda(f) + \log 2$$

and

$$(2.3) \quad \Delta(\lambda(f)^p) = 4p\{(p-1)|f|^2 + \lambda(f)\}\lambda(f)^{p-2}(f^\#)^2,$$

where Δ denotes the Laplacian:

$$\Delta = 4 \frac{\partial^2}{\partial z \partial \bar{z}}.$$

From (2.2) and (2.3), it should be noted that

$$f \in H_\sigma^p \quad \text{if and only if} \quad \|\lambda(f)\|_p < \infty$$

and

$$(2.4) \quad \Delta(\lambda(f)^p) \sim \lambda(f)^{p-1}(f^\#)^2.$$

Here and after $\psi \sim \phi$ means that either both sides are zero or the quotient ψ/ϕ lies between two positive uniform constants. See, for example, [K1], [K2], [Y1], and [Y2] for the theory of H_σ^p .

In the remaining sections, we confine f to being a holomorphic self-map of D and denote f_r , $0 \leq r \leq 1$, as the function defined by $f_r(z) = f(rz)$, $z \in D$. Positive constants depending on p (or q) will be denoted by C_p (or C_q), whose quantities may vary at each occurrence.

3. HYPERBOLIC g -FUNCTION

For h holomorphic in D , g -function of Paley defined by

$$(3.1) \quad g(\theta) := g(h)(\theta) = \left(\int_0^1 |h'|^2(re^{i\theta})(1-r)dr \right)^{1/2}, \quad 0 \leq \theta < 2\pi,$$

satisfies $\|g(h)\|_{L^p} \sim \|h\|_p$ if $h(0) = 0$ ([Z]). Consider Green's theorem of the form

$$r \int_0^{2\pi} \frac{\partial \psi}{\partial r} d\theta = \int \int_{|z| \leq r} \Delta \psi \, dx dy,$$

valid for $\psi \in C^2(D)$. If we integrate both sides with respect to dr after dividing them by r and applying $\psi = \lambda(f)^p$, then we obtain, by use of (2.4),

$$(3.2) \quad \begin{aligned} & \|\lambda(f_\rho)\|_p^p - \lambda(f)^p(0) \\ & \sim \frac{1}{2\pi} \int_0^\rho \frac{dr}{r} \int \int_{|z| \leq r} \lambda(f)^{p-1}(f^\#)^2(z) \, dx dy \\ & = \frac{1}{2\pi} \int \int_{|z| < \rho} \lambda(f)^{p-1}(f^\#)^2(z) \log \frac{\rho}{|z|} \, dx dy, \quad 0 \leq \rho \leq 1. \end{aligned}$$

In particular, we see from (3.2) that $f \in H_\sigma^1$ if and only if

$$\infty > \int \int_D (f^\#)^2(z) \log \frac{1}{|z|} dx dy.$$

This suggests we define the hyperbolic version of g -function using $f^\#$. We define

$$(3.3) \quad g_\sigma(\theta) := g_\sigma(f)(\theta) = \int_0^1 (f^\#)^2(re^{i\theta}) (1-r) dr, \quad 0 \leq \theta < 2\pi.$$

It is not surprising to see the absence of the square root in the definition of g_σ in (3.3) when we compare it to that of g -function in (3.1), because there is a known parallelism (see [Y2]) between H^2 and H_σ^1 .

Theorem 2. *If $1 \leq p < \infty$, then the following are equivalent.*

(1) $f \in H_\sigma^p$.

(2) $g_\sigma(f) \in L^p(T)$.

In fact, $\|\lambda(f)\|_p \sim \|g_\sigma(f)\|_{L^p}$ provided $f(0) = 0$.

Proof. By (3.2) and (3.3), there is nothing to prove when $p = 1$. We assume $1 < p < \infty$, and let $\frac{1}{p} + \frac{1}{q} = 1$.

(1) $\implies$ (2) We begin with the identity

$$\|g_\sigma\|_{L^p} = \sup \int_0^{2\pi} g_\sigma(\theta) h(\theta) \frac{d\theta}{2\pi},$$

where the supremum is taken with respect to all nonnegative trigonometric polynomials h with $\|h\|_{L^q} \leq 1$. Since $(f^\#)^2$ is subharmonic, we have

$$(3.4) \quad (f^\#)^2(r^2 e^{i\theta}) \leq \int_0^{2\pi} P(r, \theta - t) (f^\#)^2(re^{it}) \frac{dt}{2\pi}, \quad 0 \leq r < 1,$$

where $P(r, \theta)$ is the Poisson kernel:

$$P(r, \theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}.$$

Let u be the Poisson integral of h . Then

$$\begin{aligned} & \int_0^{2\pi} g_\sigma(\theta) h(\theta) d\theta \\ &= \int_0^{2\pi} \int_0^1 (f^\#)^2(re^{i\theta}) h(\theta) (1-r) dr d\theta \\ (3.5) \quad &= \int_0^{2\pi} \int_0^1 (f^\#)^2(r^2 e^{i\theta}) h(\theta) (1-r^2) 2r dr d\theta \\ &\leq 2 \int \int_D (f^\#)^2(z) u(z) (1-|z|^2) dx dy \\ &\leq 4 \int \int_D (f^\#)^2(z) u(z) \log \frac{1}{|z|} dx dy, \end{aligned}$$

where we changed the order of integration and used (3.4) in the first inequality.

If we denote $\partial = \frac{1}{2}(\frac{\partial}{\partial x} - i\frac{\partial}{\partial y})$ and $\bar{\partial} = \frac{1}{2}(\frac{\partial}{\partial x} + i\frac{\partial}{\partial y})$, $z = x + iy$, then it follows from elementary calculation and (2.3) that

$$4(f^\#)^2 u = \Delta(\lambda(f)u) - 4(\partial\lambda(f)\bar{\partial}u + \bar{\partial}\lambda(f)\partial u),$$

so that by (3.5) we have

$$\begin{aligned} & \int_0^{2\pi} g_\sigma(\theta)h(\theta)d\theta \\ (3.6) \quad & \leq \left| \int \int_D \Delta(\lambda(f)u)(z) \log \frac{1}{|z|} dx dy \right| \\ & \quad + 4 \int \int_D |\partial\lambda(f)\bar{\partial}u + \bar{\partial}\lambda(f)\partial u|(z) \log \frac{1}{|z|} dx dy \\ & = (I) + (II). \end{aligned}$$

Now, using Green's theorem (as in (3.2) with $p = 1$) with limiting process and Hölder's inequality, we obtain

$$\begin{aligned} (3.7) \quad (I) &= \left| \lim_{\rho \rightarrow 1} \int_0^{2\pi} (\lambda(f)u)(\rho e^{i\theta}) d\theta - 2\pi(\lambda(f)u)(0) \right| \\ &\leq 2\pi \|\lambda(f)\|_p \|u\|_q \leq 2\pi \|\lambda(f)\|_p. \end{aligned}$$

On the other hand, if we let ϕ be a holomorphic function in D whose real part is u , it then follows from direct differentiation that

$$|\partial\lambda(f)| = |\bar{\partial}\lambda(f)| = |f|f^\#$$

and

$$|\partial u| = |\bar{\partial}u| = \frac{1}{2}|\phi'|.$$

Hence

$$\begin{aligned} (II) &\leq 4 \int \int_D |\phi'(z)| |f(z)| f^\#(z) \log \frac{1}{|z|} dx dy \\ &\leq 4 \int_0^{2\pi} \int_0^1 |\phi'(re^{i\theta})| |f(re^{i\theta})| f^\#(re^{i\theta}) (1-r) dr d\theta. \end{aligned}$$

Since $|f(re^{i\theta})| \leq \sqrt{\lambda(f)(re^{i\theta})} \leq M_\lambda^{1/2}(f, \theta)$, we have, by the Schwarz inequality,

$$(3.8) \quad (II) \leq 4 \int_0^{2\pi} M_\lambda^{1/2}(f, \theta) \sqrt{g_\sigma(\theta)} g_\phi(\theta) d\theta,$$

where $g_\phi(\theta)$ is Paley g -function of ϕ . Applying Hölder's inequality with indices $2p$, $2p$, q to the right side of (3.8) and using maximal theorem (2.1), we arrive at

$$(II) \leq C_p \|\lambda(f)\|_p^{1/2} \|g_\sigma(f)\|_{L^p}^{1/2} \|g_\phi\|_{L^q}.$$

From the theory of g -function, we know $\|g_\phi\|_{L^q} \leq C_q \|\phi\|_q$, and it follows from the theorem of M. Riesz ([Z]) that $\|\phi\|_q \leq C_q \|u\|_q \leq C_q$. Thus

$$(3.9) \quad (II) \leq C_p \|\lambda(f)\|_p^{1/2} \|g_\sigma(f)\|_{L^p}^{1/2}.$$

Hence, combining estimates (3.6), (3.7), and (3.9), we have

$$\begin{aligned} \int_0^{2\pi} g_\sigma(\theta) h(\theta) d\theta &\leq (I) + (II) \\ &\leq 2\pi \|\lambda(f)\|_p + C_p \|\lambda(f)\|_p^{1/2} \|g_\sigma(f)\|_{L^p}^{1/2}, \end{aligned}$$

for all positive trigonometric polynomials h with $\|h\|_q \leq 1$. Therefore we obtain

$$(3.10) \quad \|g_\sigma(f)\|_{L^p} \leq \|\lambda(f)\|_p + C_p \|\lambda(f)\|_p^{1/2} \|g_\sigma(f)\|_{L^p}^{1/2}, \quad f \in H_\sigma^p.$$

Now we conclude from (3.10) that

$$(3.11) \quad \|g_\sigma(f)\|_{L^p} \leq C_p \|\lambda(f)\|_p.$$

In fact, if $f \equiv 0$, then there is nothing to prove; otherwise, setting $X(r) = X(f, p, r) = (\|g_\sigma(f_r)\|_{L^p} / \|\lambda(f_r)\|_p)^{\frac{1}{2}}$, $0 < r < 1$, (3.10) with f_r in place of f becomes

$$X^2(r) \leq 1 + C_p X(r),$$

and this means, by comparing the order of $X(r)$, that $X(r)$, $0 < r < 1$, does not exceed the larger root of the equation $X^2 = 1 + C_p X$. This proves (3.11) with f_r , $0 < r < 1$, in place of f , and so (3.11) follows by the monotonicity of both sides.

(2) $\implies$ (1) It follows from (3.2) that

$$\begin{aligned} \|\lambda(f_r)\|_p^p - \lambda(f)^p(0) &\sim C_p \int_0^{2\pi} \frac{d\theta}{2\pi} \int_0^r (f^\#)^2 \lambda(f)^{p-1} (\rho e^{i\theta}) \log \frac{r}{\rho} \rho d\rho \\ &\leq C_p \int_0^{2\pi} M_\lambda^{p-1}(f_r, \theta) g_\sigma(f)(\theta) \frac{d\theta}{2\pi}. \end{aligned}$$

Hölder's inequality and the maximal theorem show the last quantity to be bounded by

$$C_p \|\lambda(f_r)\|_p^{p-1} \|g_\sigma(f)\|_{L^p},$$

so that we have

$$(3.12) \quad \|\lambda(f_r)\|_p^p - \lambda(f)^p(0) \leq C_p \|\lambda(f_r)\|_p^{p-1} \|g_\sigma(f)\|_{L^p}.$$

If $g_\sigma \equiv 0$, there remains nothing to prove. Otherwise, under the assumption $0 < \|g_\sigma\|_{L^p} < \infty$, by considering the order of

$$Y(r) = Y(f, p, r) = (\|\lambda(f_r)\|_p / \|g_\sigma(f_r)\|_{L^p}), \quad 0 < r < 1,$$

we conclude that $Y(r)$, $0 < r < 1$, does not exceed the largest root of the equation

$$Y^p - \frac{\lambda(f)^p(0)}{\|g_\sigma(f)\|_{L^p}^p} = C_p Y^{p-1},$$

and from this follows

$$\|\lambda(f)\|_p \leq C_p(f) \|g_\sigma(f)\|_{L^p}.$$

Here, $C_p(f)$ denotes a constant depending on p and f .

The last assertion of Theorem 2 follows from (3.11) and (3.12).

4. BLOCH TO H^p PULLBACKS

Now we prove our main theorem, Theorem 1. By the aid of Theorem 2, it suffices to prove the following.

Theorem 3. *If $1 \leq p < \infty$, then the following are equivalent.*

- (1) $g_\sigma(f) \in L^p(T)$.
- (2) $\mathcal{C}_f$ takes Blochs to H^{2p} .

Proof. (1) $\implies$ (2) Let $h \in \mathcal{B}$. Then

$$(4.1) \quad |(\mathcal{C}_f h)'| = |(h \circ f)'| \leq \|h\|_{\mathcal{B}} f^\#.$$

Hence

$$(4.2) \quad \begin{aligned} & \|g_{h \circ f}\|_{L^{2p}}^{2p} \\ &= \int_0^{2\pi} \left(\int_0^1 |(h \circ f)'(re^{i\theta})|^2 (1-r) dr \right)^p \frac{d\theta}{2\pi} \\ &\leq \|h\|_{\mathcal{B}}^{2p} \int_0^{2\pi} \left(\int_0^1 (f^\#)^2(re^{i\theta}) (1-r) dr \right)^p \frac{d\theta}{2\pi} \\ &= \|h\|_{\mathcal{B}}^{2p} \|g_\sigma(f)\|_{L^p}^p. \end{aligned}$$

Therefore $\|h \circ f\|_{2p} < \infty$ if $g_\sigma(f) \in L^p(T)$.

(2) $\implies$ (1) Using g -function, (2) says that

$$(4.3) \quad \int_0^{2\pi} \left(\int_0^1 |(h \circ f)'|^2(re^{i\theta}) (1-r) dr \right)^p d\theta < \infty \quad \text{if} \quad h \in \mathcal{B}.$$

On the other hand, W. Ramey and D. Ullrich ([RU], Proposition 5.4) constructed two Bloch functions $h_j, j = 1, 2$, such that

$$(4.4) \quad (1 - |z|^2)(|h'_1(z)| + |h'_2(z)|) \geq 1, \quad z \in D.$$

From (4.4) it follows that $(|h'_1 \circ f| + |h'_2 \circ f|) \geq (1 - |f|^2)^{-1}$, so that

$$(4.5) \quad \begin{aligned} & \left(\int_0^1 |(h_1 \circ f)'|^2 (1-r) dr \right)^p + \left(\int_0^1 |(h_2 \circ f)'|^2 (1-r) dr \right)^p \\ &\geq 2^{-2p} \left(\int_0^1 |f'|^2 (|h'_1 \circ f| + |h'_2 \circ f|)^2 (1-r) dr \right)^p \\ &\geq 2^{-2p} \left(\int_0^1 \frac{|f'|^2}{(1 - |f|^2)^2} (1-r) dr \right)^p \\ &= 2^{-2p} g_\sigma^p(f). \end{aligned}$$

Now, integrating (4.5) with respect to $d\theta$ and applying (4.3) with $h_j, j = 1, 2$, in place of h , we obtain

$$\|g_\sigma(f)\|_p \leq C_p (\|h_1 \circ f\|_{2p} + \|h_2 \circ f\|_{2p}).$$

This completes the proof.

5. EXTREME POINTS OF H^∞

H^∞ denotes, as usual, the space of bounded holomorphic functions on D . A well-known theorem of deLeeuw and Rudin ([D], Theorem 7.9) says that

$$\int_0^{2\pi} \log \frac{1}{1 - |f(e^{i\theta})|^2} d\theta = \infty$$

is necessary and sufficient for a holomorphic f with $\|f\|_\infty = \sup_{z \in D} |f(z)| = 1$ to be an extreme point of the closed unit ball of H^∞ . The following is a direct corollary of Theorem 1.

Corollary 4. *Let $f \in H^\infty$, $\|f\|_\infty = 1$. Then the following are equivalent.*

- (1) *f is an extreme point of the closed unit ball of H^∞ .*
- (2) *$h \circ f \notin H^2$ for some $h \in \mathcal{B}$.*

ACKNOWLEDGMENT

The author wishes to express his sincere gratitude to the referee for his helpful suggestion.

REFERENCES

- [A] Patrick R. Ahern, *On the behavior near torus of functions holomorphic in the ball*, Pacific J. Math. **107** (1983), 267–278. MR **84i**:32023
- [AR] Patrick R. Ahern and Walter Rudin, *Bloch functions, BMO, and boundary zeros*, Indiana Univ. Math. J. **36** (1987), 131–148. MR **88d**:42036
- [D] Peter. L. Duren, *The theory of H^p functions*, Academic Press, New York, 1970. MR **42**:3552
- [G] John. B. Garnett, *Bounded analytic functions*, Academic Press, New York, 1981. MR **83g**:30037
- [K1] E. G. Kwon, *Fractional integration and the hyperbolic derivative*, Bull. Austral. Math. Soc. **38** (1988), 357–364. MR **90a**:30096
- [K2] ———, *Mean growth of the hyperbolic Hardy class functions*, Math. Japonica **35** (1990), 451–460. MR **91e**:30064
- [RU] Wade Ramey and David Ullrich, *Bounded mean oscillations of Bloch pullbacks*, Math. Ann. **291** (1991), 591–606. MR **92i**:32004
- [Y1] Shinji Yamashita, *Functions with H^p hyperbolic derivative*, Math. Scand. **13** (1983), 238–244. MR **85f**:30055
- [Y2] ———, *Hyperbolic Hardy classes and hyperbolically Dirichlet finite functions*, Hokkaido Math. J. **10** (1981), 709–722, Special Issue.
- [Y3] ———, *Holomorphic functions of hyperbolically bounded mean oscillations*, Bollentino U.M.I. **5-B** (1986), no. 6, 983–1000. MR **88e**:30092
- [Z] A. Zygmund, *Trigonometric series*, Cambridge Univ. Press, London, 1959. MR **21**:6498

DEPARTMENT OF MATHEMATICS-EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG 760-749, KOREA

E-mail address: egkwon@anu.andong.ac.kr