THE STRUCTURE OF MEASURABLE MAPPINGS WITH VALUES IN LOCALLY CONVEX SPACES

JUN KAWABE

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. The purpose of this paper is to show that a theorem of A. Wisniewski remains valid without the approximation property.

1. INTRODUCTION

Let X and Y be Hausdorff topological spaces. Denote by $\mathcal{B}(X)$ and $\mathcal{B}(Y)$ respectively the Borel σ -algebras on these spaces. Let μ be a finite Borel measure on X . By $\mathcal{B}_\mu(X)$ we denote the completion of the σ -algebra $\mathcal{B}(X)$ with respect to the measure μ . A mapping f from X into Y is said to be μ -measurable if it is measurable with respect to $(\mathcal{B}_\mu(X), \mathcal{B}(Y))$. Recall that a Suslin space is a Hausdorff topological space which is the continuous image of some Polish space. It is well known but not easy to prove that every finite Borel measure on a Suslin space is Radon, and hence it is tight (see, e.g., [4, Theorem 10 of Chapter II in Part I]).

In [5], Wisniewski extended a theorem of Gihman and Skorohod [2, p. 544] which clarifies the connection between continuous and μ -measurable mappings on Hilbert spaces. Namely, he proved that if μ is a finite Borel measure on a metric space X , and if Y is a separable Banach space *with the approximation property*, then every μ -measurable mapping f from X into Y is the limit of a sequence of continuous mappings with respect to μ -almost everywhere convergence.

In this paper, we show that the above result remains valid in the case that Y is a Suslin locally convex topological linear space which *does not* necessarily have the approximation property. Consequently it turns out that Theorem 2 of [5] remains valid without the approximation property.

2. MAIN RESULTS

Theorem 1. *Let μ be a finite Borel measure on a metric space X , and let Y be a locally convex Hausdorff topological linear space. Assume that every finite Borel measure on Y is tight (this is satisfied, for instance, if Y is Suslin). If f is a μ -measurable mapping from X into Y , then there exists a sequence $\{f_n\}$ of continuous mappings from X into Y such that $f_n \rightarrow f$ μ -a.e.*

Received by the editors October 25, 1994.

1991 *Mathematics Subject Classification.* Primary 28C15, 60B05; Secondary 28A20, 28C20, 60B11.

Key words and phrases. μ -measurable mappings, continuous mappings, Suslin spaces, Banach spaces, Fréchet spaces, nuclear spaces, locally convex spaces, approximation property.

Proof. To prove the theorem it suffices to show that for any $\varepsilon > 0$ and any closed neighborhood U of the origin of Y , there exists a continuous mapping $g : X \rightarrow Y$ such that

$$\mu \{x : f(x) - g(x) \notin U\} < \varepsilon.$$

Since Y is locally convex, we may assume that U is absorbent and absolutely convex (see, e.g., [3, p. 12]). Put $\nu(B) = \mu(f^{-1}(B))$ for every $B \in \mathcal{B}(Y)$. Then ν is a finite Borel measure on Y , and hence it is tight by the assumption. Thus we can find a compact subset K of Y such that $\nu(Y - K) < \varepsilon/2$. Put $K' = f^{-1}(K)$. Then we have

$$(1) \quad \mu(X - K') < \frac{\varepsilon}{2}.$$

Since K is compact, there exist $y_1, y_2, \dots, y_m \in K$ such that

$$K \subset \left(y_1 + \frac{1}{2}U\right) \cup \left(y_2 + \frac{1}{2}U\right) \cup \dots \cup \left(y_m + \frac{1}{2}U\right).$$

Consider sets $B_1 = y_1 + \frac{1}{2}U, \dots, B_i = \left\{\bigcup_{k=1}^{i-1} \left(y_k + \frac{1}{2}U\right)\right\}^c \cap \left(y_i + \frac{1}{2}U\right), \dots, B_m = \left\{\bigcup_{k=1}^{m-1} \left(y_k + \frac{1}{2}U\right)\right\}^c \cap \left(y_m + \frac{1}{2}U\right)$. Then the B_i 's are Borel subsets of Y with $K \subset \bigcup_{i=1}^m B_i$ and $B_i \cap B_j = \emptyset$ if $i \neq j$. Hence

$$f(x) - \sum_{i=1}^m 1_{B_i}(f(x))y_i \in \frac{1}{2}U \quad \text{for all } x \in K',$$

where 1_B denotes the indicator function of a set B . Consequently, using (1), we infer that

$$(2) \quad \mu \left\{x : f(x) - \sum_{i=1}^m 1_{B_i}(f(x))y_i \notin \frac{1}{2}U\right\} \leq \mu(X - K') \leq \frac{\varepsilon}{2}.$$

For every $i = 1, 2, \dots, m$, we put $f_i(x) = 1_{B_i}(f(x))$ for all $x \in X$. Then the f_i 's are μ -measurable mapping from X into the real line R . On the other hand, since U is absorbent, there exists $\lambda > 0$ such that $y_i \in \lambda U$ for all $i = 1, 2, \dots, m$. Therefore, in view of Theorem 1 of [5], for every $i = 1, 2, \dots, m$, we can find continuous mappings g_i from X into R such that

$$(3) \quad \mu \left\{x : |f_i(x) - g_i(x)| > \frac{1}{2m\lambda}\right\} < \frac{\varepsilon}{2m}.$$

Put $g(x) = \sum_{i=1}^m g_i(x)y_i$ for every $x \in X$. Then g is a continuous mapping from X into Y . Moreover, from (2) and (3), together with the fact that U is convex, we

have

$$\begin{aligned}
 \mu \{x : f(x) - g(x) \notin U\} &= \mu \left\{ x : f(x) - \sum_{i=1}^m g_i(x)y_i \notin U \right\} \\
 &\leq \mu \left\{ x : f(x) - \sum_{i=1}^m f_i(x)y_i \notin \frac{1}{2}U \right\} \\
 &\quad + \mu \left\{ x : \sum_{i=1}^m f_i(x)y_i - \sum_{i=1}^m g_i(x)y_i \notin \frac{1}{2}U \right\} \\
 &\leq \frac{\varepsilon}{2} + \sum_{i=1}^m \mu \left\{ x : |f_i(x) - g_i(x)| > \frac{1}{2m\lambda} \right\} \\
 &\leq \varepsilon/2 + m \cdot \varepsilon/(2m) = \varepsilon.
 \end{aligned}$$

This completes the proof of the theorem. $\square$

Remark. We can show that Theorem 1 also holds in the case that X is a normal space and μ is a finite Borel measure on X satisfying the condition that $\mu(B) = \sup \{\mu(F) : F \subset B \text{ and } F \text{ is a closed subset of } X\}$ for all $B \in \mathcal{B}(X)$.

Corollary. *In the following cases, Theorem 1 holds:*

- (a) Y is a separable Banach space.
- (b) Y is a separable Fréchet space.
- (c) Y is the strict inductive limit of an increasing sequence of separable Fréchet spaces.
- (d) Y is the strong dual of a nuclear Fréchet space.
- (e) Y is the strong dual of the strict inductive limit of an increasing sequence of nuclear Fréchet spaces.

Proof. (a)–(e) are Suslin (in fact, Lusin) locally convex topological linear spaces by [1, Theorem I.5.1]. $\square$

REFERENCES

- [1] X. Fernique, *Processus linéaires, processus généralisés* Ann. Inst. Fourier (Grenoble) **17** (1967), 1-92. MR **36**:4628
- [2] I. I. Gihman and A. V. Skorohod, *The theory of stochastic processes*. I, Springer-Verlag, Berlin, Heidelberg and New York, 1974. MR **49**:11603
- [3] A. P. Robertson and W. Robertson, *Topological vector spaces*, Cambridge Univ. Press, 1964. MR **28**:5318
- [4] L. Schwartz, *Radon measures on arbitrary topological spaces and cylindrical measures*, Oxford Univ. Press, 1973. MR **54**:14030
- [5] A. Wisniewski, *The structure of measurable mappings on metric spaces*, Proc. Amer. Math. Soc. **122** (1994), 147-150. MR **94k**:28006

DEPARTMENT OF MATHEMATICS, FACULTY OF ENGINEERING, SHINSHU UNIVERSITY, WAKASATO, NAGANO 380, JAPAN

E-mail address: jkawabe@gipwc.shinshu-u.ac.jp