MULTINOMIAL EXPANSIONS AND THE PYTHAGOREAN THEOREM

ALLAN FRYANT

(Communicated by J. Marshall Ash)

Dedicated to the memory of Professor Morris Marden

ABSTRACT. An addition formula for homogeneous polynomials is used to obtain a generalization of the Pythagorean theorem and a new view of the multinomial expansion.

Let $x = (x_1, x_2, \dots, x_n) \in \mathbf{R}^n$, and let P denote the set of all polynomials

$$p(x) = \sum_{j=1}^m a_j x_1^{\alpha_j^1} x_2^{\alpha_j^2} \dots x_n^{\alpha_j^n}, \quad a_j \in \mathbf{R}.$$

Letting $\partial/\partial x = (\partial/\partial x_1, \partial/\partial x_2, \dots, \partial/\partial x_n)$, we define on P the inner product

$$\begin{aligned} (p, q) &= p \left(\frac{\partial}{\partial x} \right) q(x) \Big|_{x=0} \\ (1) \quad &= \left(\sum_{j=1}^m a_j \frac{\partial^{\alpha_j^1}}{\partial x_1^{\alpha_j^1}} \frac{\partial^{\alpha_j^2}}{\partial x_2^{\alpha_j^2}} \dots \frac{\partial^{\alpha_j^n}}{\partial x_n^{\alpha_j^n}} \right) q(x) \Big|_{x=0}. \end{aligned}$$

Let $H_k \subset P$ denote the vector space of homogeneous polynomials of degree k . Note that the dimension of H_k is

$$\dim H_k = d_k = \frac{(n+k-1)!}{(n-1)!k!}.$$

We first obtain a simple analog of the Funk-Hecke theorem [1, p. 247] for homogeneous polynomials.

Theorem 1. *If $p(x)$ is a homogeneous polynomial of degree k , then*

$$((x_1 y_1 + x_2 y_2 + \dots + x_n y_n)^k, p(y)) = k! p(x).$$

Proof. Expanding the power,

$$\begin{aligned} (x_1 y_1 + \dots + x_n y_n)^k &= \sum_{\alpha} \frac{k!}{\alpha_1! \alpha_2! \dots \alpha_n!} (x_1 y_1)^{\alpha_1} (x_2 y_2)^{\alpha_2} \dots (x_n y_n)^{\alpha_n} \\ &= k! \sum_{\alpha} \frac{x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}}{\alpha_1! \alpha_2! \dots \alpha_n!} y_1^{\alpha_1} y_2^{\alpha_2} \dots y_n^{\alpha_n}. \end{aligned}$$

Received by the editors October 7, 1994.

1991 *Mathematics Subject Classification.* Primary 33E99; Secondary 26C99.

Key words and phrases. Homogeneous polynomials, addition formula.

Thus if $p(y) = \sum_{\alpha} a_{\alpha} y_1^{\alpha_1} y_2^{\alpha_2} \dots y_n^{\alpha_n}$, we have

$$\begin{aligned} & ((x_1 y_1 + \dots + x_n y_n)^k, p(y)) \\ &= \left(k! \sum_{\alpha} \frac{x_1^{\alpha_1} \dots x_n^{\alpha_n}}{\alpha_1! \dots \alpha_n!} \frac{\partial^{\alpha_1}}{\partial y_1^{\alpha_1}} \dots \frac{\partial^{\alpha_n}}{\partial y_n^{\alpha_n}} \sum_{\beta} a_{\beta} y_1^{\beta_1} \dots y_n^{\beta_n} \right) \\ &= k! \sum_{\alpha} a_{\alpha} x_1^{\alpha_1} \dots x_n^{\alpha_n} = k! p(x). \end{aligned}$$

Our next result is an addition formula for orthonormal homogeneous polynomials.

Theorem 2 (Addition Formula). *Let $\{p_j(x)\}_{j=1}^{d_k}$ be an orthonormal basis for H_k . Then*

$$(2) \quad (x_1 y_1 + x_2 y_2 + \dots + x_n y_n)^k = k! \sum_{j=1}^{d_k} p_j(x) p_j(y).$$

Proof. Let $x = (x_1, x_2, \dots, x_n)$ and $\alpha_j = (\alpha_j^1, \alpha_j^2, \dots, \alpha_j^n)$. For convenience we write $x^{\alpha_j} = x_1^{\alpha_j^1} x_2^{\alpha_j^2} \dots x_n^{\alpha_j^n}$. Then expanding the power, we have

$$(x_1 y_1 + \dots + x_n y_n)^h = \sum c_j x^{\alpha_j} y^{\alpha_j},$$

where x^{α_j} and y^{α_j} are homogeneous monomials of degree k . Since $\{p_j(x)\}_{j=1}^{d_k}$ is a basis for H_k , there exist constants a_{jm} such that

$$y^{\alpha_j} = \sum_{m=1}^{d_k} a_{jm} p_m(y).$$

Thus

$$\begin{aligned} (x_1 y_1 + \dots + x_n y_n)^k &= \sum_j c_j x^{\alpha_j} \left(\sum_m a_{jm} p_m(y) \right) \\ &= \sum_j g_j(x) p_j(y), \end{aligned}$$

where the last expression is a rearrangement of the previous sum. The $g_j(x)$ are linear combinations of the monomials x^{α_j} , and thus are homogeneous polynomials of degree k in x . Further, since the $p_j(x)$ are orthonormal,

$$((x_1 y_1 + \dots + x_n y_n)^k, p_l(y)) = \left(\sum_j g_j(x) p_j(y), p_l(y) \right) = g_l(x).$$

But by the identity given in Theorem 1,

$$((x_1 y_1 + \dots + x_n y_n)^k, p_l(y)) = k! p_l(x).$$

Thus,

$$g_j(x) = k! p_j(x), \quad j = 1, 2, \dots, d_k,$$

which completes the proof.

If $\alpha_j = (\alpha_j^1, \alpha_j^2, \dots, \alpha_j^n)$, for convenience we write $\alpha_j! = \alpha_j^1! \alpha_j^2! \dots \alpha_j^n!$. The result of Theorem 2 shows that the multinomial expansion

$$(x_1 y_1 + x_2 y_2 + \dots + x_n y_n)^k = \sum \frac{k!}{\alpha_j!} x^{\alpha_j} y^{\alpha_j} = k! \sum \frac{x^{\alpha_j}}{\sqrt{\alpha_j!}} \frac{y^{\alpha_j}}{\sqrt{\alpha_j!}}$$

is no more than an addition formula for the orthonormal basis $\{x^{\alpha_j} / \sqrt{\alpha_j!}\}_{j=1}^{d_k}$ of H_k . The significance of the result (2) lies with the fact that it holds for any orthonormal basis $\{p_j(x)\}_{j=1}^{d_k}$, not merely the monomials $\{x^{\alpha_j} / \sqrt{\alpha_j!}\}_{j=1}^{d_k}$.

The addition formula of Theorem 2 yields an identity which contains the Pythagorean theorem $\sin^2 \theta + \cos^2 \theta = 1$ as a special case:

Theorem 3 (Pythagorean Identity). *Suppose $\{p_j(x)\}_{j=1}^{d_k}$ is an orthonormal basis for H_k . Then on the unit sphere $|s| = \sqrt{s_1^2 + s_2^2 + \dots + s_n^2} = 1$,*

$$\sum_{j=1}^{d_k} [p_j(s)]^2 = \frac{1}{k!}.$$

Proof. Letting $x = y$ in the addition formula yields

$$(x_1^2 + x_2^2 + \dots + x_n^2)^k = k! \sum_{j=1}^{d_k} [p_j(x)]^2,$$

from which the result is immediate.

Most inner products are integrals. Although the inner product we have used here is defined by a derivative, it is far from artificial. Indeed, this inner product allows us to view Taylor series as Fourier series. That is, suppose we have the Taylor series

$$f(x) = \sum_{j=0}^{\infty} a_j x^{\alpha_j}.$$

Rewriting the series as

$$f(x) = \sum_{j=1}^{\infty} b_j \frac{x^{\alpha_j}}{\sqrt{\alpha_j!}},$$

and noting that the monomials $\{x^{\alpha_j} / \sqrt{\alpha_j!}\}_{j=1}^{\infty}$ are orthonormal with respect to the inner product (1), we have

$$(x^{\alpha_j} / \sqrt{\alpha_j!}, f(x)) = b_j.$$

These simple results appear to have considerable analytic significance. To see this, one need only consider the kernel

$$\sum_{k=0}^{\infty} \sum_{j=1}^{d_k} p_k^j(x) p_k^j(y) = \sum_{k=0}^{\infty} \frac{(x \cdot y)^k}{k!} = e^{x \cdot y},$$

and the reproducing formula

$$(f(y), e^{x \cdot y}) = f(x).$$

Here $\{p_k^j(x)\}_{j=1}^{d_k}$, $k = 0, 1, 2, \dots$, are arbitrary orthonormal bases for the homogeneous polynomials H_k of degree k .

REFERENCES

1. A. Erdelyi, *Higher transcendental functions*, Vol. 2, McGraw-Hill, New York, 1953. MR **15**:419i

DEPARTMENT OF MATHEMATICS, GREENSBORO COLLEGE, GREENSBORO, NORTH CAROLINA
27401