FINITE CW COMPLEXES WITH MAXIMAL TORSION GAPS

HUALE HUANG

(Communicated by Thomas Goodwillie)

ABSTRACT. We investigate some properties of finite CW complexes with maximal homotopy torsion gaps and prove a revision of the Halperin conjecture under an additional condition.

1. INTRODUCTION

Let X be a simply connected finite CW complex and $n_X = \max\{i | H_i(X, \mathbb{Q}) \neq 0\}$. On the other hand, $\pi_i(X)$ is the direct sum of finitely many copies of $\mathbb{Z}$ and a finite abelian group. We call an interval $[k, l]$ a torsion gap for X if $\pi_k(X)$ and $\pi_l(X)$ both contain copies of $\mathbb{Z}$ and $\pi_i(X)$ ($k < i < l$) is finite. In [4], S. Halperin proved

Theorem 1. *Let X be a simply connected finite CW complex. If $[k, l]$ is a torsion gap for X , then $l - k < n_X$.*

Then, Halperin proposed the

Conjecture. *If a simply connected finite CW complex X has at least one torsion gap of the form $[k, k + n_X - 1]$, then X has the rational homotopy type of a wedge of n_X -spheres.*

In [6], the author resolved the conjecture in the negative. However, those CW complexes with maximal torsion gaps are very strict. In fact, we have

Theorem A. *If a simply connected finite CW complex X has at least one torsion gap of the form $[k, k + n_X - 1]$, then the k -connected covering $X(k)$ of X has the rational homotopy type of a wedge of spheres.*

Corollary 1. *If X has infinitely many torsion gaps of the form $[k, k + n_X - 1]$, let $k_0 = \min\{k | [k, k + n_X - 1] \text{ is a torsion gap for } X\}$. Then if $[k, l]$ is a torsion gap for X and $k \geq k_0$, then $l = k + n_X - 1$ and $k \equiv 1 \pmod{n_X - 1}$. In particular, if all the torsion gaps for X have the form $[k, k + n_X - 1]$, then X has the rational homotopy type of a wedge of n_X -spheres.*

Proof. Since $[k_0, k_0 + n_X - 1]$ is a torsion gap for X , the k_0 -connected covering $X(k_0)$ of X is $(k_0 + n_X - 2)$ -connected. By Theorem 1, there exists $m \leq n_X - 1$, s.t. $\pi_{k_0+n_X-1+m}(X) \supset \mathbb{Z}$. As $k_0 \geq 2$, $(k_0, n_X - 1 + m) + 1 < 2(k_0 + n_X - 1)$; hence

$$H^{k_0+n_X-1}(X(k_0), \mathbb{Q}) \neq 0, \quad H^{k_0+n_X-1+m}(X(k_0), \mathbb{Q}) \neq 0.$$

Received by the editors August 24, 1994 and, in revised form, January 19, 1995.

1991 *Mathematics Subject Classification.* Primary 55P62.

By Theorem A, $X(k_0)$ has the rational homotopy type of a wedge of spheres. So we have

$$X(k_0)_Q \cong (S^{k_0+n_X-1} \vee S^{k_0+n_X-1+m} \vee Y)_Q.$$

Let $d = (k_0 + n_X - 2, k_0 + n_X - 2 + m)$; then $d|m$. Then there is some integer l s.t. for all $k \geq 0$, $\pi_{l+kd}(X) \supset \mathbb{Z}$. Hence $d = m = n_X - 1$, and $k_0 \equiv 1 \pmod{n_X - 1}$. Using the same argument, we can prove that if $[k, l]$ is a torsion gap for X and $k \geq k_0$, then $l = k + n_X - 1$ and $k \equiv 1 \pmod{n_X - 1}$. $\square$

As a revision of the Halperin conjecture, we have

Theorem B. *If a simply connected finite CW complex X has infinitely many torsion gaps of the form $[k, k + n_X - 1]$, and if $\text{cat}(X) \leq 2$, then X has the rational homotopy type of a wedge of n_X -spheres.*

2. PROOF OF THEOREM A

The main tool in this paper is Sullivan's theory of minimal models. A commutative differential graded algebra (CDGA) over the rational number field Q is a differential graded algebra (A, d) , where $A = \sum_{p \geq 0} A^p$, $ab = (-1)^{pq}ba$ for $a \in A^p$, $b \in A^q$, and $d: A^p \rightarrow A^{p+1}$ is a derivation. Its cohomology is denoted by $H^*(A, d)$. If $V = \sum_{p \geq 0} V^p$ is a graded vector space over Q where V^p is finite dimensional for all p , then we denote by

$$\bigwedge V = \text{Exterior algebra } (V^{\text{odd}}) \otimes \text{Symmetric algebra } (V^{\text{even}})$$

the free commutative graded algebra on V . A Sullivan minimal model is a CDGA of the form $(\bigwedge V, d)$, where V^i is finite dimensional for $i \geq 1$ and $\text{Im } d \subset (\bigwedge V)^+ \cdot (\bigwedge V)^+$, where $(\bigwedge V)^+$ is the ideal generated by V . To any simply connected space X s.t. each $H_i(X, Q)$ is finite dimensional, $i \geq 2$, Sullivan associates a unique minimal model $(\bigwedge V, d)$, satisfying the following properties:

$$H(\bigwedge V, d) \cong H^*(X, Q); \quad V \cong \text{Hom}_{\mathbb{Z}}(\pi_*(X), Q).$$

To prove Theorem A, we need two lemmas. Let $(\bigwedge V, d)$ be a minimal model, then the augmentation $\bigwedge V \rightarrow Q$ has a minimal model of the form $(\bigwedge V \otimes \bigwedge \overline{V}, d) \cong Q$, in which the differential in the quotient model $Q \otimes_{\bigwedge V} (\bigwedge V \otimes \bigwedge \overline{V}) = \bigwedge \overline{V}$ is zero and $V^i \cong \overline{V}^{i-1}$. Thus each $\bigwedge V \otimes (\bigwedge \overline{V})^{\leq r}$ is a sub- $(\bigwedge V, d)$ differential model of $\bigwedge V \otimes \bigwedge \overline{V}$. Thus we get a spectral sequence E_i with $E_0^{p,q} = (\bigwedge V)^p \otimes (\bigwedge \overline{V})^q$.

Lemma 1. *If V is finite dimensional and we let $m = \max\{i | V^i \neq 0\}$, then the spectral sequence E_i degenerates at E_{m+1} , that is, $E_{m+1} = E_{m+2} = \cdots = E_{\infty}$.*

Proof of Lemma 1. It is enough to prove by induction on r that the spectral sequence for $\bigwedge V^{\leq r} \otimes (\bigwedge \overline{V}^{\leq r})$ collapses at the E_{r+1} -term. The inductive hypothesis then gives the isomorphism

$$(E_{r+1}(\bigwedge V^{\leq r+1} \otimes (\bigwedge \overline{V}^{\leq r})), d_{r+1}) \cong (E_{r+1}(\bigwedge V^{r+1} \otimes (\bigwedge \overline{V}^r)), d),$$

with $d(\overline{v}) = v$. This clearly implies the result. $\square$

From Lemma 1, we easily get

Corollary 2. *Let V be finite dimensional and $m = \max\{i | V^i \neq 0\}$. If $H^k(\bigwedge V, d) \neq 0$, $H^l(\bigwedge V, d) \neq 0$, and $H^i(\bigwedge V, d) = 0$ for $k < i < l$, then $l - k \leq m$.*

Proof. If $l - k > m$, then by Lemma 1 $E_\infty^{l,0} \neq 0$, which is impossible. $\square$

Like homotopy torsion gap, we define homology torsion gap as follows. Let X be a simply connected CW complex, with finite Betti numbers. If $H_k(X, Q) \neq 0$, $H_l(X, Q) \neq 0$, and $H_i(X, Q) = 0$ for $k < i < l$, then we call $[k, l]$ a homology torsion gap for X .

By Corollary 2, we obtain

Corollary 3. *Let X be a simply connected rational elliptic CW complex and $m = \max\{i | \pi_i(X) \otimes Q \neq 0\}$. If $[k, l]$ is a homology torsion gap for X , then $l - k \leq m$.*

Remark. Corollary 3 is a dual form of Theorem 1 (Halperin), in the sense of Eckman-Hilton duality; furthermore, they are equivalent.

We also need the following lemma, which easily followed from S. Halperin and G. Levin [5].

Lemma 2. *Let X be a simply connected finite CW complex with $(\bigwedge V, d)$ as minimal model. Denote by J the differential ideal of $(\bigwedge V, d)$ defined by $J = (\bigwedge V)^{>n_X} \oplus W$, where W is the complement in $(\bigwedge V)^{n_X}$ of the cocycles. Let $(\bigwedge V^{\leq k}, d)$ be a submodel of $(\bigwedge V, d)$ and $\bigwedge V^{\leq k} \otimes \bigwedge U$ be a minimal of $\bigwedge V^{\leq k} \rightarrow Q$. Then the minimal model of $(\bigwedge V^{\leq k} \otimes \bigwedge U) \otimes \bigwedge_{V \leq k} \bigwedge V/J = \bigwedge U \otimes \bigwedge V/J$ has the same minimal model of $X(k)$, the k -connected covering of X .*

A CDGA morphism, ϕ , is a quism if $H(\phi)$ is an isomorphism. The equivalence class of a CDGA (under the equivalence relation generated by quisms) is called its homotopy type. A CDGA is said to be a wedge of spheres if it has the homotopy type of a connected CDGA H with differential zero and satisfies $H^+ \cdot H^+ = 0$.

Now we prove Theorem A.

Proof of Theorem A. For $\bigwedge V^{\leq k} \otimes \bigwedge U$ and $\bigwedge U \otimes \bigwedge V/J$, we have two spectral sequences $\overline{E}_m, E_m$ respectively (filtered by the degree of $\bigwedge U$) and morphism $\phi_m : \overline{E}_m \rightarrow E_m$ induced by $\bigwedge V^{\leq k} \rightarrow \bigwedge V^{\leq k}/J$. Because $\overline{E}_2^{i,*} \rightarrow E_2^{i,*}$ is identical for $i < k + n_X$, $\phi_{k+1} : \overline{E}_{k+1}^{i,*} \rightarrow E_{k+1}^{i,*}$ is an isomorphism for $i \leq n_X - 1$, while $\overline{E}_{k+1}^{i,*} = 0$ for all $(i, *) \neq (0, 0)$. Thus $E_{k+1}^{i,*} \neq 0$ only for $i = n_X$ or $(i, *) = (0, 0)$. Therefore, we can choose the representatives of $H^+(\bigwedge U \otimes \bigwedge V/J)$ from $\bigwedge U \otimes (\bigwedge V/J)^{n_X}$. Obviously their product is zero; thus $\bigwedge U \otimes \bigwedge V/J$ is a wedge of spheres.

By Lemma 2, we know that the minimal model of $X(k)$ is a wedge of spheres; thus, we get Theorem A. $\square$

3. THE SECOND MAIN THEOREM

Before proving Theorem B, we recall some definitions. In [2], Felix and Halperin define the rational L.S. category, $\text{Cat}_0(X)$, via the Sullivan minimal model $(\bigwedge V, d)$ for X , $\text{Cat}_0(X) \leq m$ if and only if in the diagram

$$\begin{array}{ccc} \bigwedge V & \xrightarrow{\pi} & \bigwedge V / \bigwedge^{>m+1} V \\ & \searrow i & \uparrow \phi \cong \\ & & \bigwedge V \otimes \bigwedge W \end{array}$$

there exists r , so that ri is identity, where π is the projection and ϕ is a quism. When X is a simply connected space with finite Betti numbers, they prove $\text{Cat}_0(X) = \text{Cat}(X_Q)$.

Let L be a graded Lie algebra. Then the radical of L , $\text{rad } L$, is the sum of all resolvable ideals of L .

$$\text{gl. dim } L = \text{gl. dim } UL = \sup\{i \mid \text{Ext}_{UL}^i(Q, Q) \neq 0\},$$

$$\text{depth } L = \text{depth } UL = \inf\{i \mid \text{Ext}_{UL}^i(Q, UL) \neq 0\}.$$

We need the following theorems from Felix et al. [4] and Felix [1].

Theorem 2. *Let X be a simply connected space with finite Betti numbers. Then*

- (1) $\text{depth } H_*(\Omega X, Q) \leq \text{Cat}_0(X) \leq \text{gl. dim } H_*(\Omega X, Q)$,
- (2) *if $\text{depth } H_*(\Omega X, Q) = \text{Cat}_0(X)$, then*

$$\text{Cat}_0(X) = \text{gl. dim } H_*(\Omega X, Q).$$

Theorem 3. *Let L be a graded Lie algebra and $\text{depth } L = m < \infty$. Then*

- (1) $\dim(\text{rad } L)_{\text{even}} \leq m$,
- (2) *$\text{rad } L$ is finite dimensional,*
- (3) *if $\dim(\text{rad } L)_{\text{even}} = m$, then $L = \text{rad } L$.*

Theorem 4. *If $\text{Cat}_0(X) \leq 2$ and $\text{gl. dim } H_*(\Omega X, Q) = 2$, then X is a coformal space, that is, if $(\bigwedge V, d)$ is the Sullivan minimal model, then $d = d_2$.*

Proof of Theorem B. First, $L = \pi_*(\Omega X) \otimes Q$ is a graded Lie algebra by the Samelson Product. We claim that $\text{depth } L \leq 1$. Otherwise $\text{depth} = 2 = \text{Cat}_0(X)$; hence, by Theorem 2, $\text{gl. dim } L = 2$. According to Theorem 4, X is a coformal space; hence, the Sullivan minimal model has the form $(\bigwedge V, d)$, $d = d_2$. Now let $k_0 = \min\{k \mid [k - n_X + 1, k] \text{ is a torsion gap for } X\}$ and take $\alpha \in V$ s.t. $|\alpha| > 2k_0$. Then

$$d\alpha = \beta_1\gamma_1 + \beta_2\gamma_2 + \cdots + \beta_m\gamma_m.$$

We may suppose $|\beta_1| = \max\{|\beta_i|, |\gamma_i|\}$. Then $|\beta_1| \geq \frac{|\alpha|}{2} > k_0$; thus $|\beta_1| \equiv 1 \pmod{n_X - 1}$ (by Corollary 1). We can suppose that $d\alpha = \beta_1\gamma + \cdots$ (the other terms having no β_1) in view of $dd\alpha = 0$, $d\gamma \cdot \beta_1 = 0$, hence $d\gamma = 0$ and $|\gamma| = n_X$ because $|\beta_1| \equiv |\alpha| \equiv 1 \pmod{n_X - 1}$. Therefore $X_Q \cong (S^{n_X} \vee Y)_Q$, but this is impossible (because of $\text{depth } \pi_*(\Omega(S^{n_X} \vee Y)) \otimes Q = 1$). So we have $\text{depth } L \leq 1$.

Next, we claim that if $x \in \pi_*(X) \otimes Q \cong \pi_{*-1}(\Omega X) \otimes Q$ and $|x| \not\equiv 1 \pmod{n_X - 1}$, then $x \in \text{rad } L$.

In fact, we denote by I the Lie ideal generated by x ; then we say $I_{\geq k_0} = 0$. Otherwise there is some $\alpha \neq 0$, $\alpha \in I_{\geq k_0}$. Since $I_{\geq k_0}$ is a free Lie algebra, there is some $\omega \in I_{\geq k_0}$ s.t. $[\alpha, \omega] \neq 0$. We write $\alpha = \sum_i [\beta_{1i}, \beta_{2i}]$, with $\beta_{1i} \in I$. The Jacobi identity implies that there is some i and some $u \in I_{\geq k_0}$, s.t. $[\beta_{1i}, u] \neq 0$. Clearly $|\beta_{1i}| < |\alpha|$. We continue the same procedure with β_{1i} in place of α . After finitely many steps, we get $[x, u] \neq 0$, where $|u| \equiv 1 \pmod{n_X - 1}$ and $|u| > k_0$, but $\pi_{|u|+|X|}(X) \otimes Q = 0$. This is a contradiction, so $I_{\geq k_0} = 0$, hence $x \in \text{rad } L$.

We know $\text{depth } L = 1$, by Theorem 3, and $(\text{rad } L)_{\text{even}} = 0$, hence $\pi_{2k+1}(X) \otimes Q = 0$ if $2k+1 \not\equiv 1 \pmod{n_X - 1}$. If n_X is odd, then we must have some $\alpha \in \pi_{n_X}(X) \otimes Q$, and $d\alpha = 0$, so $X_Q = (S^{n_X} \vee Y)_Q$. If $Y_Q \neq \bigvee S^{n_X}$, by the following Lemma 3, X has no torsion gap of the form $[k, k + n_X - 1]$, so that $Y_Q \cong \bigvee S^{n_X}$, and X has the rational homotopy type of a wedge of n_X -spheres. If n_X is even and if $m < n_X$, then $\bigwedge V^{\leq n_X}$ is a polynomial algebra and $dV^{\leq n_X} = 0$. Let $\alpha \in \pi_m(X) \otimes Q$, where $m = \min\{k \mid \pi_k(x) \otimes Q \neq 0\}$, and $\beta \in (\bigwedge V)^{n_X}$ representing a cohomology of $H^{n_X}(\bigwedge V, d)$. Then $d(\alpha\beta) = 0$, $|\alpha\beta| = m + n_X < 2n_X - 1$, so that $\alpha\beta$ cannot be a

coboundary; hence $H^{m+n_X}(\bigwedge V, d) \neq 0$. This is in contradiction with the definition of n_X . So $m = n_X$, hence, by the rational Hurewicz Theorem, X has the rational homotopy type of a wedge of n_X -spheres. $\square$

Lemma 3. *If $X = S^{n_X} \vee Y$ and $Y_Q \neq (\bigvee S^{n_X})_Q$, then X has no torsion gap of the form $[k, k + n_X - 1]$.*

Proof. As a Lie algebra, $\pi_*(\Omega X) \otimes Q$ is the free product of $\pi_*(S^{n_X}) \otimes Q$ and $\pi_*(Y) \otimes Q$. It is easy to show that X has no torsion gap of the form $[k, k + n_X - 1]$. $\square$

ACKNOWLEDGMENT

Finally, the author would like to acknowledge Professor Shen Xinyao for his helpful comments, and to also acknowledge the referee for offering a corrective proof of Lemma 1 and other useful remarks.

REFERENCES

- [1] Y. Félix, *La dichotomie elliptique-hyperbolique en homotopie rationnelle*, Asterisque **176** (1989). MR **91c**:55016
- [2] Y. Félix and S. Halperin, *Rational L.S. category and its application*, Trans. Amer. Math. Soc. **273** (1982), 1–37. MR **84h**:55011
- [3] Y. Félix, S. Halperin, C. Jacobson, C. Löfwall, and J.-C. Thomas, *The radical of the homotopy Lie algebra*, Amer. J. Math. **110** (1988), 301–322. MR **89d**:55029
- [4] S. Halperin, *Torsion gaps in the homotopy of finite complexes. II*, Topology **30** (1991), 471–478. MR **92f**:55014
- [5] S. Halperin and G. Levin, *High skeleta of CW-complexes*, Lecture Notes in Math, vol. 1183, Springer-Verlag, Berlin, 1986, pp. 211–217. MR **87m**:55013
- [6] H. Huang, *On a conjecture of S. Halperin*, Northeast Math. J. **10** (1994), 517–524.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, BEIJING 100080, PEOPLE'S REPUBLIC OF CHINA

Current address: Ph.D. Program in Mathematics, Graduate Center, CUNY, 33 W 42 St., New York, New York 10036

E-mail address: `hhuang@email.gc.cuny.edu`