EMBEDDING OF A BANACH ALGEBRA A INTO $L(A')$

ETIENNE DESQUITH

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Given a Banach space X , we have shown in 1994 that a product can be defined on it in such a way that the resulting Banach algebra is isomorphic to a compact subalgebra of the algebra $L(X')$ of all bounded linear operators on the topological dual X' of X . Our purpose here is to prove that, more generally, any Banach algebra A admitting a *left approximate identity*, is isomorphic to a subalgebra of $L(A')$, the isomorphism being isometric, provided the approximate identity is bounded by 1. As a consequence, we get a factorization through $L(A')$, of the elements in A' .

An abstract C^* -algebra is isometrically isomorphic to a uniformly closed subalgebra of $L(H)$, the algebra of all bounded linear operators on some Hilbert space H . When A is no longer a C^* -algebra, but some Banach algebra with a left approximate identity, we still get an isomorphism from A into an algebra of bounded linear operators on some Banach space E which is no longer a Hilbert space. More precisely, E is the topological dual A' of A . When the approximate identity is a subset of the unit sphere of A , we get a similar identification of A as in the case of a C^* -algebra, in the following way:

Any Banach algebra A with a left approximate identity bounded by 1 is isometrically isomorphic to a uniformly closed subalgebra of $L(A')$.

When A is a commutative Banach $*$ -algebra, then the isomorphism is an involution-preserving map.

We mainly derive from the above identification of A with a subalgebra of $L(A')$, a particular factorization of the elements of A' , through $L(A')$.

Let A and B be Banach spaces each of which is endowed with a linear involution denoted by “ $*$ ” and “ $\bullet$ ” respectively. We consider:

(1) The linear involution “ $\otimes$ ” defined on the space $L(A, B)$ by:

$$(1) \quad P^{\otimes}(a) = P(a^*)^{\bullet} \quad (P \in L(A, B); a \in A).$$

(2) The linear involution “ $\ltimes$ ” defined on the space A' by:

$$(2) \quad T^{\ltimes}(x) = \overline{T(x^*)} \quad (x \in A; T \in A')$$

($T^{\ltimes}(x) = \overline{T(x)}$, if A does not have an involution, where the bar denotes complex conjugation).

The following is the main result of the paper.

Theorem 1. *Let A be a Banach algebra with a left approximate identity $(u_{\alpha})_{\alpha \in I}$ bounded by 1. Then*

Received by the editors September 29, 1994 and, in revised form, March 13, 1995.
 1991 *Mathematics Subject Classification.* Primary 46H15.

- (i) *There exists an isometric isomorphism σ from A onto a uniformly closed subalgebra of $L(A')$.*
- (ii) *If A is a commutative Banach $*$ -algebra, then the isometry σ is $*$ -map in the following sense: $\sigma(x^*) = \sigma(x)^\otimes, \forall x \in A$.*

Proof. For all $x, y \in A$, and all $T \in A'$, we set:

$$(3) \quad (\sigma(x)(T))(y) = T(yx).$$

Then easy computations show that the map $x \mapsto \sigma(x)$ from A into $L(A')$ induced from relation (3) above is a homomorphism. Moreover, σ is one-to-one as follows: if $x, y \in A$ are such that $\sigma(x) = \sigma(y)$, then for all $z \in A$ and $T \in A'$, we have $T(zx) = T(zy)$. But since A admits a left approximate identity $(u_\alpha)_{\alpha \in I}$, we get

$$(4) \quad T(u_\alpha x) = T(u_\alpha y) \quad (\alpha \in I, T \in A'),$$

that is,

$$\begin{aligned} \|T(x - y)\| &= \|T(u_\alpha(x - y)) - T(x - y)\| \\ &\leq \|T\| \|u_\alpha(x - y) - (x - y)\|, \end{aligned}$$

from which it follows that

$$(5) \quad T(x) = T(y) \quad \forall T \in A',$$

and therefore, $x = y$.

Moreover, for all $x \in A$, it is easily checked that

$$(6) \quad \|\sigma(x)\| \leq \|x\|.$$

Conversely, if A_1 (resp. A'_1) denotes the closed unit ball of A (resp. A'), then given $x \in A$, we have, for all $\alpha \in I$:

$$\begin{aligned} \|u_\alpha x\| &\leq \sup\{\|yx\|; y \in A_1\} \\ &\leq \sup\{\sup\{|T(yx)|; T \in A'_1\}; y \in A_1\} \\ &\leq \sup\{\sup\{|(\sigma(x)T)(y)|; T \in A'_1\}; y \in A_1\}. \end{aligned}$$

It follows, when α tends to infinity, that

$$(7) \quad \|x\| \leq \|\sigma(x)\|,$$

which, together with relation (6), shows that σ is isometric. The range $\sigma(A)$ of σ is clearly closed under the uniform norm of $L(A')$.

(ii) If A is in addition a commutative $*$ -algebra, then σ is an involution-preserving map, for if $x, y \in A$ and $T \in A'$, then

$$\begin{aligned} (\sigma(x^*)(T))(y) &= T(yx^*) = T(x^*y) \quad (A \text{ is commutative}) \\ &= \overline{T^\times(y^*x)} = \overline{(\sigma(x)(T^\times))(y^*)} \\ &= (\sigma(x)(T^\times))^\times(y) = ([\sigma(x)]^\otimes(T))(y), \end{aligned}$$

from which it follows that:

$$(8) \quad \sigma(x^*) = [\sigma(x)]^\otimes,$$

and the proof is complete. $\square$

Corollary 2. *If A is a commutative C^* -algebra, then $\sigma(A)$ endowed with the involution $\otimes$ is a commutative C^* -algebra.*

Proof. Each C^* -algebra admits a two-sided approximate identity bounded by 1. According to Theorem 1 (ii) above, $\sigma(A)$ is a commutative $\otimes$ -sub-algebra of $L(A')$ satisfying, for all $x \in A$:

$$\|\sigma(x) \otimes \sigma(x)\| = \|\sigma(x^*x)\| = \|x^*x\| = \|x^2\| = \|\sigma(x)\|^2. \quad \square$$

Corollary 3. *Let G be a topological locally compact group and $C_0(G)$ the algebra of all continuous complex functions on G vanishing at infinity.*

- (i) *If $M^1(G)$ denotes the space of all bounded Radon measures on G , then $L(M^1(G))$ contains an isometric image of $C_0(G)$.*
- (ii) *If G is abelian, then $L^1(G)$ is isometrically isomorphic to a subalgebra of $L(L^\infty(G))$.*

Proof. (i) $A = C_0(G)$ endowed with the sup-norm and the involution defined by complex conjugation is a commutative C^* -algebra. Hence according to Theorem 1 and Corollary 2, A may be identified with a commutative C^* -algebra in $L(A')$. But the Riesz representation theorem ensures that A' is isometrically isomorphic to $M^1(G)$.

(ii) If G is abelian, the group algebra $L^1(G)$ endowed with the usual convolution of functions is a commutative Banach $*$ -algebra with a two-sided approximate identity bounded by 1. Hence Theorem 1 holds, and the result follows. $\square$

We shall need the following lemma for the next result.

Lemma 4. *Let X, Y, Z be Banach spaces, and P, Q be linear operators from X into Y and from X into Z respectively, such that $\text{Ker}(P)$ and $\text{Ker}(Q)$ satisfy $\text{Ker}(P) \subset \text{Ker}(Q)$. Then there exists a linear map S from the range $R(P)$ of P into Z satisfying $Q = SP$.*

Proof. Straightforward. $\square$

We have the following particular decomposition of the elements of A' :

Theorem 5. *Let A be a Banach algebra with a left approximate identity bounded by 1. Then, each $T \in A'$ admits the following factorization:*

$$(9) \quad T = \Phi \Theta \sigma,$$

where

- σ is the isometry in Theorem 1,
- Θ is a continuous linear map from $\sigma(A) \subset L(A')$ into A' , and
- Φ is an element in the bidual A'' of A .

Proof. Let $(u_\alpha)_{\alpha \in I}$ be a left approximate identity. For each $T \in A'$, we define the map Ψ_T as follows:

$$(10) \quad x \mapsto \psi_T(x) = \sigma(x)(T) \quad (x \in A).$$

ψ_T is well-defined, linear and continuous ($\|\psi_T\| \leq \|T\|$), and takes its values in A' . Moreover, using the existence of a (bounded) left approximate identity in A , one shows that

$$(11) \quad \text{Ker}(\psi_T) \subset \text{Ker}(T)$$

$[(\sigma(x)(T))(\cdot) = T(\cdot x)]$. But then, according to Lemma 4, there exists a linear map φ from the range $\psi_T(A)$ of ψ_T onto $\mathbb{C}$ such that $T = \varphi \psi_T$. φ is continuous on

$\psi_T(A)$. Indeed, let $\varepsilon > 0$ be given and let $x, y \in A$ be such that $\|\psi_T(x) - \psi_T(y)\| < \varepsilon' = \frac{\varepsilon}{2(1+\|T\|)}$. This implies, for all $z \in A_1$, $|T(z(x-y))| < \varepsilon'$.

It follows that, for each element u_α of a left approximate identity $(u_\alpha)_{\alpha \in I}$ bounded by 1, the inequality $|T(u_\alpha(x-y))| < \varepsilon'$ holds, and therefore:

$$\begin{aligned} |T(x-y)| &\leq |T(u_\alpha(x-y))| + |T(x-y) - T(u_\alpha(x-y))| \\ &< \varepsilon' + \|T\| \|(x-y) - u_\alpha(x-y)\| \\ &< (1 + \|T\|) \varepsilon' \quad \left(\text{since } \lim_{\alpha \rightarrow +\infty} [(x-y) - u_\alpha(x-y)] = 0 \right) \\ &< (1 + \|T\|) \frac{\varepsilon}{2(1 + \|T\|)} = \frac{\varepsilon}{2}. \end{aligned}$$

Hence $|\varphi(\psi_T(x)) - \varphi(\psi_T(y))| < \frac{\varepsilon}{2}$, and φ is continuous on $\psi_T(A)$. Then, the Hahn-Banach continuous extension theorem yields a continuous linear map Φ from A' onto $\mathbb{C}$ ($\Phi \in A''$) which coincides with φ on $\psi_T(A)$ and satisfies:

$$(12) \quad T = \Phi \psi_T.$$

Furthermore, $\text{Ker}(\sigma) = \{0\} \subset \text{Ker}(\psi_T)$, and once again Lemma 4 yields a linear map Θ from $\sigma(A)$ onto $\psi_T(A)$, which is continuous from the isometric nature of σ , and such that:

$$(13) \quad \psi_T = \Theta \sigma.$$

Hence T finally fulfills $T = \Phi \Theta \sigma$, which is the desired factorization. $\square$

Now, let X be a Banach space and F a continuous linear form on X . If for $x, y \in X$ one defines:

$$(14) \quad xy = F(x)y,$$

then X , endowed with its Banach space norm together with the above product, becomes a complete normed algebra $A(X, F)$, which is a Banach algebra if $\|F\| \leq 1$.

If $f \in X'$, the nonzero vector $x \in X$ is called a *maximal element* of f , if $f(x) = \|f\| \|x\|$ (cf. [5], Definition 15.2.1, page 337).

We have proved in [3] the following theorem and its corollary:

Theorem 6. *Let X be a Banach space. If F is in the unit sphere of X' and admits a maximal element $u \in X$, then $A(X, F)$ is isometrically isomorphic to a uniformly closed subalgebra of the algebra $\mathcal{K}(X')$ of all compact operators on X' (cf. [3], Théorème 2.19).*

Corollary 7. *If X is a reflexive Banach space or the dual space Y' of a Banach space Y , and if F (resp. Φ) is an element of the unit sphere of X' (resp. Y''), then $A(X, F)$ (resp. $A(Y', \Phi)$) is isometrically isomorphic to a uniformly closed subalgebra of $\mathcal{K}(X')$ (resp. $\mathcal{K}(Y'')$) (cf. [3], Corollaire 2.20).* $\square$

The isomorphism ρ in the above theorem satisfies

$$(15) \quad (\rho(x)(T))(y) = T(yx) = F(y)T(x), \quad x, y \in X.$$

Moreover, the range, say $A_\rho(X, F)$ of ρ in $\mathcal{K}(X')$, consists of rank-one operators on X' . Namely,

$$(16) \quad \rho(x)(T) = T(x)F = \hat{x}(T)F, \quad x \in X, T \in X',$$

where $x \mapsto \hat{x}$ denotes the natural linear isometry of X into X'' .

Remark 8. It turns out that Theorem 6 above is a particular case of Theorem 1, according to the following:

Proposition 9. *The Banach algebra $A(X, F)$ in Theorem 6 admits infinitely many left approximate identities bounded by 1. Moreover precisely, $(u_\alpha)_{\alpha \in I}$ is a left approximate identity for $A(X, F)$ if and only if one has:*

$$(17) \quad \lim_{\alpha \rightarrow +\infty} F(u_\alpha) = 1.$$

In particular, if $u \in X$ is such that $F(u) = 1$, then the sequence $(u_n)_{n \geq 1}$ such that $u_n = z_n u$, where $(z_n)_{n \geq 1}$ is any complex sequence in the unit disk ($\|z_n\| \leq 1; n \geq 1$) satisfying $\lim_{n \rightarrow +\infty} z_n = 1$, is a countable left approximate identity for $A(X, F)$, bounded by 1.

Proof. Straightforward. $\square$

We close the sequel with the following two properties of $A_\rho(X, F)$:

Proposition 10. (i) *For all $x \in X$ and $P \in L(X')$, the following condition holds:*

$$(18) \quad P \circ \rho(x)(\cdot) = \hat{x}(\cdot)P(F).$$

(ii) *For all $x \in X$ and $P \in L(X')$, there exists, for each $T \in X'$, some $z_{x,T,P} \in X$ (depending on x, P and T) such that:*

$$(19) \quad (\rho(x) \circ P)|_{\mathbb{C}T} = \rho(z_{x,T,P}),$$

where $(\rho(x) \circ P)|_{\mathbb{C}T}$ stands for the restriction of $(\rho(x) \circ P)$ to the one-dimensional subspace $\mathbb{C}T$ of X' .

Proof. (i) is obvious.

(ii) Let $x \in X$ and $P \in L(X')$. Then, for each $T \in X'$ and $\lambda \in \mathbb{C}$, we have $(\rho(x) \circ P)(\lambda T) = \lambda P(T)(x)F$. But we may choose $z_{x,T,P} \in X$ such that $P(T)(x) = T(z_{x,T,P})$, and it follows that

$$\begin{aligned} (\rho(x) \circ P)(\lambda T) &= \lambda T(z_{x,T,P})F = \lambda \hat{z}_{x,T,P}(T)F \\ &= \lambda \rho(z_{x,T,P})(T) = \rho(z_{x,T,P})(\lambda T). \end{aligned} \quad \square$$

Corollary 11. $A_\rho(X, F)$ is a proper subspace of $\mathcal{K}(X')$.

Proof. $A_\rho(X, F)$ is clearly not reduced to $\{0\}$. Since $\mathcal{K}(X')$ is a two-sided ideal of $L(X')$, the assumption $A_\rho(X, F) = \mathcal{K}(X')$ would imply that, given $x \in X$ and $P \in L(X')$, there exists some $u_{x,P} \in X$ satisfying:

$$(20) \quad P \circ \rho(x) = \rho(u_{x,P}).$$

Let $\text{id}_{L(X')}$ denote the identity map on $L(X')$. We may assume that $P \notin \mathbb{C}\text{id}_{L(X')}$ in relation (20) above. According to relation (15), we get from relation (20) that

$$(21) \quad T(x)P(F)(y) = T(u_{x,P})F(y)$$

for all $y \in X$ and for some nonzero T in X' .

Now, for all $x \in X$ and for any y in the kernel $\text{Ker}(F)$ of F , relation (21) above implies:

$$0 = T(x)P(F)(y) \quad (x \in X, P \in L(X')),$$

and hence $y \in \text{Ker}(P(F))$. Therefore, $\text{Ker}(F) \subset \text{Ker}(P(F))$ and the following holds:

$$(22) \quad P(F) = \lambda_P F, \quad \lambda_P \in \mathbb{C}.$$

But this forces P to satisfy $P \in \mathbb{C} \text{id}_{L(X')}$, in contradiction with the choice of P , and the conclusion follows. $\square$

Remark 12. In the proof of Corollary 11, we have in fact showed that $A_\rho(X, F)$ cannot be a left ideal of $L(X')$. Whether $A_\rho(X, F)$ is a right ideal of $L(X')$ or not, is a question to which we do not yet have an answer.

REFERENCES

- [1] F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer-Verlag, New York, 1973. MR **54**:11013
- [2] E. Desquith, *Structures d'algèbre de Banach sur le dual topologique d'un espace de Banach et applications*, Thèse de Doctorat D'Etat, Université d'Abidjan (FAST), 1992.
- [3] ———, *Sur quelques propriétés de certaines algèbres de Banach dont tout sous-espace vectoriel est un idéal*, Algebras, Groups and Geometries (AGG), vol. 11, no. 3, 1994, pp. 309–327. MR **95k**:46076
- [4] J. Dixmier, *Les C^* -algèbres et leurs représentations*, Gauthier-Villars, Paris, 1964. MR **30**:1404
- [5] L. Narici and E. Beckenstein, *Topological vector spaces*, Marcel Dekker, New York, 1985. MR **87c**:46003

INSTITUT DE RECHERCHES MATHÉMATIQUES (IRMA), 08 BP 2030 ABIDJAN 08, COTE D'IVOIRE