EQUIVALENT CONDITIONS INVOLVING COMMON FIXED POINTS FOR MAPS ON THE UNIT INTERVAL

JACEK R. JACHYMSKI

(Communicated by James E. West)

ABSTRACT. Let g be a continuous self-map of the unit interval I . Equivalent conditions are given to ensure that g has a common fixed point with every continuous map $f : I \mapsto I$ that commutes with g on a suitable subset of I . This extends a recent result of Gerald Jungck.

1. INTRODUCTION

Let f and g be two commuting continuous self-maps of I , the unit interval. It is known that f and g need not have then a common fixed point (for counterexamples, see, e.g., [6]). However, if one of the maps, say g , has appropriate additional properties then f and g possess a common fixed point. In particular, W. Boyce [1, Corollary 5] has shown that it suffices to assume the family $\{g^n : n \in \mathbf{N}\}$, iterates of g , is equicontinuous on I . This result has been extended by J. Cano [2, Theorems 1 and 2] who has required that either g has a closed interval for its set of fixed points $F(g)$, or $F(g)$ coincides with $P(g)$, the set of all periodic points of g . It is worth underlining here that Corollary 5 [1] as well as Theorem 1 [2] give only sufficient conditions for the existence of a common fixed point of f and g .

On the other hand, recently, Gerald Jungck [7, Theorem 3.6] established the following interesting equivalence: a continuous self-map g of I has a common fixed point with every continuous map $f : I \mapsto I$ that nontrivially commutes with g on the set of coincidence points of f and g if and only if $P(g) = F(g)$.

Our purpose here is to give other necessary and sufficient criteria of this type (see Theorems 1, 2 and 3). We also obtain a variant of Jungck's Theorem in more abstract settings as compact and convex subsets of a normed linear space (see Proposition 1).

2. EQUIVALENT CONDITIONS

Following Boyce [1] and Cano [2] we define the classes of maps:

$$\begin{aligned} \mathbf{B} &\doteq \{g : I \mapsto I \mid \{g^n : n \in \mathbf{N}\} \text{ is equicontinuous on } I\}, \\ \mathbf{C}_1 &\doteq \{g : I \mapsto I \mid g \text{ is continuous and } F(g) \text{ is a closed interval}\}, \\ \mathbf{C}_2 &\doteq \{g : I \mapsto I \mid g \text{ is continuous and } F(g) = P(g)\}. \end{aligned}$$

Received by the editors September 9, 1993 and, in revised form, March 29, 1995.

1991 *Mathematics Subject Classification*. Primary 54H25; Secondary 54E45.

Key words and phrases. Common fixed points, equicontinuity, commuting maps, iterates, unit interval.

Let f and g be continuous self-maps of I . Then we have :

- (1) If f and g commute on I and $g \in \mathbf{B}$ then $F(f) \cap F(g) \neq \emptyset$ [1, Corollary 5].
- (2) If f and g commute on I and $g \in \mathbf{C}_1 \cup \mathbf{C}_2$ then $F(f) \cap F(g) \neq \emptyset$; moreover, $\mathbf{B} \subseteq \mathbf{C}_1$. If $g \in \mathbf{B}$ and $F(g)$ is not a singleton then $g \in \mathbf{C}_2$ [2, Theorems 1 and 2].

Next, by [7, Theorem 3.6], $g \in \mathbf{C}_2$ if and only if $F(f) \cap F(g) \neq \emptyset$ for every continuous map $f : I \mapsto I$ such that the set of coincidence points of f and g is non-empty, and f and g commute on it.

Inspired by the last result we give other characterizations of the classes $\mathbf{C}_1$, $\mathbf{C}_2$ and $\mathbf{B}$.

Theorem 1. *Let g be a continuous self-map of I . Then the following conditions are equivalent:*

- (i) $g \in \mathbf{C}_1$;
- (ii) the family $\{g^n : n \in \mathbf{N}\}$ is equicontinuous on $F(g)$, or $F(g)$ is a singleton;
- (iii) g has a common fixed point with every continuous map $f : I \mapsto I$ that commutes with g on $F(g)$.

Proof. (i) $\Rightarrow$ (ii). If $F(g)$ is a singleton, we are done. So suppose $F(g) = [a, b]$ and $a < b$. Obviously, it suffices to show that $\{g^n : n \in \mathbf{N}\}$ is equicontinuous at the points a and b . And because of symmetry, we consider only the point a . The case when $a = 0$ is trivial. So suppose $a > 0$. Fix an $\epsilon \in (0, b - a)$. By the continuity, there exists a $\delta \in (0, \epsilon)$ such that

$$(1) \quad a - \epsilon < g(x) < a + \epsilon \text{ for all } x \in (a - \delta, a) \cap I.$$

We shall apply induction on n to show that, for all $n \in \mathbf{N}$,

$$(2) \quad a - \epsilon < g^n(x) < a + \epsilon \text{ for all } x \in (a - \delta, a) \cap I.$$

By (1), (2) holds if $n = 1$. Assuming (2) holds for $n = 1, 2, \dots, k$, we shall prove it for $k + 1$. Fix an $x \in (a - \delta, a) \cap I$. If $a \leq g^k(x) < a + \epsilon$ then $g^k(x) \in F(g)$ since $a + \epsilon < b$ so (2) is fulfilled for $n = k + 1$. Assume now that $g^k(x) < a$. Then $g^i(x) < a$ for $i = 1, 2, \dots, k$; for otherwise, by induction hypothesis, $a \leq g^i(x) < a + \epsilon$ for some i , $1 \leq i \leq k$ so $g^i(x) \in F(g)$, which implies $g^k(x) \in F(g)$ and hence $g^k(x) \geq a$, a contradiction. Since $F(g) = [a, b]$, we have $g(y) > y$ for all $y \in [0, a)$. In particular, $g^i(x) > g^{i-1}(x)$ for $i = 1, 2, \dots, k$, which implies $g^k(x) > x$. Since $x > a - \delta$ and $g^k(x) < a$, we obtain that $g^k(x) \in (a - \delta, a) \cap I$. By (1), $a - \epsilon < g^{k+1}(x) < a + \epsilon$, which completes the induction.

Since $\delta < \epsilon < b - a$, we have $g^n(x) = x$ for $x \in [a, a + \delta)$ and $n \in \mathbf{N}$. So finally, $a - \epsilon < g^n(x) < a + \epsilon$ for all $x \in (a - \delta, a + \delta)$ and $n \in \mathbf{N}$. This proves $\{g^n : n \in \mathbf{N}\}$ is equicontinuous at the point a .

(ii) $\Rightarrow$ (i). This implication follows from the proof of Cano's Theorem 1 [2].

(i) $\Rightarrow$ (iii). If f commutes with g on $F(g)$ then $F(g)$ is f -invariant so $f|_{F(g)}$ has a fixed point since $F(g)$ is a closed interval.

(iii) $\Rightarrow$ (i). Suppose $F(g)$ is not an interval. There exist $a, b \in F(g)$, $a < b$, such that $(a, b) \cap F(g) = \emptyset$. Define the map f : $f(x) \doteq b$ for $x \in [0, a]$, $f(x) \doteq -x + a + b$ for $x \in (a, b)$, and $f(x) \doteq a$ for $x \in [b, 1]$. Then f is continuous and for $x \in F(g)$, either $x \in [0, a]$ and then $f(g(x)) = g(f(x)) = b$, or $x \in [b, 1]$ and then $f(g(x)) = g(f(x)) = a$. Thus f and g commute on $F(g)$, but $F(f) \cap F(g) = \emptyset$, which contradicts (iii). $\square$

The following example shows that we cannot omit the condition “ $F(g)$ is a singleton” in (ii) of Theorem 1.

Example 1. Define the map g on I as follows:

$$g(x) \doteq 1 \text{ for } x \in [0, \frac{1}{4}], \quad g(x) \doteq -2x + \frac{3}{2} \text{ for } x \in (\frac{1}{4}, \frac{3}{4}), \quad g(x) \doteq 0 \text{ for } x \in [\frac{3}{4}, 1].$$

Clearly, $F(g) = \{\frac{1}{2}\}$ so $g \in \mathbf{C}_1$. However, it is easy to verify that the family $\{g^n : n \in \mathbf{N}\}$ is not equicontinuous at the point $\frac{1}{2}$.

Theorem 2. *Let g be a continuous self-map of I . Then the following conditions are equivalent:*

- (i) $g \in \mathbf{C}_2$;
- (ii) the sequence $\{g^n\}_{n=1}^\infty$ is pointwise convergent on I ;
- (iii) g has a common fixed point with every continuous map $f : I \mapsto I$ that commutes with g on $F(f)$.

Proof. That (i) implies (ii) was proved by S. C. Chu and R. D. Moyer [3, Theorem 1] and, independently, by E. M. Coven and G. A. Hedlund [4, Theorem 2]. To prove (ii) implies (iii) fix an $x \in F(f)$. Since, by the commutativity, $F(f)$ is g -invariant we have $g^n(x) \in F(f)$ for $n \in \mathbf{N}$. By (ii), $\{g^n(x)\}_{n=1}^\infty$ converges to some $z \in I$. Then $z \in F(f) \cap F(g)$ since $F(f)$ is closed and g is continuous. To prove (iii) implies (i) it suffices to show that for any non-empty closed g -invariant set $C \subseteq I$, $C \cap F(g) \neq \emptyset$ and then apply [3, Theorem 1]. Fix such a set C . There exists a continuous map $f : I \mapsto I$ such that $F(f) = C$. If $x \in F(f)$ then $g(f(x)) = g(x)$ and $f(g(x)) = g(x)$ since C is g -invariant. Thus f and g commute on $F(f)$ so, by (iii), $F(f) \cap F(g) \neq \emptyset$, i.e., $C \cap F(g) \neq \emptyset$. $\square$

Remark 1. The sufficiency part of Jungck’s Theorem 3.6 [7] is easily subsumed by Theorem 2: if f and g commute at their coincidence points, $P(g) = F(g)$ and $f(a) = g(a)$ then $f^n(a) = g^n(a)$ for $n \in \mathbf{N}$. By Theorem 2 ((i) $\Rightarrow$ (ii)), $\{g^n(a)\}_{n=1}^\infty$ is convergent to some b . Then, by the continuity, b is a common fixed point of f and g .

Before stating the next theorem let us notice that a common fixed point theorem for a family of commuting maps would be trivial if we assumed one of them had a *unique* fixed point. This justifies a use of the assumption “ $F(g)$ is not a singleton” in Theorem 3 below.

Theorem 3. *Let g be a continuous self-map of I such that $F(g)$ is not a singleton. Then the following conditions are equivalent:*

- (i) $g \in \mathbf{B}$;
- (ii) the sequence $\{g^n\}_{n=1}^\infty$ is uniformly convergent on I ;
- (iii) g has a common fixed point with every continuous map $f : I \mapsto I$ that commutes with g either on $F(f)$, or on $F(g)$.

Proof. (i) $\Rightarrow$ (ii). If $g \in \mathbf{B}$ and $F(g)$ is not a singleton then, by [2, Theorem 2], $g \in \mathbf{C}_2$. By Theorem 2, $\{g^n\}_{n=1}^\infty$ is pointwise convergent on I which implies (ii), since $\{g^n : n \in \mathbf{N}\}$ is equicontinuous.

(ii) $\Rightarrow$ (iii). This implication easily follows from Theorems 1 and 2.

(iii) $\Rightarrow$ (i). By Theorem 2, $\{g^n\}_{n=1}^\infty$ is pointwise convergent, which implies $F(g) = F(g^2)$. On the other hand, by Theorem 1, $F(g)$ is an interval. Therefore, $F(g^2)$ is

an interval, so by [1, Lemma 1 and Theorem 5], $\{g^n : n \in \mathbf{N}\}$ is equicontinuous, i.e., $g \in \mathbf{B}$. $\square$

Remark 2. The condition that $F(g)$ is not a singleton is necessary in Theorem 3 (consider the map $g(x) \doteq 1 - x$ ($x \in I$), for which (i) holds but (ii) is not fulfilled). Further, Example 1 shows that one cannot modify Theorem 3 similarly to Theorem 1. More precisely, we cannot add in (ii) of Theorem 3 the text “or $F(g)$ is a singleton” and, simultaneously, remove the assumption “ $F(g)$ is not a singleton” occurring at the beginning of Theorem 3.

We now give a partial extension of Theorem 2 for maps on subsets of a normed linear space.

Proposition 1. *Let A be a non-empty compact and convex subset of a normed linear space and let g be a continuous self-map of A . Then the following conditions are equivalent:*

- (i) *for any non-empty closed and g -invariant set $C \subseteq A$, $C \cap F(g) \neq \emptyset$;*
- (ii) *g has a common fixed point with every continuous map $f : A \mapsto A$ that commutes with g on $F(f)$.*

Remark 3. It follows from [3, Theorem 1] that in case when $A = I$, the conditions (i) of Proposition 1 and (i) of Theorem 2 are equivalent.

Proof of Proposition 1. (i) $\Rightarrow$ (ii). Let a continuous map $f : A \mapsto A$ commute with g on $F(f)$. Then $F(f)$ is g -invariant and closed. Moreover, by Schauder’s Fixed Point Theorem, $F(f)$ is non-empty. So by (i), we get $F(f) \cap F(g) \neq \emptyset$.

(ii) $\Rightarrow$ (i). Let C be a non-empty closed g -invariant subset of A . We show that $C \cap F(g) \neq \emptyset$. The case when $C = A$ is trivial. So assume $C \neq A$ and fix a point $a \in A \setminus C$. There exists a continuous function $\phi : A \mapsto I$ such that $\phi^{-1}(0) = a$ and $\phi^{-1}(1) = C$ (see, e.g., [5, Theorem 1.5.19, p.69]). Assume further that $0 \in C$. Define a map f by $f(x) \doteq \phi(x)x$ for $x \in A$. Then $f(A) \subseteq A$ by convexity, and $F(f) = C$, since $x = f(x)$ iff $x = 0$ or $\phi(x) = 1$. For $x \in C$, $g(f(x)) = g(x)$ since $f(x) = x$, and $f(g(x)) = g(x)$ since C is g -invariant and $F(f) = C$. Thus f and g commute on $F(f)$. By (ii), $F(f) \cap F(g) \neq \emptyset$, i.e., $C \cap F(g) \neq \emptyset$. In case when $0 \notin C$, fix a point $c \in C$, consider the sets $A' \doteq A - c$, $C' \doteq C - c$, and repeat the above argument to deduce there exists a continuous map $f' : A' \mapsto A'$ such that $F(f') = C'$. Next, define a map f as $f(x) \doteq f'(x - c) + c$ for $x \in A$ and apply (ii) for such a map f to obtain that $C \cap F(g) \neq \emptyset$. $\square$

ACKNOWLEDGEMENT

The author is grateful to the referee for suggesting some stylistic and expositional changes of the text of this paper.

REFERENCES

- [1] W. M. BOYCE, Γ -compact maps on an interval and fixed points, Trans. Amer. Math. Soc. **160** (1971), 87-102. MR **43**:6374
- [2] J. CANO, Common fixed points for a class of commuting mappings on an interval, Proc. Amer. Math. Soc. **86** (1982), 336-338. MR **83h**:54054
- [3] S. C. CHU and R. D. MOYER, On continuous functions, commuting functions, and fixed points, Fund. Math. **59** (1966), 90-95. MR **34**:805
- [4] E. M. COVEN and G. A. HEDLUND, Continuous maps of the interval whose periodic points form a closed set, Proc. Amer. Math. Soc. **79** (1980), 127-133. MR **81a**:54042

- [5] R. ENGELKING, *General topology*, Polish Scientific Publishers, Warszawa, 1977. MR **58**:18316b
- [6] J. P. HUNEKE, *Two counterexamples to a conjecture on commuting continuous functions of the closed unit interval*, Abstract no. 67T-2311, Notices Amer. Math. Soc. **14** (1967), 284.
- [7] G. JUNGCK, *Common fixed points for compatible maps on the unit interval*, Proc. Amer. Math. Soc. **115** (1992), 495-499. MR **92m**:54076

INSTITUTE OF MATHEMATICS, TECHNICAL UNIVERSITY OF ŁÓDŹ, ŻWIRKI 36, 90-924 ŁÓDŹ,
POLAND

E-mail address: jachymusk@lodz1.p.lodz.pl