ALMOST EVERYWHERE CONVERGENCE OF LACUNARY PARTIAL SUMS OF VILENKIN-FOURIER SERIES

WO-SANG YOUNG

(Communicated by J. Marshall Ash)

ABSTRACT. We prove that if $f \in L^p$, $p > 1$, and $\{n_k\}$ is any lacunary sequence of positive integers, then the sequence of n_k th partial sums of Vilenkin-Fourier series of f converges almost everywhere to f .

1. INTRODUCTION

Let $G = \prod_{i=0}^{\infty} Z_{p_i}$ be the direct product of cyclic groups of order p_i , where $\{p_i\}_{i \geq 0}$ is a sequence of integers with $p_i \geq 2$. Let μ be the Haar measure on G normalized by $\mu(G) = 1$. For $x = \{x_k\} \in G$, let $\phi_k(x) = \exp(2\pi i x_k / p_k)$, $k = 0, 1, 2, \dots$. The Vilenkin system $\{\chi_n\}$ is the set of all finite products of $\{\phi_k\}$, and is enumerated in the following manner. Let $m_0 = 1$, $m_k = \prod_{i=0}^{k-1} p_i$, $k = 1, 2, \dots$. We express each nonnegative integer n as a finite sum in the form $n = \sum_{k=0}^{\infty} \alpha_k m_k$, where $0 \leq \alpha_k < p_k$, and define $\chi_n = \prod_{k=0}^{\infty} \phi_k^{\alpha_k}$. The functions $\{\chi_n\}$ are the characters of G , and they form a complete orthonormal system on G . If $p_i = 2$, $i = 0, 1, 2, \dots$, then $\{\phi_k\}$ are the Rademacher functions and $\{\chi_n\}$ are the Walsh functions. In this paper there is no restriction on the orders $\{p_i\}$.

We consider Fourier series with respect to $\{\chi_n\}$. For $f \in L^1$, let $\widehat{f}(j) = \int_G f(t) \overline{\chi_j(t)} d\mu(t)$, $j = 0, 1, 2, \dots$, and $S_n f = \sum_{j=0}^{n-1} \widehat{f}(j) \chi_j$, $n = 1, 2, \dots$. Not much is known about the almost everywhere convergence of partial sums when there is no restriction on the orders $\{p_i\}$. We have the following result.

Theorem. *Let $1 < p < \infty$ and $\{n_k\}_{k \geq 1}$ be a lacunary sequence of positive integers, i.e., there is $q > 1$ such that $n_{k+1}/n_k \geq q$, $k = 1, 2, \dots$. Then there is a constant C_p such that*

$$(1.1) \quad \left\| \sup_k |S_{n_k} f| \right\|_p \leq C_p \|f\|_p, \quad f \in L^p.$$

It follows that $\lim_{k \rightarrow \infty} S_{n_k} f(x) = f(x)$ a.e. for all $f \in L^p$.

Received by the editors March 8, 1995 and, in revised form, June 15, 1995.

1991 *Mathematics Subject Classification.* Primary 42C10; Secondary 42B25, 43A75.

In the proof that follows, C and C_p will denote absolute constants which may vary from line to line.

2. PROOF OF THE THEOREM

We first prove (1.1). For $k = 0, 1, \dots$, let L_k be the positive integer such that $2^{L_k} \leq p_k < 2^{L_k+1}$. Since every lacunary sequence can be decomposed into a finite number of lacunary subsequences with ratio $q = 2$, we may assume, by adding more terms to the sequence if necessary, that $\{n_j\}$ can be relabelled $\{n_{k,\ell}\}_{k=0,1,\dots,\ell=0,1,\dots,L_k}$, such that for $k = 0, 1, \dots$, $2^\ell m_k \leq n_{k,\ell} < 2^{\ell+1} m_k$ if $\ell = 0, 1, \dots, L_k - 1$, and $2^{L_k} m_k \leq n_{k,L_k} < m_{k+1}$. (There is no n_{k,L_k} term if $2^{L_k} m_k = m_{k+1}$.) Also, it is sufficient to show that there is a constant C_p such that

$$(2.1) \quad \left\| \sup_{\substack{\ell=0,\dots,L_k \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f| \right\|_p \leq C_p \|f\|_p$$

for all $f \in L^p$, $N = 1, 2, \dots$.

Let $f_k = S_{m_{k+1}} f - S_{m_k} f$, $k = 0, 1, \dots$, and $f_{-1} = S_1 f$. We observe that

$$\sup_{\substack{\ell=0,\dots,L_k \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f| \leq \sup_{\substack{\ell=0,\dots,L_k \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f_k| + \sup_{k=0,\dots,N-1} |S_{m_k} f|.$$

Since $\{S_{m_k} f\}$ is a martingale (see, e.g., [5]), it follows from Doob's inequality ($\|\sup_{k \geq 0} |S_{m_k} f|\|_p \leq C_p \|f\|_p$, $f \in L^p$) that (2.1) will be proved if we have

$$(2.2) \quad \left\| \sup_{\substack{\ell=0,\dots,L_k \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f_k| \right\|_p \leq C_p \|f\|_p,$$

for all $f \in L^p$, $N = 1, 2, \dots$.

Now,

$$\begin{aligned} \sup_{\substack{\ell=0,\dots,L_k \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f_k| &\leq \left(\sum_{k=0}^{N-1} |S_{n_{k,0}} f_k|^2 \right)^{1/2} + \sup_{\substack{\ell=1,\dots,L_k-2 \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f_k| \\ &\quad + \left(\sum_{k=0}^{N-1} |S_{n_{k,L_k-1}} f_k|^2 \right)^{1/2} + \left(\sum_{k=0}^{N-1} |S_{n_{k,L_k}} f_k|^2 \right)^{1/2}. \end{aligned}$$

For each $\ell = 0, L_k - 1$ and L_k , we apply [5, Theorem 2] and Burkholder's result for martingales [1, Theorem 3.2] to get

$$\begin{aligned} \left\| \left(\sum_{k=0}^{N-1} |S_{n_{k,\ell}} f_k|^2 \right)^{1/2} \right\|_p &\leq C_p \left\| \left(\sum_{k=0}^{N-1} |f_k|^2 \right)^{1/2} \right\|_p \\ &\leq C_p \|f\|_p. \end{aligned}$$

To prove (2.2) it remains to show

$$(2.3) \quad \left\| \sup_{\substack{\ell=1,\dots,L_k-2 \\ k=0,\dots,N-1}} |S_{n_{k,\ell}} f_k| \right\|_p \leq C_p \|f\|_p$$

for all $f \in L^p$, $N = 1, 2, \dots$.

We shall use the following operators. Let $k = 0, 1, \dots$. If $L_k > 2$, define, for $\ell = 1, \dots, L_k - 2$, the sequence $\{a_{k,\ell}(n)\}_{n \geq 0}$ by

$$a_{k,\ell}(n) = \begin{cases} 1 & \text{if } 2^\ell m_k \leq n < 2^{\ell+1} m_k, \\ \frac{j}{2^{\ell-1}} & \text{if } (2^{\ell-1} + j)m_k \leq n < (2^{\ell-1} + j + 1)m_k, \\ & j = 0, 1, \dots, 2^{\ell-1} - 1, \\ 1 - \frac{j+1}{2^{\ell-1}} & \text{if } (2^{\ell+1} + j)m_k \leq n < (2^{\ell+1} + j + 1)m_k, \\ & j = 0, 1, \dots, 2^{\ell-1} - 1, \\ 0 & \text{otherwise,} \end{cases}$$

and set

$$A_{k,\ell}f = \sum_{n=0}^{\infty} a_{k,\ell}(n) \hat{f}(n) \chi_n.$$

It is proved in [6, pp. 665-666] that

$$(2.4) \quad \left\| \left(\sum_{k=0}^{N-1} \sum_{\ell=1}^{L_k-2} |A_{k,\ell}f|^2 \right)^{1/2} \right\|_p \leq C_p \|f\|_p, \quad f \in L^p, \quad N = 1, 2, \dots$$

(We interpret a sum $\sum_{j=k}^{\ell}$ with $\ell < k$ as zero.)

The second operator is defined as follows. Let $k = 0, 1, \dots$. If $L_k \geq 2$, define, for $\ell = 1, \dots, L_k - 1$, the sequence $\{b_{k,\ell}(n)\}_{n \geq 0}$ by

$$b_{k,\ell}(n) = \begin{cases} 1 & \text{if } m_k \leq n < 2^{\ell-1} m_k \text{ or } m_{k+1} - (2^{\ell-1} - 1)m_k \leq n < m_{k+1}, \\ 1 - \frac{j}{2^{\ell-1}} & \text{if } (2^{\ell-1} + j)m_k \leq n < (2^{\ell-1} + j + 1)m_k \text{ or} \\ & m_{k+1} - (2^{\ell-1} + j)m_k \leq n < m_{k+1} - (2^{\ell-1} + j - 1)m_k, \\ & j = 0, 1, \dots, 2^{\ell-1} - 1, \\ 0 & \text{otherwise.} \end{cases}$$

For $f \in L^1$, set

$$B_{k,\ell}f = \sum_{n=0}^{\infty} b_{k,\ell}(n) \hat{f}(n) \chi_n.$$

To estimate $B_{k,\ell}f$, we need the following definitions. Let $\{G_k\}$ be the sequence of subgroups of G defined by

$$G_0 = G, \quad G_k = \prod_{i=0}^{k-1} \{0\} \times \prod_{i=k}^{\infty} Z_{p_i}, \quad k = 1, 2, \dots$$

We shall identify G with the unit interval $(0, 1)$ by associating with each $\{x_i\} \in G$, $0 \leq x_i < p_i$, the point $\sum_{i=0}^{\infty} x_i m_{i+1}^{-1} \in (0, 1)$. If we disregard the countable set of p_i -rationals, this mapping is one-to-one, onto and measure preserving. On the

interval $(0, 1)$, cosets of G_k are intervals of the form $(jm_k^{-1}, (j+1)m_k^{-1})$, $j = 0, 1, \dots, m_k - 1$. A set I is called a generalized interval if $I \subset x + G_k$ for some $x \in G$, $k = 0, 1, \dots$, I is a union of cosets of G_{k+1} , and I is an interval if we consider $x + G_k$ as a circle.

Let

$$Mf(x) = \sup_{\substack{x \in I \\ I \text{ generalized} \\ \text{interval}}} \frac{1}{\mu(I)} \int_I |f| d\mu$$

be the Hardy-Littlewood maximal function for the Vilenkin system. The following pointwise estimate will be obtained in §3.

Lemma. *There is a constant C such that*

$$\sup_{\substack{\ell=1, \dots, L_k-1 \\ k=0, 1, \dots}} |B_{k,\ell}f(x)| \leq CMf(x), \quad f \in L^1, \quad x \in G.$$

To estimate $S_{n_{k,\ell}}f_k$, $\ell = 1, \dots, L_k - 2$, $k = 0, 1, \dots$, we shall use the part of $b_{k,\ell}(n)$ with $n \in [m_k, 2^\ell m_k)$. In order to get rid of the remaining part of $b_{k,\ell}(n)$, we define

$$H^N f = \sum_{k=0}^{N-1} (S_{2^{L_k-1}m_k} f - S_{m_k} f).$$

Since $H^N f = S_{m_N} f - \sum_{k=0}^{N-1} (S_{m_{k+1}} f - S_{2^{L_k-1}m_k} f)$, and it is proved in [4, Theorem 1*] that

$$\left\| \sum_{k=0}^{N-1} (S_{m_{k+1}} f - S_{2^{L_k-1}m_k} f) \right\|_p \leq C_p \|f\|_p$$

for all $f \in L^p$, $N = 1, 2, \dots$, we have

$$(2.5) \quad \|H^N f\|_p \leq C_p \|f\|_p, \quad f \in L^p, \quad N = 1, 2, \dots$$

(See also [2].)

We are now ready to prove (2.3). We observe that for $k = 0, \dots, N-1$, $\ell = 1, \dots, L_k - 2$,

$$S_{n_{k,\ell}}f_k = B_{k,\ell}(H^N f) + S_{n_{k,\ell}}(A_{k,\ell}f).$$

Hence

(2.6)

$$\sup_{\substack{\ell=1, \dots, L_k-2 \\ k=0, \dots, N-1}} |S_{n_{k,\ell}}f_k| \leq \sup_{\substack{\ell=1, \dots, L_k-2 \\ k=0, \dots, N-1}} B_{k,\ell}(H^N f) + \left(\sum_{k=0}^{N-1} \sum_{\ell=1}^{L_k-2} |S_{n_{k,\ell}}(A_{k,\ell}f)|^2 \right)^{1/2}.$$

To estimate the first term on the right, we apply the lemma, the fact that the Hardy-Littlewood maximal operator M is bounded in L^p (see [3]) and (2.5). We have

$$\begin{aligned} \left\| \sup_{\substack{\ell=1, \dots, L_k-2 \\ k=0, \dots, N-1}} |B_{k,\ell}(H^N f)| \right\|_p &\leq C \|M(H^N f)\|_p \\ &\leq C_p \|H^N f\|_p \leq C_p \|f\|_p \end{aligned}$$

for all $f \in L^p$, $N = 1, 2, \dots$. For the last term in (2.6) we use [5, Theorem 2] and (2.4). We obtain

$$\begin{aligned} \left\| \left(\sum_{k=0}^{N-1} \sum_{\ell=1}^{L_k-2} |S_{n_{k,\ell}}(A_{k,\ell}f)|^2 \right)^{1/2} \right\|_p &\leq C_p \left\| \left(\sum_{k=0}^{N-1} \sum_{\ell=1}^{L_k-2} |A_{k,\ell}f|^2 \right)^{1/2} \right\|_p \\ &\leq C_p \|f\|_p \end{aligned}$$

for all $f \in L^p$, $N = 1, 2, \dots$. This proves (2.3). The proof of (1.1) will be complete once we prove the lemma.

Finally, if $f \in L^p$, $\lim_{k \rightarrow \infty} \|S_{m_k}f - f\|_p = 0$ since $S_{m_k}f$ is the average of f over the cosets of G_k . As a consequence of this and (1.1), we have $\lim_{k \rightarrow \infty} S_{n_k}f(x) = f(x)$ a.e.

3. PROOF OF THE LEMMA

We shall use the following notation. For each generalized interval I , we define the generalized interval $3I$ as follows. If $I = G$, let $3I = G$. Suppose I is a proper subset of $x + G_k$, $x \in G$, $k = 0, 1, 2, \dots$, and is the union of cosets of G_{k+1} . If $\mu(I) \geq \mu(G_k)/3$, let $3I = x + G_k$. If $\mu(I) < \mu(G_k)/3$, consider $x + G_k$ as a circle and define $3I$ to be the interval in this circle which has the same center as I and has measure $\mu(3I) = 3\mu(I)$. For all cases, we have $\mu(3I) \leq 3\mu(I)$.

For $k = 0, 1, \dots$, $\ell = 1, \dots, L_k - 1$ and $f \in L^1$, we have

$$B_{k,\ell}f(x) = \int_G f(t) \left[\sum_{n=0}^{\infty} b_{k,\ell}(n) \chi_n(x-t) \right] d\mu(t).$$

Since $b_{k,\ell}(n)$ vanishes for $n \notin [m_k, m_{k+1})$ and is constant for $n \in [\alpha m_k, (\alpha+1)m_k)$, $\alpha = 0, 1, \dots$, we get

$$\begin{aligned} \sum_{n=0}^{\infty} b_{k,\ell}(n) \chi_n &= \sum_{\alpha=1}^{p_k-1} b_{k,\ell}(\alpha m_k) \sum_{n=\alpha m_k}^{(\alpha+1)m_k-1} \chi_n \\ &= \sum_{\alpha=1}^{p_k-1} b_{k,\ell}(\alpha m_k) \phi_k^\alpha D_{m_k}, \end{aligned}$$

where $D_n = \sum_{j=0}^{n-1} \chi_j$, $n = 1, 2, \dots$, denotes the n^{th} Dirichlet kernel. Since $D_{m_k} = \mu(G_k)^{-1} \chi_{G_k}$, we have

$$(3.1) \quad B_{k,\ell}f(x) = \frac{1}{\mu(G_k)} \int_{x+G_k} f(t) M_{k,\ell}(x-t) d\mu(t),$$

where

$$\begin{aligned} M_{k,\ell} &= \sum_{\alpha=1}^{p_k-1} b_{k,\ell}(\alpha m_k) \phi_k^\alpha \\ &= \sum_{\alpha=1}^{2^\ell-1} b_{k,\ell}(\alpha m_k) \phi_k^\alpha + \phi_k^0 + \sum_{\alpha=-(2^\ell-1)}^{-1} b_{k,\ell}((p_k+\alpha)m_k) \phi_k^\alpha - \phi_k^0. \end{aligned}$$

Let

$$D_{k,j} = \sum_{\alpha=-j}^j \phi_k^\alpha, \quad j = 0, 1, \dots, 2^{L_k-1} - 1,$$

and

$$K_{k,n} = \frac{1}{n} \sum_{j=0}^{n-1} D_{k,j}, \quad n = 1, \dots, 2^{L_k-1}.$$

Then

$$\begin{aligned} (3.2) \quad M_{k,\ell} &= \frac{1}{2^{\ell-1}} (D_{k,2^{\ell-1}} + D_{k,2^{\ell-1}+1} + \dots + D_{k,2^\ell-1}) - \phi_k^0 \\ &= 2K_{k,2^\ell} - K_{k,2^{\ell-1}} - \phi_k^0. \end{aligned}$$

For $n = 1, \dots, 2^{L_k-1}$, let

$$\sigma_{k,n}f(x) = \frac{1}{\mu(G_k)} \int_{x+G_k} f(t) K_{k,n}(x-t) d\mu(t).$$

From (3.1) and (3.2) we obtain

$$|B_{k,\ell}f(x)| \leq 2|\sigma_{k,2^\ell}f(x)| + |\sigma_{k,2^{\ell-1}}f(x)| + Mf(x).$$

The lemma will be proved if we show

$$(3.3) \quad |\sigma_{k,n}f(x)| \leq C Mf(x),$$

for all $k = 0, 1, \dots$, $n = 1, \dots, 2^{L_k-1}$, $x \in G$.

By a direct computation we have

$$K_{k,n}(x) = \begin{cases} \frac{1}{n} \left(\sin^2 \frac{\pi n x_k}{p_k} / \sin^2 \frac{\pi x_k}{p_k} \right) & \text{if } x_k \neq 0, \\ n & \text{if } x_k = 0, \end{cases}$$

and hence

$$(3.4) \quad 0 \leq K_{k,n}(x) \leq \min \left\{ \left(n \sin^2 \frac{\pi x_k}{p_k} \right)^{-1}, n \right\}.$$

Let I be a generalized interval containing x such that $I \subset x + G_k$, I is a union of cosets of G_{k+1} and

$$\frac{p_k}{n} - 1 < \frac{\mu(I)}{\mu(G_{k+1})} \leq \frac{p_k}{n}.$$

(Note that $p_k/n \geq 2$ since $n \leq 2^{L_k-1}$.) For $j = 1, 2, \dots$, let $3^{j+1}I = 3(3^jI)$, and $J = \min\{j \geq 1 : 3^jI = x + G_k\}$. Then

$$\begin{aligned} (3.5) \quad |\sigma_{k,n}f(x)| &\leq \frac{1}{\mu(G_k)} \int_{3I} |f(t)| K_{k,n}(x-t) d\mu(t) \\ &\quad + \sum_{j=1}^{J-1} \frac{1}{\mu(G_k)} \int_{3^{j+1}I \setminus 3^jI} |f(t)| K_{k,n}(x-t) d\mu(t). \end{aligned}$$

By (3.4),

$$\frac{1}{\mu(G_k)} \int_{3I} |f(t)| K_{k,n}(x-t) d\mu(t) \leq \frac{3\mu(I)}{\mu(G_k)} n \frac{1}{\mu(3I)} \int_{3I} |f(t)| d\mu(t) \leq CM f(x).$$

To estimate the last term in (3.5), we apply the other estimate in (3.4). We observe that when $x \in I$, $t \notin 3^j I$, $j = 1, \dots, J-1$,

$$\left| \sin \frac{\pi(x_k - t_k)}{p_k} \right| \geq C\mu(3^{j-1}I)/\mu(G_k).$$

Hence

$$\begin{aligned} & \frac{1}{\mu(G_k)} \int_{3^{j+1}I \setminus 3^j I} |f(t)| K_{k,n}(x-t) d\mu(t) \\ & \leq \frac{\mu(3^{j+1}I)}{\mu(G_k)} \frac{1}{\mu(3^{j+1}I)} \int_{3^{j+1}I \setminus 3^j I} |f(t)| \left(n \sin^2 \frac{\pi(x_k - t_k)}{p_k} \right)^{-1} d\mu(t) \\ & \leq C \frac{\mu(3^{j+1}I)}{\mu(G_k)} \frac{1}{n} \left[\frac{\mu(G_k)}{\mu(3^{j-1}I)} \right]^2 \frac{1}{\mu(3^{j+1}I)} \int_{3^{j+1}I} |f(t)| d\mu(t) \\ & \leq C 3^{-j} \frac{\mu(G_k)}{n\mu(I)} Mf(x) \\ & \leq C 3^{-j} Mf(x). \end{aligned}$$

Substituting these estimates into (3.5) and summing over j , we obtain (3.3). The lemma follows.

This completes the proof of the theorem.

REFERENCES

1. D.L. Burkholder, *Distribution function inequalities for martingales*, Ann. Probab. **1** (1973), 19–42. MR **51**:1944
2. P. Simon, *On the concept of a conjugate function*, Colloq. Math. Soc. J. Bolyai, Fourier Analysis and Approximation Theory, 1976 **19** (1978), 747–755. MR **81b**:42084
3. P. Simon, *On a maximal function*, Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **21** (1978), 41–44. MR **81b**:42083
4. W.-S. Young, *Mean convergence of generalized Walsh-Fourier series*, Trans. Amer. Math. Soc. **218** (1976), 311–320. MR **52**:14828
5. W.-S. Young, *Almost everywhere convergence of Vilenkin-Fourier series of H^1 -functions*, Proc. Amer. Math. Soc. **108** (1990), 433–441. MR **90g**:42057
6. W.-S. Young, *Littlewood-Paley and multiplier theorems for Vilenkin-Fourier series*, Canad. J. Math. **46** (1994), 662–672. MR **95c**:42031

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA
T6G 2G1