AN ABSTRACT ERGODIC THEOREM AND SOME INEQUALITIES FOR OPERATORS ON BANACH SPACES

YUAN-CHUAN LI AND SEN-YEN SHAW

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We prove an abstract mean ergodic theorem and use it to show that if $\{A_n\}$ is a sequence of commuting m -dissipative (or normal) operators on a Banach space X , then the intersection of their null spaces is orthogonal to the linear span of their ranges. It is also proved that the inequality $\|x + Ay\| \geq \|x\| - 2\sqrt{\|Ax\| \|y\|}$ ($x, y \in D(A)$) holds for any m -dissipative operator A . These results either generalize or improve the corresponding results of Shaw, Mattila, and Crabb and Sinclair, respectively.

1. INTRODUCTION

Let X be a real (or complex) Banach space and $B(X)$ the Banach algebra of all bounded linear operators on X . Given a family $\mathbf{A}$ of closed linear operators on X , a net $\{A_\alpha\}$ in $B(X)$ will be called an **A-ergodic net** if the following conditions hold:

- (a) There is an $M > 0$ such that $\|A_\alpha\| \leq M$ for all α ;
- (b) $\|(A_\alpha - I)x\| \rightarrow 0$ for all $x \in \bigcap_{A \in \mathbf{A}} N(A)$ and $R(A_\alpha - I) \subset \overline{\sum_{A \in \mathbf{A}} R(A)}$ eventually α ;
- (c) For every $A \in \mathbf{A}$, $R(A_\alpha) \subset D(A)$ eventually α and $w\text{-}\lim_\alpha AA_\alpha x = 0$ for all $x \in X$, and $\|A_\alpha Ax\| \rightarrow 0$ for all $x \in D(A)$.

Note that **A-ergodic nets** with $\mathbf{A} = \{T - I; T \in \mathbf{S}\}$ for some set $\mathbf{S} \subset B(X)$ were first studied by Eberlein [8]; such an operator net is named a right, weakly left **S-ergodic net** in [12, p. 75].

An abstract mean ergodic theorem from [17, Theorem 1.1] asserts that, for an **A-ergodic net** with $\mathbf{A}$ consisting of a single closed operator A , the operator P , defined by

$$\begin{cases} D(P) := \{x \in X; s\text{-}\lim_\alpha A_\alpha x \text{ exists}\}, \\ Px = s\text{-}\lim_\alpha A_\alpha x, \quad x \in D(P), \end{cases}$$

Received by the editors February 14, 1995 and, in revised form, May 18, 1995.

1991 *Mathematics Subject Classification.* Primary 47A35, 47B15, 47B44.

Key words and phrases. Abstract mean ergodic theorem, hermitian operator, hyponormal operator, m -dissipative operator, normal operator, orthogonality, strictly c -convex space.

This research was supported in part by the National Science Council of the R.O.C.

is a bounded linear projection with norm $\|P\| \leq M$, range $R(P) = N(A)$, and null space $N(P) = \overline{R(A)}$. Applications of it to studies of ergodic properties of many particular operator families, such as integrated semigroups, cosine operator functions, and tensor product semigroups were also discussed in [18] and [19]. The purposes of this paper are: (1) to extend the above ergodic theorem to the general case that $\mathbf{A}$ is not a singleton; (2) to deduce from the generalized ergodic theorem Dunford's ergodic theorem [7] for multi-parameter semigroups; (3) using the abstract ergodic theorem to investigate orthogonality properties of some operators.

The generalized abstract mean ergodic theorem states as follows.

Theorem 1. *Let $\mathbf{A}$ be a family of closed linear operators on X , and let $\{A_\alpha\}$ be an $\mathbf{A}$ -ergodic net. Then the operator P is a linear projection with norm $\|P\| \leq M$, range $R(P) = \bigcap_{A \in \mathbf{A}} N(A)$, null space $N(P) = \overline{\sum_{A \in \mathbf{A}} R(A)}$, and domain*

$$\begin{aligned} D(P) &= \bigcap_{A \in \mathbf{A}} N(A) \oplus \overline{\sum_{A \in \mathbf{A}} R(A)} \\ &= \{x \in X; \{A_\alpha x\} \text{ has a weak cluster point}\}. \end{aligned}$$

Here $\sum_{A \in \mathbf{A}} R(A)$ denotes the linear space spanned by the spaces $R(A)$, $A \in \mathbf{A}$. If the space X is reflexive, then $D(P) = X$.

The proof of Theorem 1 and some consequences including an application to multi-parameter semigroups will be given in Section 2. Section 3 will be concerned with applications to orthogonality properties of m -dissipative operators and normal operators on Banach spaces.

We recall some necessary definitions. Let X be a complex Banach space. The (spatial) numerical range of a linear operator $A : D(A) \subset X \rightarrow X$ is the set $W(A) := \{\langle Ax, x^* \rangle; x \in D(A), x^* \in X^*, \|x\| = \|x^*\| = \langle x, x^* \rangle = 1\}$. An operator $H \in B(X)$ is said to be *hermitian* if its numerical range $W(H)$ is contained in the real line R , or equivalently, if $\|\exp(itH)\| = 1$ for all $t \in R$ (see [2], [3]). H is said to be *positive* if $W(H) \subset [0, \infty)$. If an operator T can be expressed as $T = H + iK$ with H and K hermitian, then $i(HK - KH)$ is hermitian. T is called *normal* if $T = H + iK$ for some commuting hermitian operators H and K . These definitions generalize those familiar concepts of operators on Hilbert spaces. An operator $A : D(A) \subset X \rightarrow X$ is called *dissipative* if its numerical range $W(A)$ is contained in the half plane $\{z \in C, \operatorname{Re} z \leq 0\}$. A dissipative operator A is called *m -dissipative* if $\rho(A) \cap (0, \infty) \neq \emptyset$.

Mattila [16] and Fong [9] proved that the null space $N(T)$ of a normal operator $T \in B(X)$ is *orthogonal* to the range $R(T)$ in the sense that $\|x + Ty\| \geq \|x\|$ for all $x \in N(T)$ and all $y \in X$. In particular, a scalar multiple of a hermitian operator has this orthogonality property. This special case is readily contained in a theorem of Crabb and Sinclair [3, Theorem 20.6], which says that if 0 is not an interior point of the closed convex hull $\overline{\operatorname{co}} W(T)$ of the numerical range $W(T)$ of a bounded operator T (in other words, T is a scalar multiple of a bounded m -dissipative operator), then $\|x + Ty\| \geq \|x\| - \sqrt{8\|Tx\|\|y\|}$ for all $x, y \in X$.

In Section 3 we establish the following generalizations or improvements of the above known results:

(1) Let $\{A_1, A_2, \dots\}$ be a countable family of m -dissipative operators or normal operators on X . If the resolvents of the A_k 's are commutative, then $\bigcap_{k=1}^{\infty} N(A_k)$ is orthogonal to the linear span $\sum_{k=1}^{\infty} R(A_k)$ of $\{R(A_k)\}$ (Theorem 7, Corollary 8).

(2) If A is an m -dissipative operator, then $\|x + Ay\| \geq \|x\| - 2\sqrt{\|Ax\| \|y\|}$ for $x, y \in D(A)$ (Theorem 9).

(3) If 0 is not an interior point of $\overline{\text{co}} W(T)$, $T \in B(X)$, then $\|x + Ty\| \geq \|x\| - 2\sqrt{\|Tx\| \|y\|}$ for all $x, y \in X$, and $\|T\|^2 \leq 4\|T^2\|$ (Corollary 10, Corollary 11).

2. PROOF OF THEOREM 1 AND APPLICATIONS TO PRODUCTS OF A -ERGODIC NETS

Proof of Theorem 1. Clearly, (a) implies that P is a bounded linear operator with $\|P\| \leq M$, and both $D(P)$ and $N(P)$ are closed. The first part of (c) and the closedness of A imply that $R(P) \subset N(A)$ for all $A \in \mathbf{A}$. Hence we have $R(P) \subset \bigcap_{A \in \mathbf{A}} N(A)$. Conversely, if $y \in \bigcap_{A \in \mathbf{A}} N(A)$, then by the first part of (b) we have $y \in D(P)$ and $Py = y$. Therefore $R(P) = \bigcap_{A \in \mathbf{A}} N(A)$. Moreover, since $Px \in \bigcap_{A \in \mathbf{A}} N(A)$, one has $PPx = Px$ for all $x \in D(P)$, i.e. P is a projection.

Next, we show that $N(P) = \overline{\sum_{A \in \mathbf{A}} R(A)}$. The second part of (c) implies that $R(A) \subset N(P)$ for all $A \in \mathbf{A}$, so that $\sum_{A \in \mathbf{A}} R(A) \subset N(P)$. Conversely, if $x \in N(P)$, the second part of (b) implies that

$$x = s\text{-}\lim_{\alpha} (I - A_{\alpha})x \in \overline{\sum_{A \in \mathbf{A}} R(A)}.$$

Therefore $N(P) = \overline{\sum_{A \in \mathbf{A}} R(A)}$.

Finally, if $x \in X$ is such that $\{A_{\alpha}x\}$ has a weakly convergent subnet $\{A_{\beta}x\}$, say $y = w\text{-}\lim_{\beta} A_{\beta}x$, it follows from the first part of (c) and the closedness of each $A \in \mathbf{A}$ that $y \in D(A)$ and $Ay = w\text{-}\lim_{\beta} AA_{\beta}x = 0$. That is, $y \in \bigcap_{A \in \mathbf{A}} N(A) = R(P)$. On the other hand, the second part of (b) implies that

$$y - x = w\text{-}\lim_{\beta} (A_{\beta} - I)x \in \overline{\sum_{A \in \mathbf{A}} R(A)} = N(P).$$

Hence $x = y - (y - x) \in R(P) \oplus N(P) = D(P)$, and the proof has been completed.

From Theorem 1 we now deduce some ergodic theorems for (countable) products of A -ergodic nets.

Corollary 2. *Let $\{A^{(k)}\}$ be a sequence of closed linear operators on X . For each k , let $\{A_{\alpha}^{(k)}\}$ be an $A^{(k)}$ -ergodic net satisfying*

(a') $M := \sup\{\|\prod_{k \in F} A_{\alpha}^{(k)}\|; F \text{ a finite set of natural numbers and for all } \alpha\} < \infty$;

(b') $\|(A_{\alpha}^{(k)} - I)x\| \rightarrow 0$ for all $x \in N(A^{(k)})$ and $R(A_{\alpha}^{(k)} - I) \subset \overline{R(A^{(k)})}$ for all α ;

(c') $R(A_{\alpha}^{(k)}) \subset D(A^{(k)})$ and $A_{\alpha}^{(k)} A^{(k)} \subset A^{(k)} A_{\alpha}^{(k)}$ for all α , and $\|A^{(k)} A_{\alpha}^{(k)}\| \rightarrow 0$.

(d') $A_{\alpha}^{(j)} A_{\alpha}^{(k)} = A_{\alpha}^{(k)} A_{\alpha}^{(j)}$ for all α, j , and k .

For a nondecreasing net $\{r_{\alpha}\}$ of positive integers, let $P : D(P) \subset X \rightarrow X$ be the operator defined by

$$\begin{cases} D(P) := \{x \in X; s\text{-}\lim_{\alpha} A_{\alpha}^{(1)} A_{\alpha}^{(2)} \cdots A_{\alpha}^{(r_{\alpha})} x \text{ exists}\}, \\ Px = s\text{-}\lim_{\alpha} A_{\alpha}^{(1)} A_{\alpha}^{(2)} \cdots A_{\alpha}^{(r_{\alpha})} x, x \in D(P). \end{cases}$$

Then P is a linear projection with norm $\|P\| \leq M$, range $R(P) = \bigcap_{k=1}^{\infty} N(A^{(k)})$, null space $N(P) = \overline{\sum_{k=1}^{\infty} A^{(k)}}$, and domain

$$D(P) = \{x \in X; \{A_{\alpha}^{(1)} A_{\alpha}^{(2)} \cdots A_{\alpha}^{(r_{\alpha})} x\} \text{ has a weak cluster point}\}.$$

Proof. Define $A_\alpha := A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(r_\alpha)}$ for all α . It suffices to check that the net $\{A_\alpha\}$ is an $\mathbf{A}$ -ergodic net with $\mathbf{A} = \{A^{(k)}\}$, i.e. (a)–(c) of Theorem 1 are satisfied. (a') implies (a). For every $x \in X$ and α , we have, by the commutativity of $A_\alpha^{(j)}$ and $A_\alpha^{(k)}$, that

$$\begin{aligned} (A_\alpha - I)x &= A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(r_\alpha)} x - x \\ &= \sum_{k=2}^{r_\alpha} A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(k-1)} (A_\alpha^{(k)} - I)x + (A_\alpha^{(1)} - I)x \\ &= \sum_{k=2}^{r_\alpha} (A_\alpha^{(k)} - I) A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(k-1)} x + (A_\alpha^{(1)} - I)x, \end{aligned}$$

from which and (a') it is seen that parts 1 and 2 of (b) follow respectively from the corresponding parts of (b').

Also we have for every k

$$A_\alpha = A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(r_\alpha)} = A_\alpha^{(k)} \prod_{1 \leq j \neq k \leq r_\alpha} A_\alpha^{(j)} = \left\{ \prod_{1 \leq j \neq k \leq r_\alpha} A_\alpha^{(j)} \right\} A_\alpha^{(k)},$$

which and (c') imply that $R(A_\alpha) \subset D(A^{(k)})$,

$$\begin{aligned} \|A^{(k)} A_\alpha\| &\leq \|A^{(k)} A_\alpha^{(k)}\| \left\| \prod_{1 \leq j \neq k \leq r_\alpha} A_\alpha^{(j)} \right\| \rightarrow 0, \\ \|A_\alpha A^{(k)} x\| &\leq \left\| \prod_{1 \leq j \neq k \leq r_\alpha} A_\alpha^{(j)} \right\| \|A_\alpha^{(k)} A^{(k)}\| \|x\| \rightarrow 0, \quad x \in D(A^{(k)}). \end{aligned}$$

Therefore $\{A_\alpha\}$ is an $\mathbf{A}$ -ergodic net, and the conclusion now follows from Theorem 1.

In the special cases $r_\alpha \equiv m$ and $r_n = n$, Corollary 2 becomes the following two corollaries.

Corollary 3. *Let $\{A^{(k)}\}$, $k = 1, 2, \dots, m$, be closed linear operators on X . For each k , let $\{A_\alpha^{(k)}\}$ be an $A^{(k)}$ -ergodic net satisfying conditions (b'), (c'), and (d'). Let $P : D(P) \subset X \rightarrow X$ be the operator defined by*

$$\begin{cases} D(P) := \{x \in X; s\text{-}\lim_\alpha A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(m)} x \text{ exists}\}, \\ Px = s\text{-}\lim_\alpha A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(m)} x, \quad x \in D(P). \end{cases}$$

Suppose that $\|A_\alpha^{(k)}\| \leq M_k$ for all $1 \leq k \leq m$. Then P is a linear projection with norm $\|P\| \leq M_1 \cdots M_m$, range $R(P) = \bigcap_{k=1}^m N(A^{(k)})$, null space $N(P) = \overline{\sum_{k=1}^m A^{(k)}}$, and domain

$$D(P) = \{x \in X; \{A_\alpha^{(1)} A_\alpha^{(2)} \cdots A_\alpha^{(m)} x\} \text{ has a weak cluster point}\}.$$

Corollary 4. *Let $\{A^{(k)}\}$, $k = 1, 2, \dots$, be a sequence of closed linear operators on X . For each k , let $\{A_n^{(k)}\}$ be an $A^{(k)}$ -ergodic sequence satisfying (b'), (c'), (d') (with α replaced by n). Let $P : D(P) \subset X \rightarrow X$ be the operator defined by*

$$\begin{cases} D(P) := \{x \in X; s\text{-}\lim_n A_n^{(1)} A_n^{(2)} \cdots A_n^{(n)} x \text{ exists}\}, \\ Px = s\text{-}\lim_n A_n^{(1)} A_n^{(2)} \cdots A_n^{(n)} x, \quad x \in D(P). \end{cases}$$

Suppose that $M := \sup\{\|\prod_{k \in F} A_n^{(k)}\|; F \text{ a finite set of natural numbers and } n \geq 1\} < \infty$. Then P is a linear projection with norm $\|P\| \leq M$, range $R(P) = \bigcap_{k=1}^{\infty} N(A^{(k)})$, null space $N(P) = \overline{\sum_{k=1}^{\infty} A^{(k)}}$, and domain

$$D(P) = \{x \in X; \{A_n^{(1)} A_n^{(2)} \cdots A_n^{(n)} x\} \text{ has a weak cluster point}\}.$$

Next, we give two illustrative examples.

Example 1. Let $A^{(1)}, A^{(2)}, \dots$ be a sequence of closed operators on a Banach space X satisfying $(0, \infty) \subset \rho(A^{(k)})$, $(\lambda - A^{(j)})^{-1}(\lambda - A^{(k)})^{-1} = (\lambda - A^{(k)})^{-1}(\lambda - A^{(j)})^{-1}$, and $\|\lambda(\lambda - A^{(k)})^{-1}\| \leq M_k$ for all $\lambda > 0$ and $j, k = 1, 2, \dots$. Suppose further that $M = \sup\{\prod_{k \in F} M_k; F \text{ a finite set of natural numbers}\} < \infty$ (particularly, $M_k = 1$ for all k). If we put $A_n^{(k)} := \frac{1}{n}(\frac{1}{n} - A^{(k)})^{-1}$ for $n \geq 1$, then we have $A_n^{(k)} A^{(k)} \subset A^{(k)} A_n^{(k)} = \frac{1}{n}(A_n^{(k)} - I)$, so that conditions in Corollary 4 are satisfied. Hence we can formulate the next theorem.

Theorem 5. Under the above assumption on closed operators $A^{(1)}, A^{(2)}, \dots$, the limits $\lim_{n \rightarrow \infty} n^{-n}(\frac{1}{n} - A^{(1)})^{-1} \cdots (\frac{1}{n} - A^{(n)})^{-1}x$ define a linear projection P with norm $\|P\| \leq M$, range $R(P) = \bigcap_{k=1}^{\infty} N(A^{(k)})$, null space $N(P) = \overline{\sum_{k=1}^{\infty} R(A^{(k)})}$, and domain

$$\begin{aligned} D(P) &= \left[\bigcap_{k=1}^{\infty} N(A^{(k)}) \right] \oplus \overline{\sum_{k=1}^{\infty} R(A^{(k)})} \\ &= \left\{ x \in X; \exists n_k \rightarrow \infty \ni w\text{-}\lim_{k \rightarrow \infty} n_k^{-n_k} \left(\frac{1}{n_k} - A^{(1)} \right)^{-1} \right. \\ &\quad \left. \cdots \left(\frac{1}{n_k} - A^{(n_k)} \right)^{-1} x \text{ exists} \right\}. \end{aligned}$$

Example 2. For α, β in Euclidean N -space R^N , $\alpha > \beta$ means that $\alpha_1 > \beta_1, \dots, \alpha_N > \beta_N$; and $\alpha \rightarrow \infty$ means that $\alpha_1 \rightarrow \infty, \dots, \alpha_N \rightarrow \infty$. Let $\{T(t); t \in R^N, t > 0\} \subset B(X)$ be a strongly continuous semigroup (see [11]) such that $\|T(u)\| \leq M$ for all $0 < u \in R^N$. The averages $A(\alpha)$; $\alpha = (a, \dots, a)$, are defined in terms of $T(u)$ and the N -dimensional interval $\sigma(\alpha) \equiv (0, a]^N$, by the equation

$$A(\alpha)x = \frac{1}{a^N} \int_{\sigma(\alpha)} T(u)x \, du, \quad x \in X.$$

Let $\mathbf{A} = \{T(t) - I; t \in R^N, t > 0\}$.

Since $(A(\alpha) - I)x = \frac{1}{a^N} \int_{\sigma(\alpha)} (T(u) - I)x \, du$, $\{A(\alpha)\}$ satisfies condition (b). Since

$$(T(t) - I)A(\alpha)x = \frac{1}{a^N} \left[\int_{\sigma(\alpha)' \cap (t + \sigma(\alpha))} T(u)x \, du - \int_{\sigma(\alpha) \cap (t + \sigma(\alpha))'} T(u)x \, du \right],$$

and since the measures of the sets $\sigma(\alpha)' \cap (t + \sigma(\alpha))$ and $\sigma(\alpha) \cap (t + \sigma(\alpha))'$ are both $O(a^{N-1})$ as $\alpha \rightarrow \infty$ (see [7]), we have that $\|(T(t) - I)A(\alpha)x\| \rightarrow 0$ as $\alpha \rightarrow \infty$. Hence $\{A(\alpha)\}$ is an $\mathbf{A}$ -ergodic net.

In case the semigroup $T(\cdot)$ is strongly continuous on $\{\alpha \in R^N; \alpha_k \geq 0, 0 \leq k \leq N\}$ (i.e. an N -parameter C_0 -semigroup), the families $T_k(\cdot) := \{T(te_k); 0 \leq t < \infty\}$, $k = 1, \dots, N$, are commuting one-parameter C_0 -semigroups. Let A_k be the infinitesimal generator of $T_k(\cdot)$. It can be verified that $\{A(\alpha)\}$ is an $\mathbf{A}$ -ergodic net with $\mathbf{A} = \{A_1, A_2, \dots, A_N\}$.

Now we can deduce the following theorem from Theorem 1.

Theorem 6. *Let $\{T(t); t \in \mathbb{R}^N, t > 0\}$ be a uniformly bounded, strongly continuous semigroup. The operator $P : x \rightarrow \lim_{\alpha \rightarrow \infty} A(\alpha)x$ is a bounded linear projection with range $R(P) = \bigcap_{0 < u \in \mathbb{R}^N} N(T(u) - I)$, null space $N(P) = \overline{\bigcup_{0 < u \in \mathbb{R}^N} R(T(u) - I)}$, and domain*

$$D(P) = \{x \in X; \{A(\alpha)x\} \text{ has a weak cluster point}\}.$$

In case that $T(\cdot)$ is a uniformly bounded N -parameter C_0 -semigroup, one also has $R(P) = \bigcap_{k=1}^N N(A_k)$ and $N(P) = \overline{\sum_{k=1}^N R(A_k)}$.

3. ORTHOGONALITY PROPERTIES OF SOME OPERATORS

Let Y and Z be two subspaces of the Banach space X . We say that Y is *orthogonal* to Z , and denote this by $Y \perp Z$, if $\|y + z\| \geq \|y\|$ for all $y \in Y$ and $z \in Z$. This definition of *orthogonality* is consistent with the usual concept of orthogonality in Hilbert spaces, and is equivalent to $Y \cap Z = \{0\}$ and the projection P onto Y along Z has norm $\|P\| = 1$.

Theorem 7. *Let $\{A_n\}$ be a sequence of m -dissipative operators such that their resolvents are commutative. Then*

$$(3.1) \quad \left[\bigcap_{k=1}^{\infty} N(A_k) \right] \perp \overline{\sum_{k=1}^{\infty} R(A_k)}.$$

If the space X is reflexive, then

$$(3.2) \quad X = \left[\bigcap_{k=1}^{\infty} N(A_k) \right] \perp \overline{\sum_{k=1}^{\infty} R(A_k)}.$$

Proof. Since an operator A is dissipative if and only if for each $\lambda > 0$, $\lambda - A$ is injective and $\|\lambda(\lambda - A)^{-1}\| \leq 1$ (see [10, p. 26]), the theorem follows immediately from Theorem 5. Noting that A and $\lambda(\lambda - A)^{-1} - I$ ($\lambda > 0$) have the same null space and range, one can also deduce the theorem directly from Corollary 4 by setting $A^{(k)} = \lambda(\lambda - A_k)^{-1} - I$ and $A_n^{(k)} = \frac{1}{n} \sum_{j=0}^{n-1} (\lambda(\lambda - A_k)^{-1})^j$.

The conclusion of Theorem 7 holds in particular for any sequence of commutative hermitian operators, because i times a hermitian operator is m -dissipative. The next corollary shows that normal operators share the same property. For the proof of it we need the generalized Fuglede theorem (see [5], [6], [9]), which states that if $T = H + iK$ is a normal operator, where H and K are commuting hermitian operators, then $N(T) = N(H) \cap N(K)$. Applying this to the normal derivation $\Delta_{T,T} = \Delta_{H,H} + i\Delta_{K,K}$ (see [1], [13], [17]), one obtains the Fuglede theorem for normal operators on Banach spaces, that is, $TB = BT$ if and only if $HB = BH$ and $KB = BK$.

Corollary 8. *Let $\{T_n\}$ be a sequence of commuting normal operators on X . Then*

$$\left[\bigcap_{k=1}^{\infty} N(T_k) \right] \perp \overline{\sum_{k=1}^{\infty} R(T_k)}.$$

If the space X is reflexive, then

$$(3.3) \quad X = \left[\bigcap_{k=1}^{\infty} N(T_k) \right] \overset{\perp}{\oplus} \overline{\sum_{k=1}^{\infty} R(T_k)}.$$

Proof. Since normal operators $T_k = H_k + iK_k$, $k = 1, 2, \dots$, are commutative, the above observation shows that the hermitian operators $H_1, H_2, \dots, K_1, K_2, \dots$, are commutative. It follows from Theorem 7 that

$$\left\{ \bigcap_{k=1}^{\infty} [N(H_k) \cap N(K_k)] \right\} \perp \overline{\sum_{k=1}^{\infty} [R(H_k) + R(K_k)]},$$

and in case X is reflexive,

$$X = \left\{ \bigcap_{k=1}^{\infty} [N(H_k) \cap N(K_k)] \right\} \overset{\perp}{\oplus} \overline{\sum_{k=1}^{\infty} [R(H_k) + R(K_k)]}.$$

Since $N(T_k) = N(H_k) \cap N(K_k)$ and $R(T_k) \subset R(H_k) + R(K_k)$, we have proved the first part of the corollary. In particular, $X_0 := [\bigcap_{k=1}^{\infty} N(T_k)] \overset{\perp}{\oplus} \overline{\sum_{k=1}^{\infty} R(T_k)} = \overset{\perp}{\oplus} [\sum_{k=1}^{\infty} R(T_k^*)] \overset{\perp}{\oplus} \overline{\sum_{k=1}^{\infty} R(T_k)}$ is a closed linear subspace of X .

Since the dual operators T_k^* are also normal operators, the first part of this theorem implies that

$$\left[\bigcap_{k=1}^{\infty} N(T_k^*) \right] \perp \overline{\sum_{k=1}^{\infty} R(T_k^*)}.$$

If X is reflexive, then $X_0^\perp = \overline{\sum_{k=1}^{\infty} R(T_k^*)} \cap [\bigcap_{k=1}^{\infty} N(T_k^*)] = \{0\}$. Hence $X_0 = X$, i.e. (3.3) holds.

Remarks. (1) In the case of bounded operators, Theorem 7 and Corollary 8 both follow also from [14, Theorem 3.3] and [15, Theorem 2].

(2) If X is smooth, i.e. the norm of X is Gâteaux differentiable, then, because $x \perp y$ and $x \perp z$ imply $x \perp y+z$, the first part of Theorem 7 and that of Corollary 8 follow immediately from the case of a single operator, even when the resolvents of the concerned family of operators are not commutative.

(3) Mattila [15] showed that if $T = H + iK$ is a hyponormal operator on a strictly c -convex Banach space, then $N(T) = N(H) \cap N(K)$. Thus when X is a smooth and strictly c -convex Banach space (e.g. L_p , $1 < p < \infty$, or Hilbert spaces), the first part of Corollary 8 also holds for noncommutative hyponormal operators.

Theorem 7 implies that if A is an m -dissipative operator, then $\|x + Ay\| \geq \|x\|$ for all $x \in N(A)$ and $y \in D(A)$. The next theorem gives a generalized inequality.

Theorem 9. *If A is an m -dissipative operator in a Banach space X , then*

$$(3.4) \quad \|x + Ay\| \geq \|x\| - 2\sqrt{\|Ax\| \|y\|} \quad \text{for all } x, y \in D(A).$$

Proof. For each $n \geq 1$ we define an operator function $T_n(\cdot)$ by

$$T_n(t)x := t^{-1} \int_0^t (I - \frac{s}{n}A)^{-n} x \, ds, \quad x \in X, t > 0.$$

The m -dissipativity of A implies $\|T_n(t)\| \leq 1$ for all n and t . Then we have for $x, y \in D(A)$, $n \geq 2$ and $t > 0$

$$\begin{aligned}
\|x + Ay\| &\geq \|T_n(t)(x + Ay)\| \\
&\geq \|x\| - \|(T_n(t) - I)x + T_n(t)Ay\| \\
&\geq \|x\| - \left\| \frac{1}{t} \int_0^t \left[\left(I - \frac{s}{n} A \right)^{-n} - I \right] x \, ds \right\| - \left\| \frac{1}{t} \int_0^t \left(I - \frac{s}{n} A \right)^{-n} Ay \, ds \right\| \\
&= \|x\| - \left\| \frac{1}{t} \int_0^t \frac{s}{n} \sum_{k=0}^{n-1} \left(I - \frac{s}{n} A \right)^{-k-1} Ax \, ds \right\| \\
&\quad - \left\| \frac{1}{t} \int_0^t \frac{n}{n-1} \frac{d}{ds} \left[\left(I - \frac{s}{n} A \right)^{1-n} y \right] ds \right\| \\
&\geq \|x\| - \frac{1}{t} \int_0^t s \, ds \|Ax\| - \left\| \frac{1}{t} \frac{n}{n-1} \left[\left(I - \frac{t}{n} A \right)^{1-n} y - y \right] \right\| \\
&\geq \|x\| - \left(\frac{t}{2} \|Ax\| + \frac{n}{n-1} \frac{2}{t} \|y\| \right).
\end{aligned}$$

Hence $\|x + Ay\| \geq \|x\| - \left(\frac{t}{2} \|Ax\| + \frac{2}{t} \|y\| \right)$ for all $t > 0$. Minimizing the function $\frac{t}{2} \|Ax\| + \frac{2}{t} \|y\|$, we obtain its minimum $2\sqrt{\|Ax\| \|y\|}$, and hence the estimate (3.4).

Theorem 9 contains as a corollary the following slight improvement of a theorem of Crabb and Sinclair (see [4] or [3, Theorem 20.6]).

Corollary 10. *If an operator $T \in B(X)$ is such that 0 is not an interior point of $\overline{\text{co}} W(T)$, then $\|x + Ty\| \geq \|x\| - 2\sqrt{\|Tx\| \|y\|}$ for all $x, y \in X$.*

Using Corollary 10 one can deduce some consequences. For instance, by modifying the proof of Corollary 20.12 in [3] we obtain the following improvement.

Corollary 11. *Under the hypothesis of Corollary 10, the inequality $\|T\|^2 \leq 4\|T^2\|$ holds.*

ACKNOWLEDGMENT

The authors would like to thank the referee for his valuable suggestions.

REFERENCES

1. J. Anderson, *On normal derivations*, Proc. Amer. Math. Soc. **38** (1973), 135–140. MR **47**:875
2. F. F. Bonsall and J. Duncan, *Numerical Ranges of Operators on Normed Spaces and of Elements of Normed Algebras*, London Math. Soc. Lecture Note Series No. 2, Cambridge Univ. Press, Cambridge, 1971. MR **44**:5779
3. ———, *Numerical Ranges II*, London Math. Soc. Lecture Note Series No. 10, Cambridge Univ. Press, Cambridge, 1973. MR **56**:1063
4. M. J. Crabb and A. M. Sinclair, *On the boundary of the spatial numerical range*, Bull. London Math. Soc. **4** (1972), 17–19. MR **46**:7929
5. M. J. Crabb and P. G. Spain, *Commutators and normal operators*, Glasgow Math. J. **18** (1977), 197–198. MR **56**:1115
6. H. R. Dawson, T. A. Gillespie, and P. G. Spain, *A commutativity theorem for hermitian operators*, Math. Ann. **220** (1976), 215–217. MR **54**:5898
7. N. Dunford, *Some ergodic theorems*, Proc. Royal Soc. Edinburgh **85A** (1980), 111–118. MR **82a**:47006

8. W. Eberlein, *Abstract ergodic theorems and weak almost periodic functions*, Trans. Amer. Math. Soc. **67** (1949), 217–240. MR **12**:112a
9. C. K. Fong, *Normal operators on Banach spaces*, Glasgow Math. J. **20** (1979), 163–168. MR **80e**:47020
10. J. A. Goldstein, *Semigroups of Linear Operators and Applications*, Oxford University Press, Oxford, 1985. MR **87c**:47056
11. E. Hille and R. S. Phillips, *Functional Analysis and Semi-groups*, Amer. Math. Soc. Colloq. Publ., Vol. 31, Amer. Math. Soc., Providence, R.I., 1957. MR **19**:664d
12. U. Krengel, *Ergodic Theorems*, Walter de Gruyter, Berlin, 1985. MR **87i**:28001
13. J. Kyle, *Numerical ranges of derivations*, Proc. Edinburgh Math. Soc. **21** (1978), 33–39. MR **58**:7127
14. K. Mattila, *Normal operators and proper boundary points of the spectra of operators on a Banach space*, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes **19** (1978), 48 pp. MR **80j**:47031
15. ———, *On proper boundary points of the spectrum and complemented eigenspaces*, Math. Scand. **43** (1978), 363–368. MR **80i**:47003
16. ———, *Complex strict and uniform convexity and hyponormal operators*, Math. Proc. Camb. Phil. Soc. **96** (1984), 483–493. MR **86c**:47026
17. S.-Y. Shaw, *On numerical ranges of generalized derivations and related properties*, J. Austral. Math. Soc. (series A) **36** (1984), 134–142. MR **85e**:47001
18. ———, *Mean ergodic theorems and linear functional equations*, J. Funct. Anal. **87** (1989), 428–441. MR **90k**:47018
19. ———, *Uniform convergence of ergodic limits and approximate solutions*, Proc. Amer. Math. Soc. **114** (1992), 405–411. MR **92e**:47012

DEPARTMENT OF MATHEMATICS, NATIONAL CENTRAL UNIVERSITY, CHUNG-LI, TAIWAN 320

Current address, Y.-C. Li: Department of Mathematics, Chung Yuan University, Chung-Li, Taiwan 320

E-mail address: shaw@math.ncu.edu.tw