NONCOMMUTATIVE H^2 SPACES

MICHAEL MARSALLI

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let $\mathcal{M}$ be a von Neumann algebra with a faithful, finite, normal tracial state τ , and let $\mathcal{A}$ be a finite, maximal subdiagonal algebra of $\mathcal{M}$. Let H^2 be the closure of $\mathcal{A}$ in the noncommutative Lebesgue space $L^2(\mathcal{M}, \tau)$. Then H^2 possesses several of the properties of the classical Hardy space on the circle, including a commutant lifting theorem, some results on Toeplitz operators, an H^1 factorization theorem, Nehari's Theorem, and harmonic conjugates which are L^2 bounded.

1. INTRODUCTION

In [1] Arveson introduced the concept of a subdiagonal algebra in order to unify the analysis of several broad classes of nonselfadjoint operator algebras. Arveson drew a close analogy between a subdiagonal algebra and the classical Hardy space H^∞ , the boundary values of the bounded analytic functions on the disk. Roughly, a subdiagonal algebra $\mathcal{A}$ stands in relation to its von Neumann algebra $\mathcal{M}$ as H^∞ stands in relation to the Lebesgue space L^∞ of the unit circle. Subsequently, several authors studied the invariant subspaces of $\mathcal{A}$ acting on the noncommutative Lebesgue space L^p [6], [8], [10]. There has also been considerable investigation of analytic crossed products, which are a type of subdiagonal algebra introduced by McAsey, Muhly and Saito, including their invariant subspace structure [8], [10], maximality among weak* closed subalgebras of $\mathcal{M}$ [8], associated Toeplitz operators [11] and Hankel operators [5]. We shall study the closure of $\mathcal{A}$ in L^2 as an analogue of the classical Hardy space H^2 , and so obtain analogues of several classical results including a commutant lifting theorem, some results on Toeplitz operators, an H^1 factorization theorem, Nehari's Theorem on the norm of a Hankel operator, and the existence and L^2 boundedness of the harmonic conjugate.

Let $\mathcal{M}$ be a von Neumann algebra with a faithful, normal finite tracial state τ . For $1 \leq p < \infty$, let $L^p = L^p(\mathcal{M}, \tau)$ denote the noncommutative Lebesgue space which is associated with $\mathcal{M}$ and τ (cf. [3], [9], [12]). For $t \in \mathcal{M}$, $x \in L^2$, let $L_t(x) = tx$ and $R_t(x) = xt$. Then $\mathcal{L} = \{L_t : t \in \mathcal{M}\}$ and $\mathcal{R} = \{R_t : t \in \mathcal{M}\}$ are von Neumann algebras on L^2 which are each other's commutants. Furthermore, the map $t \rightarrow L_t$ (resp. $t \rightarrow R_t$) is a normal, *-isomorphism (resp. *-anti-isomorphism) of $\mathcal{M}$ onto $\mathcal{L}$ (resp. $\mathcal{R}$), and the identity 1 is a cyclic and separating vector for $\mathcal{L}$ and $\mathcal{R}$. The map $x \rightarrow x^*$ on $\mathcal{M}$ extends to a conjugate linear isometry on L^p . As is customary, we identify $\mathcal{M}$ with L^∞ while the ultraweak topology on $\mathcal{M}$ will

Received by the editors July 10, 1995 and, in revised form, July 27, 1995.

1991 *Mathematics Subject Classification*. Primary 47D15, 46L50.

be identified with the weak* topology on L^∞ regarded as the dual of L^1 . We now introduce $\mathcal{A}$, the noncommutative analogue of H^∞ (cf. [1], [8]).

Definition. Let $\mathcal{A}$ be a w^* -closed unital subalgebra of $\mathcal{M}$, and let Φ be a faithful, normal expectation from $\mathcal{M}$ onto the diagonal $\mathcal{D} = \mathcal{A} \cap \mathcal{A}^*$. Then $\mathcal{A}$ is a finite, maximal subdiagonal subalgebra of $\mathcal{M}$ with respect to Φ if:

- (1) $\mathcal{A} + \mathcal{A}^*$ is w^* -dense in $\mathcal{M}$,
- (2) $\Phi(ab) = \Phi(a)\Phi(b)$ for all $a, b \in \mathcal{A}$,
- (3) $\mathcal{A}$ is maximal among those subalgebras satisfying (1) and (2), and
- (4) $\tau \circ \Phi = \tau$.

For $\mathcal{S} \subset L^p$, $1 \leq p < \infty$, let $[\mathcal{S}]_p$ denote the closure of $\mathcal{S}$ in L^p . Let $H^p = [\mathcal{A}]_p$, $H_0^p = [\{x \in \mathcal{A}: \Phi(x) = 0\}]_p$, and P be the orthogonal projection of L^2 onto H^2 . Then Φ extends to the orthogonal projection of L^2 onto $[\mathcal{D}]_2$ and $L^2 = H^2 \oplus (H_0^2)^* = H_0^2 \oplus [\mathcal{D}]_2 \oplus (H_0^2)^*$ by [8, Proposition 1.1]. For $t \in \mathcal{M}$ we define the (left) Toeplitz operator with symbol t by $T_t = PL_tP$. We define the (left) Hankel operator with symbol t by $H_t = (1 - P)L_tP$.

2. A COMMUTANT LIFTING THEOREM

Let $\mathcal{T}$ denote the algebra $\{L_a: a \in \mathcal{A}\}$ considered as an algebra of operators on H^2 . In this section we identify $\mathcal{T}'$, the operators on H^2 that commute with $\mathcal{T}$. In particular, we show that every element of $\mathcal{T}'$ lifts to an element of $\mathcal{R}$. Our result is analogous to Yoshino's Theorem [13], which describes the commutant of a rationally cyclic subnormal operator. Indeed, if μ is a measure on the complex plane with compact support K , R is the algebra of rational functions with poles off of K , R^2 is the $L^2(\mu)$ closure of R , and S is multiplication by z restricted to R^2 , then the commutant of S is $\{M_\phi: \phi \in R^2 \cap L^\infty(\mu)\}$, where $M_\phi f = \phi f$ is the usual multiplication operator. It follows that if X is an operator on R^2 which commutes with multiplications by elements of R , then $X = M_\phi$ for some $\phi \in L^\infty(\mu)$ with $M_\phi(R^2) \subset R^2$. So the commutant of R acting on its $L^2(\mu)$ closure lifts to the von Neumann algebra $L^\infty(\mu)$ acting on $L^2(\mu)$. For $\mathcal{T}$ acting on H^2 we have the following result.

Theorem 1. *If $X \in \mathcal{T}'$, then there exists $b \in \mathcal{M}$ such that H^2 is invariant for R_b , $X = R_b$ on H^2 , and $\|X\| = \|b\|$.*

Proof. Let $X(1) = h$. First, we will show that $\|rh\|_2 \leq \|X\| \|r\|_2$ for all $r \in \mathcal{M}$. Let $\varepsilon > 0$. Then $r^*r + \varepsilon 1$ is an invertible positive operator in $\mathcal{M}$. So by [1, Corollary 4.2.4] there is an $a \in \mathcal{A}$ such that $r^*r + \varepsilon 1 = a^*a$. Thus

$$\begin{aligned} \langle (r^*r + \varepsilon 1)h, h \rangle &= \langle a^*ah, h \rangle = \|ah\|_2^2 = \|Xa\|_2^2 \\ &\leq \|X\|^2 \|a\|_2^2 = \|X\|^2 \langle a^*a, 1 \rangle = \|X\|^2 \langle r^*r + \varepsilon 1, 1 \rangle. \end{aligned}$$

Letting ε go to 0, we obtain $\|rh\|_2^2 \leq \|X\|^2 \|r\|_2^2$.

Define a map Y via $Y(r) = rh$. Clearly Y extends to a bounded operator on L^2 which commutes with $\mathcal{L}$, and $\|Y\| \leq \|X\|$. Thus there is a $b \in \mathcal{M}$ such that $Y = R_b$. Because $h \in H^2$, Y leaves H^2 invariant. Obviously Y agrees with X on H^2 , and so $\|b\| = \|Y\| = \|X\|$.

3. FACTORIZATIONS, TOEPLITZ AND HANKEL OPERATORS

Our next theorem is a useful factorization result for elements of L^2 , which might be expressed as $L^2 = L^\infty H^2 = H^2 L^\infty$. The proof in the classical case is an

application of the fact that if $\log |f|$ is integrable for an L^2 function f , then $|f| = |h|$ for some H^2 function h . However, this fact is a consequence of Szegő's Theorem, and it is not yet known if Szegő's Theorem extends to the subdiagonal algebra setting. Nevertheless [10, Proposition 1] is a sufficient substitute. As consequences we obtain some facts about Toeplitz operators, the factorization of elements of H^1 as products of H^2 elements (previously obtained by Saito), and Nehari's Theorem.

Theorem 2. *For every $\varepsilon > 0$ and $z \in L^2$, there exist $h_1, h_2 \in H^2$ and $v_1, v_2 \in \mathcal{M}$, such that $z = v_1 h_1 = h_2 v_2$, $\|v_i\| \leq 1$, $\|h_i\|_2 < (1 + \varepsilon)\|z\|_2$, and $h_i^{-1} \in \mathcal{A}$ for $i = 1, 2$.*

Proof. We prove the existence of v_1 and h_1 . The proof for v_2 and h_2 is similar. Choose $\delta < \sqrt{2\varepsilon + \varepsilon^2}\|z\|_2$. Consider the positive weak* continuous linear functional $\omega_{\delta 1 + \omega_z}$ on $\mathcal{R}$. By [7, Theorem 7.2.3] there is a vector $y \in L^2$ such that $\omega_y = \omega_{\delta 1 + \omega_z}$ on $\mathcal{R}$. Clearly, $\omega_{\delta 1} \leq \omega_y$ and $\omega_z \leq \omega_y$ on $\mathcal{R}$. It follows that there exist $r, s \in \mathcal{M}$ such that $\delta 1 = ry$ and $z = sy$. Because $\mathcal{M}$ is finite, $\delta^{-1}r$ has an inverse in L^2 .

By [10, Proposition 1] there exist a unitary $u \in \mathcal{M}$ and $g \in \mathcal{A}$ such that $\delta^{-1}r = gu$, and $h = g^{-1} \in H^2$. Thus $y = u^*h$, $z = su^*h$, and $\|h\|_2 = \|y\|_2 = \sqrt{\delta^2 + \|z\|_2^2} < (1 + \varepsilon)\|z\|_2$. Now let $v_1 = su^*$ and $h_1 = h$.

In the classical setting, one has that the map $t \rightarrow T_t$ is an isometry on L^∞ , but the proof relies heavily on the fact that L^∞ is commutative. We do not know if this map is isometric in our noncommutative setting. We do have the following partial results, including the fact that the map is very well behaved on $\mathcal{A}$.

Corollary 3. *For every $t \in \mathcal{M}$, $\|L_t P\| = \|t\|$.*

Proof. We have $\|L_t\| = \|t\|$. So given $\varepsilon > 0$, choose z such that $\|z\|_2 < 1$ and $\|tz\|_2 \geq \|t\| - \varepsilon$. By Theorem 2 we can write $z = hv$ where $v \in \mathcal{M}$, $\|v\| \leq 1$, $h \in H^2$, and $\|h\|_2 < 1$. Thus $\|tz\|_2 = \|thv\|_2 \leq \|th\|_2$, so $\|th\|_2 \geq \|t\| - \varepsilon$.

Corollary 4. *For every $t \in \mathcal{M}$, $\|T_{t^*t}\| = \|t^*t\|$.*

Proof. We have $\|T_{t^*t}\| = \|PL_{t^*t}P\| = \|PL_t^*L_tP\| = \|L_tP\|^2 = \|t\|^2 = \|t^*t\|$.

Corollary 5. *The map $a \rightarrow T_a$ is an isometric algebra isomorphism and a weak* homeomorphism of $\mathcal{A}$ onto $\{T_a : a \in \mathcal{A}\}$.*

Proof. For $a \in \mathcal{A}$, $T_a = L_a P$, so by Corollary 3 the map is an isometry. Because the map $a \rightarrow L_a$ is a weak* continuous isomorphism and H^2 is invariant for $\{L_a : a \in \mathcal{A}\}$, the map $a \rightarrow T_a$ is a weak* continuous isomorphism. It follows from [2, Theorem 1.20] that $\{T_a : a \in \mathcal{A}\}$ is weak* closed and the map is a weak* homeomorphism.

Let $\mathcal{A}_*$ denote the space of weak* continuous linear functionals on $\mathcal{A}$. The next result shows that the weak* continuous functionals on the algebra $\{T_a : a \in \mathcal{A}\}$ have a rank one structure. In the language of the theory of dual algebras [2] the algebra has property $\mathbf{A}_1(1)$.

Corollary 6. *For every $\varepsilon > 0$ and for every $\phi \in \mathcal{A}_*$, there exist $g, h \in H^2$ such that $\phi(a) = \langle ag, h \rangle$ for $a \in \mathcal{A}$, and $\|g\|_2\|h\|_2 < (1 + \varepsilon)\|\phi\|$.*

Proof. For $\varepsilon > 0$, choose $\eta > 0$ such that $(1 + \eta)^2 < 1 + \varepsilon$. There is a functional $\hat{\phi}$ in $\mathcal{M}_*$ such that $\hat{\phi}$ extends ϕ and $\|\hat{\phi}\| < (1 + \eta)\|\phi\|$ by [4, Lemma 2.4]. Because of the duality between L^1 and $\mathcal{M}$, there is an $f \in L^1$ such that $\hat{\phi}(t) = \tau(tf)$ for $t \in \mathcal{M}$ and $\|\hat{\phi}\| = \|f\|_1$. So there exist vectors $x, y \in L^2$ such that $f = xy$ and

$\|x\|_2\|y\|_2 = \|f\|_1 = \|\hat{\phi}\| < (1 + \eta)\|\phi\|$. By Theorem 2 $x = gv$ with $g \in H^2$, $v \in \mathcal{M}$, $\|v\| \leq 1$, and $\|g\|_2 < (1 + \eta)\|x\|_2$. Thus, for all $a \in \mathcal{A}$,

$$\phi(a) = \langle L_agv, y \rangle = \langle L_ag, yv^* \rangle = \langle L_ag, P(yv^*) \rangle.$$

Let $h = P(yv^*)$, and the result follows.

We now proceed to establish an analogue of the factorization theorem for H^1 . Recall that any function f in the classical H^1 space can be written $f = gh$ where g, h are in H^2 and $\|f\|_1 = \|g\|_2\|h\|_2$. The following result was previously obtained by Saito [11, Lemma 5.5], but we include it for completeness.

Corollary 7. *For every $\varepsilon > 0$ and for every $f \in H^1$, there exist $g, h \in H^2$ such that $f = gh$ and $\|g\|_2\|h\|_2 < (1 + \varepsilon)\|f\|_1$. If $f \in H_0^1$, then h is in H_0^2 .*

Proof. Because $f \in L^1$, there exist vectors $x, y \in L^2$ such that $f = xy$ and $\|x\|_2\|y\|_2 = \|f\|_1$. By Theorem 2 we can find $g \in H^2$ and $v \in \mathcal{M}$ such that $x = gv$, $\|v\| \leq 1$, $\|g\|_2 < (1 + \varepsilon)\|x\|_2$ and $g^{-1} \in \mathcal{A}$. Now by [10, Lemma 3] $H^1 = \{x \in L^1 : \tau(xy) = 0 \text{ for all } y \in H_0^\infty\}$. Thus, for $a \in \mathcal{A}_0$,

$$\tau(af) = \tau(axy) = \tau(agvy) = \langle ag, (vy)^* \rangle.$$

Let $h = vy$. Then $h^* \in [\mathcal{A}_0g]_2^\perp$. But $[\mathcal{A}_0g]_2 = [\mathcal{A}_0]_2$, because $g^{-1} \in \mathcal{A}$. Thus $h^* \in [\mathcal{A}_0]_2^\perp = (H^2)^*$, so $h \in H^2$ and $f = xy = gvy = gh$. Finally, $\|g\|_2\|h\|_2 < (1 + \varepsilon)\|x\|_2\|y\|_2 = (1 + \varepsilon)\|f\|_1$.

Note that if $f \in H_0^1$, then $\tau(af) = 0$ for all $a \in \mathcal{A}$. So $h^* \in (H^2)^\perp = (H_0^2)^*$, and thus $h \in H_0^2$.

As a consequence of the previous factorization theorem, we can obtain an exact analogue of Nehari's Theorem on the norm of a Hankel operator. In the classical setting, the norm of a Hankel operator H_f is given by the distance in L^∞ from the symbol f to H^∞ . This remains true in the noncommutative setting.

Corollary 8. *For every $t \in \mathcal{M}$, $\|H_t\| = \text{dist}(t, \mathcal{A})$.*

Proof. By [10, Lemma 4] $H_0^1 = \{f \in L^1 : \tau(fy) = 0 \text{ for all } y \in \mathcal{A}\}$. By Corollary 7 $\{f \in H_0^1 : \|f\|_1 < 1\} = \{gh : g \in H^2, h \in H_0^2, \|g\|_2\|h\|_2 < 1\}$. Because $\mathcal{A}$ is w^* -closed,

$$\begin{aligned} \text{dist}(t, \mathcal{A}) &= \sup\{|\tau(tf)| : f \in L^1, \tau(yf) = 0 \text{ for all } y \in \mathcal{A}, \|f\|_1 < 1\} \\ &= \sup\{|\tau(tf)| : f \in H_0^1, \|f\|_1 < 1\} \\ &= \sup\{|\tau(tgh)| : g \in H^2, h \in H_0^2, \|g\|_2\|h\|_2 < 1\} \\ &= \sup\{|\langle L_tg, h^* \rangle| : g \in H^2, h \in H_0^2, \|g\|_2\|h\|_2 < 1\} \\ &= \|H_t\|. \end{aligned}$$

4. HARMONIC CONJUGATES

Every harmonic function u on the unit disk has an associated harmonic function $\tilde{u}$, the harmonic conjugate of u , such that $u + i\tilde{u}$ is analytic and $\tilde{u}(0) = 0$. The map $u \rightarrow \tilde{u}$ is a real linear transformation which is L^2 bounded. We will establish an analogue of this result by thinking of $\text{Re } \mathcal{A}$, the real parts of the operators in $\mathcal{A}$, as a noncommutative version of the space of bounded harmonic functions. We first construct a real linear map from $\text{Re } \mathcal{A}$ to $\mathcal{A}$ which is the analogue of the Herglotz transform, i.e. integration against the kernel $(e^{i\theta} + z)(e^{i\theta} - z)^{-1}$.

Theorem 9. *There is a real linear transformation $T: \operatorname{Re} \mathcal{A} \rightarrow \mathcal{A}$ such that $u = \operatorname{Re}(Tu)$, $\Phi(\operatorname{Im} Tu) = 0$, and $\|Tu\|_2 \leq \sqrt{2}\|u\|_2$ for all $u \in \operatorname{Re} \mathcal{A}$. Thus T extends to a bounded operator from $[\operatorname{Re} \mathcal{A}]_2$ to H^2 .*

Proof. Let $u \in \operatorname{Re} \mathcal{A}$. So $u = \operatorname{Re} g$ for some $g \in \mathcal{A}$. Let $a = g - \frac{1}{2}\Phi(g - g^*)$. Then $a \in \mathcal{A}$, $u = \operatorname{Re} a$, and $\Phi(\operatorname{Im} a) = 0$. Thus there exists an $a \in \mathcal{A}$ with the desired properties. We now show that such an element of $\mathcal{A}$ is unique. Suppose that $u = \operatorname{Re} a = \operatorname{Re} h$ for $a, h \in \mathcal{A}$, and $\Phi(\operatorname{Im} a) = \Phi(\operatorname{Im} h) = 0$. Then $a + a^* = h + h^*$ implies that $a - h = h^* - a^*$, so $(a - h)^* = h - a$. Furthermore, $\Phi(\operatorname{Im} a) = \Phi(\operatorname{Im} h) = 0$ implies that $\Phi(a) = \Phi(a^*)$ and $\Phi(h) = \Phi(h^*)$. Consequently, $\Phi(a) = \Phi(h)$, because $\operatorname{Re} a = \operatorname{Re} h$. Thus

$$\Phi((a - h)(a - h)^*) = \Phi((a - h)(h - a)) = \Phi(a - h)\Phi(h - a) = 0.$$

So $a = h$, because Φ is faithful. Thus we can define $Tu = a$, where a is the unique element of $\mathcal{A}$ with $u = \operatorname{Re} a$ and $\Phi(\operatorname{Im} a) = 0$. It is easy to see that T is real linear.

We now proceed to establish the inequality $\|Tu\|_2 \leq \sqrt{2}\|u\|_2$. Let $Tu = d + b$, where $d \in \mathcal{D}$, $b \in \mathcal{A}_0$. Now $\Phi(\operatorname{Im} Tu) = 0$ and $\Phi(b) = \Phi(b^*) = 0$, so we have $d = \Phi(d) = \Phi(d^*) = d^*$. Thus

$$\|Tu\|_2^2 = \tau((d + b)(d + b^*)) = \|d\|_2^2 + \tau(db^*) + \tau(db) + \|b\|_2^2 = \|d\|_2^2 + \|b\|_2^2.$$

By a simple computation we have

$$0 = \tau(b^2) = \tau((\operatorname{Re} b)^2) - \tau((\operatorname{Im} b)^2) + 2i\tau((\operatorname{Re} b)(\operatorname{Im} b)).$$

Note that

$$\overline{\tau((\operatorname{Re} b)(\operatorname{Im} b))} = \tau((\operatorname{Im} b)(\operatorname{Re} b)) = \tau((\operatorname{Re} b)(\operatorname{Im} b)).$$

By equating real and imaginary parts we obtain

$$\|(\operatorname{Re} b)\|_2^2 = \|(\operatorname{Im} b)\|_2^2 \quad \text{and} \quad \tau((\operatorname{Re} b)(\operatorname{Im} b)) = 0.$$

So

$$\|b\|_2^2 = \|(\operatorname{Re} b)\|_2^2 + \|(\operatorname{Im} b)\|_2^2 = 2\|(\operatorname{Re} b)\|_2^2.$$

Thus

$$\|Tu\|_2^2 = \|\operatorname{Re} d\|_2^2 + 2\|\operatorname{Re} b\|_2^2.$$

Now because b is orthogonal to $\mathcal{D}$, we have $\tau((\operatorname{Re} d)(\operatorname{Re} b)) = 0$. Consequently,

$$\|Tu\|_2^2 \leq 2(\|\operatorname{Re} d\|_2^2 + \|\operatorname{Re} b\|_2^2) = 2\|\operatorname{Re}(d + b)\|_2^2 = 2\|u\|_2^2.$$

Corollary 10. *For each $u \in [\operatorname{Re} \mathcal{A}]_2$ there is a unique $\tilde{u}$ in $[\operatorname{Re} \mathcal{A}]_2$ such that $u + i\tilde{u} \in H^2$ and $\Phi(\tilde{u}) = 0$. Furthermore, $\|\tilde{u}\|_2 \leq \sqrt{2}\|u\|_2$.*

Proof. Let $\tilde{u} = \operatorname{Im} Tu$.

REFERENCES

1. W. Arveson, *Analyticity in operator algebras*, Amer. J. Math. **89** (1967), 578–642. MR **36**:6946
2. H. Bercovici, C. Foiaş, and C. Pearcy, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS Regional Conf. Ser. in Math., no. 56, Amer. Math. Soc., Providence, RI, 1985. MR **87g**:47091
3. J. Dixmier, *Formes linéaires sur un anneau d'opérateurs*, Bull. Soc. Math. France **81** (1953), 9–39. MR **15**:539a
4. D. Hadwin and E. Nordgren, *Subalgebras of reflexive algebras*, J. Operator Theory **7** (1983), 3–23. MR **83f**:47033

5. Y. Imin and K.-S. Saito, *Hankel operators associated with analytic crossed products*, Can. Math. Bull. **37** (1994), 75–81. MR **94k**:47024
6. N. Kamei, *Simply invariant subspace theorems for antisymmetric finite subdiagonal algebras*, Tohoku Math. J. **21** (1969), 467–473. MR **41**:839
7. R. Kadison and J. Ringrose, *Fundamentals of the Theory of Operator Algebras*, Academic Press, New York, 1983. MR **85j**:46099
8. M. McAsey, P. Muhly and K.-S. Saito, *Nonselfadjoint crossed products (invariant subspaces and maximality)*, Trans. Amer. Math. Soc. **248** (1979), 381–410. MR **80j**:46101b
9. E. Nelson, *Notes on non-commutative integration*, J. Functional Analysis **15** (1974), 103–116. MR **50**:8102
10. K.-S. Saito, *A note on invariant subspaces for finite maximal subdiagonal algebras*, Proc. Amer. Math. Soc. **77** (1979), 348–352. MR **81b**:46078
11. ———, *Toeplitz operators associated with analytic crossed products*, Math. Proc. Cambridge Phil. Soc. **108** (1990), 539–549. MR **91m**:46109
12. I. Segal, *A noncommutative extension of abstract integration*, Ann. of Math. **57** (1953), 401–457. MR **14**:991f
13. T. Yoshino, *Subnormal operators with a cyclic vector*, Tôhoku Math. J. **21** (1973), 47–55. MR **39**:7450

DEPARTMENT OF MATHEMATICS, ILLINOIS STATE UNIVERSITY, NORMAL, ILLINOIS 61790-4520
E-mail address: marsalli@math.ilstu.edu