A NOTE ON THE RELATIVE CLASS NUMBER IN FUNCTION FIELDS

MICHAEL ROSEN

(Communicated by William W. Adams)

ABSTRACT. Let F be a finite field, $A = F[T]$, and $k = F(T)$. Let $K_m = k(\Lambda_m)$ be the field extension of k obtained by adjoining the m -torsion on the Carlitz module. The class number h_m of K_m can be written as a product $h_m = h_m^+ h_m^-$. The number h_m^- is called the relative class number. In this paper a formula for h_m^- is derived which is the analogue of the Maillet determinant formula for the relative class number of the cyclotomic field of p -th roots of unity. Some consequences of this formula are also derived.

Let $\mathbf{Q}$ denote the rational numbers and consider the cyclotomic field $K_p = \mathbf{Q}(\zeta_p)$ with class number h_p . It is well known that this class number factors as a product $h_p^+ h_p^-$ of two integers. The number h_p^+ is the class number of the maximal real subfield of K_p . The integer h_p^- is called the relative class number. In [Ca] and [Ca-O] it is shown how the relative class number can be computed in terms of a certain classical determinant known as the Maillet determinant. A nice exposition of this is given in Chapter 3 of [L] (see Theorem 7.1).

In this note we will give an analogue of this material in the context of cyclotomic function fields. Let F be a finite field with q elements and $A = F[T]$ the polynomial ring over F . Let $k = F(T)$ be the quotient field of A . For an irreducible polynomial m of degree d we will denote by Λ_m the m -torsion on the Carlitz module and let $K_m = k(\Lambda_m)$ be the “cyclotomic” function field obtained by adjoining the elements of Λ_m to k . For the definition of the Carlitz module and its properties see [H] and [G-R]. The class number of K_m , h_m , factors as a product of two integers $h_m = h_m^+ h_m^-$, where h_m^+ is the class number of the “maximal real” subfield of K_m , i.e. the decomposition field of the prime at infinity of k in K_m , and h_m^- is called the relative class number. Our aim is to give an expression for h_m^- as a product of certain easily computed determinants related to the classical Maillet determinant.

We begin by recalling the analytic class number formula for h_m^- which follows immediately from Theorem 2 of [G-R]. A character χ of $(A/mA)^*$ is said to be *real* if its restriction to F^* is the trivial character. Otherwise it is said to be *non-real* or *imaginary*. If χ is imaginary, define $S(\chi) = \sum_a \chi(a)$, where the sum is over all monic polynomials of degree less than d . Then,

$$(1) \quad h_m^- = \prod_{\chi \text{ imaginary}} S(\chi).$$

Received by the editors July 2, 1995 and, in revised form, November 15, 1995.

1991 *Mathematics Subject Classification*. Primary 11R29; Secondary 11R58, 14H05.

This work was partially supported with a grant from the National Science Foundation.

Let $t = (q^d - 1)/(q - 1)$. Then t is the size of the set $\mathcal{M}$ of monic polynomials of degree less than d . We construct a $t \times t$ matrix $[c(a, b)]$ where $a, b \in \mathcal{M}$. Namely, for $a, b \in \mathcal{M}$ write $ab = sm + r$, where $s, r \in A$ and either $r = 0$ or $\deg(r) < d$. In fact, r cannot be zero since we have assumed that m is irreducible. Define $c(a, b)$ to be the leading coefficient of r . By construction, the matrix $[c(a, b)]$ has all its coefficients in F^* . Let λ be a character of F^* and define

$$d(\lambda) = \det[\lambda(c(a, b))].$$

Theorem 1.

$$h_m^- = \pm \prod_{\lambda \neq \lambda_o} d(\lambda),$$

where the product is over all the non-trivial characters λ of F^* .

Proof. The proof proceeds by a step by step rewriting of equation (1). Let λ be any non-trivial character of F^* , and set

$$h_\lambda = \prod_{\chi|_{F^*}=\lambda} S(\chi).$$

From equation (1) we see that

$$(2) \quad h_m^- = \prod_{\lambda \neq \lambda_o} h_\lambda.$$

Let ϕ be a fixed character whose restriction to F^* is λ . Then all the characters whose restriction to F^* is λ can be written $\phi\chi'$, where χ' is a real character. Note that a real character can also be thought of as a character of the group $G = (A/mA)^*/F^*$. We'll come back to this in a moment.

Starting with a non-trivial character λ on F^* we construct a function $\tilde{\lambda}$ on $(A/mA)^*$ as follows. Let a be a polynomial prime to m . Write $a = sm + r$ with $s, r \in A$ and $\deg(r) < d$. Define $\tilde{\lambda}(a)$ to be λ evaluated on the inverse of the leading coefficient of r . Then $\tilde{\lambda}$ is a well defined function on $(A/mA)^*$.

We rewrite h_λ using the quantities just defined:

$$\begin{aligned} h_\lambda &= \prod_{\chi|_{F^*}=\lambda} \sum_{a \text{ monic}, \deg(a) < d} \chi(a) \\ &= \prod_{\chi'} \sum_{a \text{ monic}, \deg(a) < d} \chi'(a) \phi(a) \tilde{\lambda}(a). \end{aligned}$$

Observe that the function $\phi(a)\tilde{\lambda}(a)$ is unchanged if we replace a by αa where α is a constant. It is also unchanged if we replace a by anything in the congruence class of a modulo m . Thus the summation in the above sum can be rewritten as a summation over all the elements of the group $G = (A/mA)^*/F^*$. We obtain

$$h_\lambda = \prod_{\chi'} \sum_{a \in G} \chi'(a) [\phi(a)\tilde{\lambda}(a)].$$

We apply the Dedekind determinant formula (see [L], Theorem 6.1) to derive

$$h_\lambda = \det [\phi(b^{-1}a)\tilde{\lambda}(b^{-1}a)].$$

Here, a and b vary over the elements of the group G . Replacing b^{-1} by b merely permutes the rows of the matrix, and so,

$$(3) \quad h_\lambda = \pm \det \left[\phi(b)\phi(a)\tilde{\lambda}(ba) \right].$$

We have used the fact that ϕ is a character. Each element of the b' th row of the determinant in equation (3) contains $\phi(b)$ as a factor. Similarly, each element of the a' th column contains $\phi(a)$ as a factor. By elementary properties of determinants, it follows that

$$h_\lambda = \pm \phi \left(\prod_{a \in G} a \right)^2 \det \left[\tilde{\lambda}(ab) \right].$$

The product of all the elements in an abelian group is equal to the product of all the elements of order two. Thus,

$$\phi \left(\prod_{a \in G} a \right)^2 = (\pm 1)^2 = 1,$$

and so,

$$(4) \quad h_\lambda = \pm \det \left[\tilde{\lambda}(ab) \right] = \pm d(\lambda^{-1}).$$

The last equality is obtained by letting a and b once again run through all the monic polynomials of degree less than d and simply using the definition of $\tilde{\lambda}$. From equations (2) and (4) the proof of the theorem is immediate. $\square$

I would like to thank David Goss for suggesting that the following result should hold.

Theorem 2. *Let $[c(a, b)]$ be the matrix introduced in the remarks preceding Theorem 1. Then, the relative class number h_m^- is divisible by p , the characteristic of F , if and only if there is an integer i , $1 \leq i \leq q - 2$, such that $\det[c(a, b)^i] = 0$.*

Proof. We have to reinterpret equation (2) first p -adically and then, after reduction mod p , as an equality in the ring A/mA .

Let ζ be a complex primitive $(q^d - 1)$ st root of unity and set $E = \mathbf{Q}(\zeta)$. Let $\mathcal{O}$ be the ring of integers in E . E is unramified at all primes above p . Let $\mathcal{P}$ be a fixed such prime and let $\hat{\mathcal{O}}$ be the completion of $\mathcal{O}$ at $\mathcal{P}$. Using the natural imbedding of $\mathcal{O}$ in $\hat{\mathcal{O}}$ we can (and do) interpret equation (2) as holding in $\hat{\mathcal{O}}$. Now, the residue class field, κ , of $\hat{\mathcal{O}}$ is a finite field with q^d elements and so is isomorphic to A/mA . Let $\rho: \kappa \rightarrow A/mA$ be such an isomorphism. Finally, let r denote reduction modulo $\hat{\mathcal{P}}$ on $\hat{\mathcal{O}}$. Applying the composed map $\rho \circ r$ to both sides of equation (2), we see that p divides h_m^- if and only if $\rho \circ r(h_\lambda) = 0$ in A/mA for some complex character λ of F^* .

Let λ be any complex character of F^* , and define $\tilde{\lambda} = \rho \circ r \circ \lambda$. Then $\tilde{\lambda}$ is a character on F^* with values in the $(q - 1)$ st roots of unity in $(A/mA)^*$, i.e. in F^* . It is easy to see that the multiplicative maps from F^* to itself consist of precisely the powers of the identity map under pointwise product. Now, apply the homomorphism $\rho \circ r$ to both sides of equation (4). We find

$$\rho \circ r(h_\lambda) = \pm d(\rho \circ r \circ \lambda^{-1}) = \pm \det(c(a, b)^i),$$

where i is some index between 1 and $q - 2$. This proves the theorem. $\square$

Remark 1. It is undoubtedly true that Theorem 2 can be refined so that the vanishing of $\det[c(a, b)^i]$ can be related to the p -divisibility of the order of the “ i th piece” of the class group. See [G-S] for an explanation of how to decompose the class group into pieces corresponding to powers of the Teichmüller character. We shall not enter into this here.

As in [Ca], [Ca-O], and [L], it is possible to use Theorem 1 to give an upper bound for h_m^- .

Theorem 3. *As above, let $m \in A$ be an irreducible polynomial of degree d and define $t = q^d - 1/(q - 1)$. We have*

$$(5) \quad h_m^- \leq \sqrt{t}^{t(q-2)} < (2q)^{\frac{d-1}{2}q^d}.$$

Proof. We apply the Hadamard determinant inequality which states that the absolute value of the determinant of a complex square matrix is less than or equal to the product of the lengths of the row vectors which compose the matrix. Since all the entries of $[\lambda(c(a, b))]$ have absolute value 1, we find that $|d(\lambda)| \leq \sqrt{t}^t$. The first inequality follows from this and equation (2). As for the second inequality, note that the exponent $t(q-2) < q^d - 1 < q^d$. Also, note that $t = q^{d-1} + q^{d-2} + \dots + 1 < 2q^{d-1}$. These two observations yield the second inequality. $\square$

It is interesting to compare the inequalities of Theorem 3 with those obtained from the congruence Riemann hypothesis. Recall that $h_m^- = h_m/h_m^+$. It follows from the theory of algebraic curves over finite fields and the congruence Riemann hypothesis that there are complex numbers π_i of absolute value $\sqrt{q}$ such that

$$(6) \quad h_m^- = \prod_{i=1}^{2g-2g^+} (1 - \pi_i).$$

Here, g , resp. g^+ , is the genus of the field K_m , resp. K_m^+ . The only primes of k which ramify in K_m are (m) and ∞ . The prime (m) is totally ramified of degree d , whereas ∞ splits into t primes each with ramification index $q - 1$ and degree 1. By Riemann-Hurwitz one deduces that

$$2g - 2 = (d - 1)q^d + 1 - 2d - t.$$

See [H] for more details and the case of general m . In K_m^+ only (m) is ramified, and it is totally ramified. It follows that

$$2g^+ - 2 = (d - 2)t - d.$$

Subtracting these equations, we find that

$$(7) \quad 2g - 2g^+ = (d - 1)q^d + O(q^{d-1}).$$

The error term is an explicit polynomial in q whose leading term is $(1 - d)q^{d-1}$, so for large d the error term makes a negative contribution.

Combining equations (6) and (7) and using the very coarse inequality $\sqrt{q} + 1 < 2\sqrt{q}$, we deduce that

$$(8) \quad h_m^- < (\sqrt{q} + 1)^{2g-2g^+} < (4q)^{g-g^+}.$$

Equations (7) and (8) yield

$$\log_q(h_m^-) < \left(1 + \frac{\log(4)}{\log q}\right) \frac{d-1}{2} q^d + O(q^{d-1}),$$

whereas the elementary methods used in proving Theorem 3 (see equation (5)) give the result

$$\log_q(h_m^-) < \left(1 + \frac{\log(2)}{\log(q)}\right) \frac{d-1}{2} q^d,$$

which is surprisingly good.

We conclude with two more remarks.

Remark 2. Throughout this paper we have been concerned with the class numbers associated to the fields K_m and K_m^+ . It is also of interest to consider the class numbers of the rings $\mathcal{O}_m$ and $\mathcal{O}_m^+$ which are the integral closures of A in K_m and K_m^+ respectively. Call these class numbers $h(\mathcal{O}_m)$ and $h(\mathcal{O}_m^+)$. It can be shown that the second of these numbers divides the first. Call the ratio $h(\mathcal{O}_m)^-$. This is the relative class number on the level of rings. The following relationship holds:

$$h(\mathcal{O}_m)^- = (q-1)^{1-t} h_m^-.$$

For a proof of this, in a more general setting, see [R].

The upshot is that Theorem 1 provides a formula for $h(\mathcal{O}_m)^-$ as well as h_m^- . Also, Theorem 2 gives a criterion for p -divisibility of both of these numbers.

Remark 3. The “cyclotomic” construction of ray class fields of $k = F(T)$ due to Carlitz has been extended to arbitrary global function fields by V.G. Drinfeld and D.R. Hayes. Using Hayes’ normalized rank one Drinfeld modules, L. Shu [S] has developed analytic class number formulas for the generalization of the relative class numbers h_m^- . Using these, she has been able to extend Theorems 1 and 2 to this setting. Her paper is in preparation.

REFERENCES

- [Ca] L. Carlitz, *A generalization of Maillet’s determinant and a bound for the first factor of the class number*, Proc. AMS **12** (1961), 256–261. MR **22**:12093
- [Ca-O] L. Carlitz and F.R. Olson, *Maillet’s Determinant*, Proc. AMS **6** (1955), 265–269. MR **16**:999d
- [G-R] S. Galovich and M. Rosen, *The Class Number of Cyclotomic Function Fields*, J. of Number Theory **13**, No. 3 (1981), 363–375. MR **83m**:12022
- [G-S] D. Goss and W. Sinnott, *Class-groups of Function Fields*, Duke Math. J. **52**, No. 2 (1985), 507–516. MR **87b**:11118
- [H] D. Hayes, *Explicit class field theory for rational function fields*, Trans. Amer. Math. Soc. **189** (1974), 77–91. MR **48**:8444
- [L] S. Lang, *Cyclotomic Fields*, Springer Verlag, New York-Heidelberg-Berlin, GTM 59, 1978. MR **58**:5578
- [R] M. Rosen, *The Hilbert class field in function fields*, Exposition. Math. **5** (1987), 365–378. MR **89b**:10094
- [S] L. Shu, *Narrow ray class fields and partial zeta functions*, pre-print, 1995.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RHODE ISLAND 02912-0001
E-mail address: `ma408000@brownvm.brown.edu`