A NEW PROOF OF THE TWO WEIGHT NORM INEQUALITY FOR THE ONE-SIDED FRACTIONAL MAXIMAL OPERATOR

DAVID CRUZ-URIBE, SFO

(Communicated by J. Marshall Ash)

ABSTRACT. We give a new proof of the two weight norm inequality for the one-sided, fractional maximal operator, M_α^+ , simplifying the original proof of Martín-Reyes and de la Torre.

1. INTRODUCTION

In [1], Andersen and Sawyer introduced the one-sided fractional maximal operators

$$M_\alpha^+ f(x) = \sup_{t>0} \frac{1}{t^{1-\alpha}} \int_x^{x+t} |f| dy \quad \text{and} \quad M_\alpha^- f(x) = \sup_{t>0} \frac{1}{t^{1-\alpha}} \int_{x-t}^x |f| dy,$$

$0 < \alpha < 1$, in order to study the weighted norm inequalities for the Riemann-Liouville and Weyl fractional integral operators. Using complex interpolation, they proved one-weight norm inequalities for M_α^+ and M_α^- . In [3], Martín-Reyes and de la Torre answered a question of Andersen and Sawyer by giving a geometric proof of a two-weight norm inequality for M_α^+ . (The same result holds, *mutatis mutandis*, for M_α^- .)

Theorem 1.1. *For non-negative weights u and v , and for $1 < p \leq q$, the following are equivalent:*

1. *There exists a constant C such that, for every function f in $L^p(v)$,*

$$\left(\int_{\mathbb{R}} (M_\alpha^+ f)^q u dx \right)^{1/q} \leq C \left(\int_{\mathbb{R}} |f|^p v dx \right)^{1/p};$$

2. *The pair (u, v) satisfy the $(S_{p,q,\alpha}^+)$ condition: there exists a constant C such that for every interval $I = [a, b]$ for which $u((-\infty, a)) > 0$,*

$$\left(\int_I M_\alpha^+ (\sigma \chi_I)^q u dx \right)^{1/q} \leq C \left(\int_I \sigma dx \right)^{1/p} < \infty,$$

where $\sigma = v^{1-p'}$.

Received by the editors August 29, 1995 and, in revised form, October 25, 1995 and November 16, 1995.

1991 *Mathematics Subject Classification.* Primary 42B25.

Key words and phrases. One-sided fractional maximal operator, weighted norm inequalities.

Their proof involved proving a weighted norm inequality for a dyadic variant of the fractional maximal operator, $M_{\alpha D}^+$, using a dyadic version of the Sawyer condition, $(S_{pq\alpha D}^+)$, and then showing that these were equivalent to M_{α}^+ and $(S_{pq\alpha}^+)$.

The purpose of this paper is to give a new proof of Theorem 1.1, one which eliminates the dyadic maximal operator. We do this by adapting the proof in the dyadic case using special covering properties of $\mathbb{R}$ and the continuity properties of the maximal operator. We believe that these techniques will be useful in proving other norm inequalities for maximal operators on $\mathbb{R}$.

The paper is organized as follows: Section 2 contains three lemmas and some remarks on their applicability, and Section 3 contains the actual proof. Throughout, all functions are assumed to be measurable, C denotes a positive constant whose value may be different at each appearance, $p' = p/(p-1)$ is the conjugate exponent of p , and $0 < \alpha < 1$. Given a Borel set E and a function w , let $|E|$ denote the Lebesgue measure of E and $w(E) = \int_E w \, dx$.

2. PRELIMINARY RESULTS

Lemma 2.1. *Let f be a non-negative, bounded, upper semicontinuous function of compact support. Then $M_{\alpha}^+ f$ is continuous.*

Proof. Since $M_{\alpha}^+ f$ is always lower semicontinuous, it will suffice to show that it is upper semicontinuous at each point $x \in \mathbb{R}$. We will show this by contradiction: suppose that there exist an $\epsilon > 0$, a point x_0 and a sequence of points $\{x_n\}$ converging to x_0 such that $M_{\alpha}^+ f(x_n) > M_{\alpha}^+ f(x_0) + \epsilon$. For each n there exists an interval J_n whose left endpoint is x_n such that

$$(1) \quad \frac{1}{|J_n|^{1-\alpha}} \int_{J_n} |f| \, dx > M_{\alpha}^+ f(x_0) + \epsilon.$$

Let the support of f be contained in the (finite) interval I , and let K be a finite, open interval containing I and the x_n 's, $n \geq 0$. Then we may assume that each J_n is contained in K . Therefore, after passing to a subsequence we may assume that the J_n 's converge to a possibly degenerate interval J whose left endpoint is x_0 . If $|J| > 0$ then inequality (1) implies that

$$M_{\alpha}^+ f(x_0) \geq \frac{1}{|J|^{1-\alpha}} \int_J |f| \, dx \geq M_{\alpha}^+ f(x_0) + \epsilon,$$

a contradiction. If $J = \{x_0\}$ then for n sufficiently large, by the upper semicontinuity of f , $f(x) - f(x_0) < \epsilon$ for all $x \in J_n$. Therefore,

$$\frac{1}{|J_n|^{1-\alpha}} \int_{J_n} f(x) \, dx < |J_n|^{\alpha} (f(x_0) + \epsilon).$$

If we combine this with inequality (1) and take the limit as n tends to infinity, we get (since f is bounded) that $0 \geq M_{\alpha}^+ f(x_0) + \epsilon$, which is again a contradiction. Hence $M_{\alpha}^+ f$ is upper semicontinuous at each point, and we are done. $\square$

It is worth noting that this lemma is not true if $\alpha = 0$. A simple counter-example is given by the characteristic function of $[0, 1]$. (I am grateful to Juha Kinnunen for pointing this out to me.)

The next lemma is due to Jesus Aldaz; the proof is in Bliedtner and Loeb [2].

Lemma 2.2. *If μ is a finite Borel measure on $\mathbb{R}$, and if $\mathcal{I}$ is an arbitrary collection of non-degenerate intervals, then for each $\delta > 0$ there exists a finite subcollection, $\mathcal{I}_\delta$, of disjoint intervals in $\mathcal{I}$ such that*

$$\mu\left(\bigcup_{I \in \mathcal{I}} I\right) \leq (2 + \delta) \sum_{I_k \in \mathcal{I}_\delta} \mu(I_k).$$

Below we will want to apply Lemma 2.2 with the measure $u \, dx$, where (u, v) satisfies the $(S_{pq\alpha}^+)$ condition. To do this, we need u to be locally integrable and the intervals to be contained in some compact set. However, if $I = [a, b]$ is an interval such that $u((-\infty, a)) > 0$, and if there exists an interval $J = [b, c]$ such that $\sigma(J) > 0$, then for all $x \in I$,

$$M_\alpha^+(\sigma \chi_{I \cup J})(x) \geq \sigma(J)/(c - a)^{1-\alpha}.$$

Hence, by the $(S_{pq\alpha}^+)$ condition applied to $I \cup J$,

$$u(I) \leq C \sigma(I)^{q/p} (c - a)^{1-\alpha} / \sigma(J) < \infty.$$

If no such J exists, then $\sigma \equiv 0$ on $[b, \infty)$, so $v \equiv \infty$ on the same set. But if $f \in L^p(v)$, then $f \equiv 0$ on $[b, \infty)$.

Below, we will apply Lemma 2.2 to closed intervals contained in an open set O_k . In O_k , $M_\alpha^+ f > 0$, so f cannot be identically zero to the right of these intervals. Further, by the definition of O_k , u is not identically zero to the left of these intervals. Finally, since we will also be assuming that f has compact support, the intervals will be contained in some compact set. Hence Lemma 2.2 is applicable.

The last lemma is an extension of a result of Muckenhoupt [4] for Lebesgue measure. The proof of the extension is identical to his proof and so is omitted.

Lemma 2.3. *Let μ be a Borel measure, f a function, and $\{I_\beta\}$ a collection of intervals, all contained in some interval I , with the property that*

$$\int_{I_\beta} f \, d\mu \geq N \mu(I_\beta).$$

If $J = \bigcup_\beta I_\beta$ then

$$\int_J f \, d\mu \geq (N/2) \mu(J).$$

If (u, v) satisfy the $(S_{pq\alpha}^+)$ condition, then $\sigma \, dx = v^{1-p'} \, dx$ is a Borel measure.

3. PROOF OF THEOREM 1.1

To show that the $(S_{pq\alpha}^+)$ condition is necessary for inequality (1) of Theorem 1.1 to hold, first suppose that there is some interval $I = [a, b]$ such that $u((-\infty, a)) > 0$ but $\sigma(I) = \infty$. Equivalently, the function $v^{-1} \chi_I$ is not in $L^{p'}(v)$, so there exists a function f in $L^p(v)$ such that

$$\infty = \int_I f v^{-1} v \, dx = \int_I f \, dx.$$

Then for all $x \in J$, $M_\alpha^+ f(x) = \infty$, which contradicts inequality (1). The rest of the $(S_{pq\alpha}^+)$ condition follows if we substitute $f = \sigma \chi_I$ into the norm inequality.

To prove that the $(S_{pq\alpha}^+)$ condition is sufficient, we follow the outline of the proof of Martín-Reyes and de la Torre [3], which in turn is based on a proof by Sawyer [6]. Let f be in $L^p(v)$; we will first consider the special case where f is a non-negative,

bounded, upper semicontinuous function of compact support. By Lemma 2.1, $M_\alpha^+ f$ is continuous. Further, since the set $\{x : M_\alpha^+ f(x) = \lambda\}$ has positive measure for at most a countable number of λ , by multiplying f by a suitable constant we may assume without loss of generality that the sets $\{x : M_\alpha^+ f(x) = 2^k\}$ have measure zero for all integers k .

Let $a = \sup\{x : u((-\infty, x)) = 0\}$. For each integer k define the set $O_k = \{x : 2^k < M_\alpha^+ f(x) < 2^{k+1}\} \cap (a, \infty)$. Since $M_\alpha^+ f$ is continuous, it follows that each O_k is open, and the set $\mathbb{R} \setminus \bigcup_k O_k$ has measure zero. For each $x \in O_k$, there exists an open interval $J_{xk} = (x, t_x)$ such that

$$(2) \quad 2^k < \frac{1}{|J_{xk}|^{1-\alpha}} \int_{J_{xk}} f \, dy < 2^{k+1}.$$

We claim that there exists a point $s_x \in J_{xk}$ such that if $y \in I_{xk} = [x, s_x]$ then

$$(3) \quad \frac{1}{|J_{xk}|^{1-\alpha}} \int_{J_{xk}} \sigma \, dy \leq 2M_\alpha^+(\sigma \chi_{J_{xk}})(y).$$

If $M_\alpha^+(\sigma \chi_{J_{xk}})(x) = 0$ then this is immediate. If it is positive, then since $M_\alpha^+(\sigma \chi_{J_{xk}})$ is lower semicontinuous, we can find s_x such that $M_\alpha^+(\sigma \chi_{J_{xk}})(y)$ is also positive for $y \in I_{xk}$. By the continuity of the integral, the desired inequality holds if we take s_x sufficiently close to x . Finally, since O_k is open we may take s_x so that $I_{xk} \subset O_k$.

The union of the I_{xk} 's is O_k . Therefore, by Lemma 2.2 and the remarks following it, there exists a finite, disjoint collection of intervals $\{I_{jk}\}_{j=1}^{n_k}$ such that

$$(4) \quad u(O_k) \leq 3 \sum_{j=1}^{n_k} u(I_{jk}).$$

Since the sets O_k are disjoint, the intervals I_{jk} are pairwise disjoint for all j and k .

Using inequalities (2) and (4), we can now make the following estimate:

$$\begin{aligned} \int_{\mathbb{R}} (M_\alpha^+ f)^q u \, dx &= \sum_k \int_{O_k} (M_\alpha^+ f)^q u \, dx \\ &\leq \sum_k u(O_k) 2^{q(k+1)} \\ &\leq C \sum_{j,k} u(I_{jk}) 2^{qk} \\ &\leq C \sum_{j,k} u(I_{jk}) \left(\frac{1}{|J_{jk}|^{1-\alpha}} \int_{J_{jk}} f \, dx \right)^q \\ &= C \sum_{j,k} u(I_{jk}) \left(\frac{1}{|J_{jk}|^{1-\alpha}} \int_{J_{jk}} \sigma \, dx \right)^q \left(\frac{\int_{J_{jk}} (f/\sigma) \cdot \sigma \, dx}{\int_{J_{jk}} \sigma \, dx} \right)^q \end{aligned}$$

Define the measure ω on $X = \mathbb{N} \times \mathbb{Z}$ by

$$\omega(j, k) = u(I_{jk}) \left(\frac{1}{|J_{jk}|^{1-\alpha}} \int_{J_{jk}} \sigma \, dx \right)^q$$

if $j \leq n_k$, and $\omega(j, k) = 0$ if $j > n_k$. Also define the operator T by

$$Th(j, k) = \frac{\int_{J_{j,k}} |h| \sigma \, dx}{\int_{J_{j,k}} \sigma \, dx}.$$

Then, following the argument of Sawyer, to get the desired norm inequality it will suffice to show that T is a bounded operator from $L^p(\sigma)$ into $L^q(X, \omega)$. Since T is bounded on L^∞ , by Marcinkiewicz interpolation it will suffice to show that T is weak-type $(1, q/p)$: that is, for each $\lambda > 0$

$$\sum_{(j,k) \in E_\lambda} u(I_{j,k}) \left(\frac{1}{|J_{j,k}|^{1-\alpha}} \int_{J_{j,k}} \sigma \, dx \right)^q \leq C \left(\frac{1}{\lambda} \int_{\mathbb{R}} |h| \sigma \, dx \right)^{q/p},$$

where $E_\lambda = \{(j, k) \in X : Th(j, k) > \lambda\}$. If $(j, k) \in E_\lambda$, then

$$\int_{J_{j,k}} |h| \sigma \, dx > \lambda \int_{J_{j,k}} \sigma \, dx.$$

Let G_λ be the union of all such $J_{j,k}$'s. Then, since the $J_{j,k}$'s are open, G_λ is the union of a countable number of disjoint open intervals J_i . By Lemma 2.3, for each i

$$(5) \quad \int_{J_i} |h| \sigma \, dx > \frac{\lambda}{2} \int_{J_i} \sigma \, dx.$$

Since $I_{j,k} \subset J_{j,k}$, each $I_{j,k}$ is contained in exactly one interval J_i . (Here we ignore the left endpoints of the $I_{j,k}$'s since they form a set of measure zero.) Therefore, by inequalities (3) and (5) and the $(S_{p,q,\alpha}^+)$ condition, since $q/p \geq 1$, and since the $I_{j,k}$'s and the J_i 's are disjoint,

$$\begin{aligned} \sum_{(j,k) \in E_\lambda} u(I_{j,k}) \left(\frac{1}{|J_{j,k}|^{1-\alpha}} \int_{J_{j,k}} \sigma \, dx \right)^q &\leq 2 \sum_{(j,k) \in E_\lambda} \int_{I_{j,k}} M_\alpha^+(\sigma \chi_{J_{j,k}})^q u \, dx \\ &\leq 2 \sum_i \sum_{I_{j,k} \subset J_i} \int_{I_{j,k}} M_\alpha^+(\sigma \chi_{J_i})^q u \, dx \\ &\leq 2 \sum_i \int_{J_i} M_\alpha^+(\sigma \chi_{J_i})^q u \, dx \\ &\leq C \sum_i \left(\int_{J_i} \sigma \, dx \right)^{q/p} \\ &\leq C \sum_i \left(\frac{1}{\lambda} \int_{J_i} |h| \sigma \, dx \right)^{q/p} \\ &\leq C \left(\frac{1}{\lambda} \int_{\mathbb{R}} |h| \sigma \, dx \right)^{q/p}. \end{aligned}$$

This completes the special case. To complete the proof, take any $f \in L^p(v)$; then by the Vitali-Carathéodory theorem, there exists an increasing sequence $\{f_n\}$ of non-negative, bounded, upper semicontinuous functions of compact support which converge to $|f|$. (See, for example, Rudin [5, p. 57].) By the monotone convergence theorem, $M_\alpha^+ f_n$ increases pointwise to $M_\alpha^+ f$. Therefore, again by the monotone

convergence theorem,

$$\begin{aligned} \left(\int_{\mathbb{R}} (M_{\alpha}^{+} f)^q u \, dx \right)^{1/q} &= \lim_{n \rightarrow \infty} \left(\int_{\mathbb{R}} (M_{\alpha}^{+} f_n)^q u \, dx \right)^{1/q} \\ &\leq \lim_{n \rightarrow \infty} C \left(\int_{\mathbb{R}} f_n^p v \, dx \right)^{1/p} \\ &= C \left(\int_{\mathbb{R}} |f|^p v \, dx \right)^{1/p}. \end{aligned}$$

ACKNOWLEDGMENT

I want to thank F. Martín-Reyes and A. de la Torre for their hospitality, and my wife Gabrielle for her patient generosity; without either this paper would not have been possible.

REFERENCES

- [1] K. F. Andersen and E. T. Sawyer, *Weighted norm inequalities for the Riemann-Liouville and Weyl fractional integral operators*, Trans. Amer. Math. Soc. 308 (1988), 547-557. MR **89h**:26006
- [2] J. Bliedtner and P. Loeb, *A reduction technique for limit theorems in analysis and probability theory*, Ark. Mat. 30 (1992), 25-43. MR **93k**:28005
- [3] F. J. Martín-Reyes and A. de la Torre, *Two weight norm inequalities for fractional one-sided maximal operators*, Proc. Amer. Math. Soc. 117 (1993), 483-489. MR **94b**:42010
- [4] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. 165 (1972), 207-226. MR **45**:4261
- [5] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, New York, 1974. MR **49**:8783
- [6] E. T. Sawyer, *A characterization of a two-weight norm inequality for maximal operators*, Studia Math. 75 (1982), 1-11. MR **84i**:42032

DEPARTMENT OF MATHEMATICS, TRINITY COLLEGE, HARTFORD, CONNECTICUT 06106-3100
E-mail address: david.cruzuribe@mail.trincoll.edu