APPROXIMATION OF FIXED POINTS OF A STRICTLY PSEUDOCONTRACTIVE MAPPING

LIWEI LIU

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. A fixed point of the strictly pseudocontractive mapping is obtained as the limit of an iteratively constructed sequence with error estimation in general Banach spaces.

Let X be a Banach space. A mapping T is said to be strictly pseudocontractive if there exists a number $t > 1$ such that the inequality

$$\|x - y\| \leq \|(1 + r)(x - y) - rt(Tx - Ty)\|$$

holds for all $x, y \in D(T)$ and $r > 0$. We denote by J the normalized duality mapping from X to 2^{X^*} given by

$$Jx = \{f^* \in X^* : \|f^*\|^2 = \|x\|^2 = \operatorname{Re}\langle x, f^* \rangle\}.$$

A mapping T is said to be strongly accretive (see, e.g., [1, 2]) if there exists a positive number k such that for each $x, y \in D(T)$ there is $j \in J(x - y)$ such that

$$(1) \quad \langle Tx - Ty, j \rangle \geq k\|x - y\|^2.$$

The inequality (1) implies the inequality

$$\|x - y\| \leq \|x - y + r[(T - kI)x - (T - kI)y]\|.$$

Without loss of generality, we can assume $k \in (0, 1)$.

Lemma ([6]). *Let X be a Banach space, K a subset of X , and $T : K \rightarrow K$. Then T is a strictly pseudocontractive mapping if and only if $I - T$ is a strongly accretive mapping, i.e., the inequality*

$$(2) \quad \|x - y\| \leq \|x - y + r[(I - T - kI)x - (I - T - kI)y]\|$$

holds for any $x, y \in K$ and $r > 0$, where $k = (t - 1)/t$.

In this paper, we prove by the inequality (2) that the Mann iteration process converges strongly to the unique fixed point of a Lipschitzian and strictly pseudocontractive mapping. Our results extend corresponding results of [1, 2, 3, 4, 5] to the general Banach spaces. Furthermore, error estimates are also given.

The set of fixed points of the mapping T is denoted by $F(T)$. The Lipschitzian constant of T is denoted by L (≥ 1).

Received by the editors July 11, 1995 and, in revised form, November 3, 1995.

1991 *Mathematics Subject Classification.* Primary 47H17.

Key words and phrases. Strictly pseudocontractive mapping, iterative process.

Theorem 1. *Let X be a Banach space, and let K be a nonempty closed convex and bounded subset of X . Let $T : K \rightarrow K$ be a Lipschitzian strictly pseudocontractive mapping. If $F(T) \neq \emptyset$, then $\{x_n\} \subset K$ generated by $x_1 \in K$,*

$$(3) \quad x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T x_n$$

with $\{\alpha_n\} \subset (0, 1]$ satisfying

$$\sum_{n=1}^{\infty} \alpha_n = \infty, \quad \alpha_n \rightarrow 0,$$

strongly converges to $q \in F(T)$ and $F(T)$ is a single set.

Proof. Let q be a fixed point of T . Since T is a strictly pseudocontractive mapping, then $I - T$ is strongly accretive, i.e., the inequality (2) holds for any $x, y \in K$ and $r > 0$.

From the definition of $\{x_n\}$, we have

$$\begin{aligned} x_n &= x_{n+1} + \alpha_n x_n - \alpha_n T x_n \\ &= (1 + \alpha_n)x_{n+1} + \alpha_n(I - T - kI)x_{n+1} - (2 - k)\alpha_n x_{n+1} \\ &\quad + \alpha_n x_n + \alpha_n(T x_{n+1} - T x_n) \\ &= (1 + \alpha_n)x_{n+1} + \alpha_n(I - T - kI)x_{n+1} - (2 - k)\alpha_n[(1 - \alpha_n)x_n + \alpha_n T x_n] \\ &\quad + \alpha_n x_n + \alpha_n(T x_{n+1} - T x_n) \\ &= (1 + \alpha_n)x_{n+1} + \alpha_n(I - T - kI)x_{n+1} - (1 - k)\alpha_n x_n + (2 - k)\alpha_n^2(x_n - T x_n) \\ &\quad + \alpha_n(T x_{n+1} - T x_n). \end{aligned}$$

By $Tq = q$, we have

$$\begin{aligned} x_n - q &= (1 + \alpha_n)(x_{n+1} - q) + \alpha_n(I - T - kI)(x_{n+1} - q) - (1 - k)\alpha_n(x_n - q) \\ &\quad + (2 - k)\alpha_n^2(x_n - T x_n) + \alpha_n(T x_{n+1} - T x_n). \end{aligned}$$

By using the inequality (2), we obtain

$$\begin{aligned} \|x_n - q\| &\geq (1 + \alpha_n)\|x_{n+1} - q\| - (1 - k)\alpha_n\|x_n - q\| - (2 - k)\alpha_n^2\|x_n - T x_n\| \\ &\quad - \alpha_n\|T x_{n+1} - T x_n\|. \end{aligned}$$

Since

$$\|T x_{n+1} - T x_n\| \leq L\|x_{n+1} - x_n\| \leq L(L + 1)\alpha_n\|x_n - q\|,$$

then

$$\begin{aligned} \|x_n - q\| &\geq (1 + \alpha_n)\|x_{n+1} - q\| - (1 - k)\alpha_n\|x_n - q\| - (2 - k)\alpha_n^2\|x_n - T x_n\| \\ &\quad - L(L + 1)\alpha_n^2\|x_n - q\|. \end{aligned}$$

So,

$$\begin{aligned}
 \|x_{n+1} - q\| &\leq [1 + (1 - k)\alpha_n](1 + \alpha_n)^{-1}\|x_n - q\| \\
 &\quad + (2 - k)\alpha_n^2(1 + \alpha_n)^{-1}\|x_n - Tx_n\| \\
 &\quad + L(L + 1)\alpha_n^2(1 + \alpha_n)^{-1}\|x_n - q\| \\
 (4) \quad &\leq [1 + (1 - k)\alpha_n](1 - \alpha_n + \alpha_n^2)\|x_n - q\| + (2 - k)\alpha_n^2\|x_n - Tx_n\| \\
 &\quad + L(L + 1)\alpha_n^2\|x_n - q\| \\
 &\leq (1 - k\alpha_n)\|x_n - q\| + M\alpha_n^2
 \end{aligned}$$

for some constant $M > 0$, since K is bounded. It follows from the Lemma of Dunn [7, p. 41] that the sequence $\{x_n\}$ strongly converges to the unique fixed point q . This completes the proof.

Remark. The nonlipschitzian version of Theorem 1 was proved in [1, 3] under the assumption that X is a uniformly smooth Banach space by using the inequality

$$\|x + y\|^2 \leq \|x\|^2 + 2\operatorname{Re}\langle y, Jx \rangle + \max\{\|x\|, 1\}\|y\|\beta(\|y\|).$$

Theorem 2. Let K and T be as in Theorem 1. If $\alpha_n = \frac{k}{2(3+3L+L^2)}$, where $k = (t-1)/t$, $\{q\} = F(T)$, then the sequence $\{x_n\}$ generated by (3) converges strongly to the unique fixed point of T , and we have the estimate

$$\|x_{n+1} - q\| < \rho^n \|x_1 - q\|,$$

where $\rho = 1 - \frac{k^2}{4(3+3L+L^2)}$.

Proof. Since $0 < \alpha_n < 1$ and $Tq = q$,

$$\begin{aligned}
 [1 + (1 - k)\alpha_n](1 - \alpha_n + \alpha_n^2) &= 1 - k\alpha_n + \alpha_n^2 - (1 - k)\alpha_n^2(1 - \alpha_n) \\
 &\leq 1 - k\alpha_n + \alpha_n^2,
 \end{aligned}$$

$$\|x_n - Tx_n\| \leq (1 + L)\|x_n - q\|.$$

By (4), we obtain

$$\begin{aligned}
 \|x_{n+1} - q\| &\leq (1 - k\alpha_n)\|x_n - q\| + [1 + (2 - k)(1 + L) + L(L + 1)]\alpha_n^2\|x_n - q\| \\
 &< [1 - k\alpha_n + (3 + 3L + L^2)\alpha_n^2]\|x_n - q\| \\
 &= \left[1 - \frac{k^2}{2(3 + 3L + L^2)} + \frac{k^2}{4(3 + 3L + L^2)}\right]\|x_n - q\| \\
 &= \left[1 - \frac{k^2}{4(3 + 3L + L^2)}\right]\|x_n - q\| \\
 &= \rho\|x_n - q\|.
 \end{aligned}$$

Hence $\|x_{n+1} - q\| < \rho^n \|x_1 - q\|$. The proof is complete.

REFERENCES

1. X. Weng, *Fixed point iteration for local strictly pseudocontractive mapping*, Proc. Amer. Math. Soc. **113** (1991), 727–731. MR **92b**:47099
2. C. E. Chidume, *Iterative approximation of fixed points of Lipschitzian strictly pseudocontractive mappings*, Proc. Amer. Math. Soc. **99** (1987), 283–288. MR **87m**:47122
3. ———, *Approximation of fixed points of strongly pseudocontractive mappings*, Proc. Amer. Math. Soc. **120** (1994), 545–550. MR **94d**:47056

4. ———, *An iterative process for nonlinear Lipschitzian strongly accretive mappings in L_p spaces*, J. Math. Anal. Appl. **151** (1990), 453–461. MR **91k**:47131
5. K-K. Tan and H-K. Xu, *Iterative solutions to nonlinear equation of strongly accretive operators in Banach spaces*, J. Math. Anal. Appl. **178** (1993), 9–21. MR **94g**:47085
6. J. Bogin, *On strict pseudo—contractions and a fixed point theorem*, Technion preprint series No. MT-219, Haifa, Israel, 1974.
7. J. C. Dunn, *Iterative construction of fixed points for multi-valued operators of the monotone type*, J. Funct. Anal. **27** (1978), 30–50. MR **81f**:47056

DEPARTMENT OF MATHEMATICS, NANCHANG UNIVERSITY, NANCHANG, JIANGXI 330047, PEOPLE'S REPUBLIC OF CHINA