***J*-HOLOMORPHIC CURVES IN ALMOST COMPLEX SURFACES DO NOT ALWAYS MINIMIZE THE GENUS**

G. MIKHALKIN

(Communicated by Ronald Stern)

ABSTRACT. The adjunction formula computes the genus of an almost complex curve F embedded in an almost complex surface M in terms of the homology class of F . If M is Kähler (or at least symplectic) and the self-intersection of F is non-negative then the genus of any other surface embedded in M and homologous to F is not less than the genus of F (the proof of this statement (which is a generalization of the Thom conjecture for $\mathbb{C}P^2$) was recently given by the Seiberg-Witten theory). This paper shows that the extra assumptions on M are essential for the genus-minimizing properties of embedded almost complex curves.

Let M be a connected smooth 4-manifold with the tangent bundle τM equipped with a fiberwise-linear map $J : \tau M \rightarrow \tau M$ respecting the fibers and such that $J^2 = -1$. In this case M is called an *almost complex surface*. The $\mathbb{R}$ -linear map J makes τM into a 2-dimensional complex bundle and induces an orientation on M . The *canonical class* $K \in H^2(M; \mathbb{Z})$ is the Euler class of the exterior square over $\mathbb{C}$ of τM multiplied by (-1) .

An embedded surface $F \subset M$ is called a *J -holomorphic curve* if its tangent bundle τF is invariant under J . A J -holomorphic curve gets an orientation from J . If F is J -holomorphic then the normal bundle νF can be chosen to be invariant under J , and the direct sum formula for the characteristic classes of bundles produces the *adjunction formula* for the genus $g(F)$ of F :

$$g(F) = 1 + \frac{F.F + K.F}{2},$$

where $F.F$ denotes the self-intersection of F and $K.F$ denotes the result of evaluation of K on F .

Let $E \subset M$ be an orientable surface homologous to F . The genus $g(E)$ of E is not determined by its homology class. However, in the case when M is symplectic with the symplectic form ω compatible to J so that $\omega(x, Jx) \geq 0$ for any $x \in \tau M$ and $F.F$ is non-negative, the adjunction formula turns into the adjunction inequality

$$g(E) \geq g(F) = 1 + \frac{F.F + K.F}{2}$$

recently proven by means of the Seiberg-Witten theory.

The following theorem shows that the adjunction inequality $g(E) \geq g(F)$ does not hold for all almost complex surfaces.

Received by the editors September 22, 1995.

1991 *Mathematics Subject Classification*. Primary 57R95, 53C15.

Theorem 1. *There exist an almost complex surface M and two smooth closed surfaces $E, F \subset M$, $[E] = [F] \neq 0 \in H_2(M)$, such that F is J -holomorphic and $g(E) < g(F)$.*

Proof. Let M be diffeomorphic to $\mathbb{C}P^2 \# \mathbb{C}P^2 \# \mathbb{C}P^2$. This manifold admits an almost complex structure with the canonical class $-(3, 3, 1) \in \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} = H^2(M)$ by Satz 4.6 of [1], since $(3, 3, 1) \cdot (3, 3, 1) = 19 = 2\chi(M) + 3\sigma(M)$. The next two lemmata produce the required E and F . $\square$

Lemma 1. *There exists a deformation of the almost complex structure on M such that $(4, 0, 0) \in H_2(M)$ is realizable by a closed J -holomorphic curve F of genus 3.*

Proof. Let $F \subset M$ be the orientable surface of genus 3 produced by an embedding of a nonsingular quartic curve into the first summand $\mathbb{C}P^2$ of $M \approx \mathbb{C}P^2 \# \mathbb{C}P^2 \# \mathbb{C}P^2$. We may assume (after a small deformation of F) that F is J -holomorphic in a neighbourhood of a point $x \in F$.

The restriction $TM|_F$ of the tangent bundle of M is a trivial 4-dimensional bundle (since $w_1(TM|_F) = 0$ and $w_2(TM|_F) = 0$). Choose a trivialization of $TM|_F$. The subbundle $TF \subset TM|_F$ is then given by a map $\alpha : F \rightarrow G_{4,2}$, where $G_{4,2} \approx S^2 \times S^2$ is the Grassmannian of the orientable 2-subspaces in $\mathbb{R}^4$. The homotopy class of such a map is determined by two numbers corresponding to the Euler number of the bundle and the Euler number of the normal bundle.

The J -invariant planes passing through a section of $TF|_{F-\{x\}}$ determine a complex subbundle $B \subset TM|_F$ with the first Chern number $c_1(B) = \chi(F) = -4$. The first Chern number of the bundle normal to B is $16 = F \cdot F$, since $c_1(TM|_F) = (3, 3, 1) \cdot (4, 0, 0) = 12$. Therefore, the map $\beta : F \rightarrow G_{4,2}$ corresponding to B is homotopic to α .

To finish the proof we deform the almost complex structure in the tubular neighbourhood of F by following the homotopy between α and β to make TF J -invariant. $\square$

Lemma 2. *There exists an orientable surface $E \subset M$ of genus 1 realizing the homology class $(4, 0, 0) \in H_2(\mathbb{C}P^2 \# \mathbb{C}P^2 \# \mathbb{C}P^2)$.*

Remark. By the Rokhlin-Hsiang-Szczarba inequality [5], [2] the genus of any embedded surface of homology class $(4, 0, 0)$ is at least 1.

Proof of Lemma 2. A nonsingular quartic curve $C \subset \mathbb{C}P^2$ can be obtained from a union of 4 lines and, therefore, it admits a (-1) -membrane for each of its 3 handles (i.e. a disk $M \subset \mathbb{C}P^2$ normal to C along the curve $M \cap C = \partial M$ coinciding with the cocore of the handle and such that the self-intersection number of M equipped with the framing on ∂M coming from C is -1). Making a connected sum with the pair $(\mathbb{C}P^2, \mathbb{C}P^1)$ (cf. [4]) allows us to make the self-intersection of the membrane into zero and, thus, make an embedding surgery removing the handle along this membrane. After repeating this procedure two times we get the required surface $E \subset \mathbb{C}P^2 \# \mathbb{C}P^2 \# \mathbb{C}P^2 = M$ of genus 1 (cf. getting a $(3, 0)$ -sphere in $\mathbb{C}P^2 \# \mathbb{C}P^2$ in [3]). $\square$

Acknowledgment. I am indebted to Y. Eliashberg for the discussion which led to the proof of Lemma 1.

Remark. As I was informed by the referee, an example similar to the one presented in this paper was given by D. Kotschick in a lecture in Cambridge in 1994.

REFERENCES

- [1] F. Hirzebruch, H. Hopf, *Felder von Flächenelementen in 4-dimensionalen Mannigfaltigkeiten*, Math. Ann. **136** (1958), 156-172. MR **20**:7272
- [2] W. Hsiang, R. Szczarba, *On embedded surfaces in four-manifolds*, Proc. Sympos. Pure Math. **22** (1970), 97-103. MR **49**:4000
- [3] M. Kervaire, J. Milnor, *On 2-spheres in 4-manifolds*, Proc. Nat. Acad. Sci. U.S.A. **47** (1961), 1651-1657. MR **24A**:2968
- [4] G. Mikhalkin, *Surfaces of small genus in connected sums of $\mathbb{C}P^2$ and real algebraic curves with many nests in $\mathbb{R}P^2$* , Contemp. Math. **182** (1995), 73-82. MR **95m**:57050
- [5] V. A. Rokhlin, *Two-dimensional submanifolds of four-dimensional manifolds*, Funktsional. Anal. i Prilozhen. **5**(1) (1971), 48-60. MR **45**:7733

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA M5S 1A1

E-mail address: mihalkin@math.toronto.edu