AMENABLE REPRESENTATIONS AND FINITE INJECTIVE VON NEUMANN ALGEBRAS

ALAIN VALETTE

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let $U(M)$ be the unitary group of a finite, injective von Neumann algebra M . We observe that any subrepresentation of a group representation into $U(M)$ is amenable in the sense of Bekka; this yields short proofs of two known results—one by Robertson, one by Haagerup—concerning group representations into $U(M)$.

A unitary representation π of a group Γ on a Hilbert space $\mathcal{H}_\pi$ is *amenable* if there exists on $\mathcal{B}(\mathcal{H}_\pi)$ an $\text{Ad } \pi$ -invariant state, i.e. a state ϕ on $\mathcal{B}(\mathcal{H}_\pi)$ such that, for any $T \in \mathcal{B}(\mathcal{H}_\pi)$, $g \in \Gamma$:

$$\phi(\pi(g)T\pi(g^{-1})) = \phi(T).$$

This notion was introduced and studied by Bekka in [Be].

In the present paper, M will always denote a finite, injective von Neumann algebra. We start with the observation that, if $\pi(\Gamma)$ is contained in the unitary group $U(M)$, then any subrepresentation of π is amenable. We use this to give short, hopefully new proofs of two known results. The first, due to Robertson ([Ro], Theorem C and remark (4) on p. 554), states that for any representation π of a group Γ with Kazhdan's property (T) into $U(M)$, the closure of $\pi(\Gamma)$ in the L^2 -norm topology on $U(M)$ is compact. The second, due to Haagerup ([Ha], Lemma 2.2), says that for any $n \in \mathbb{N}$, $U_1, U_2, \dots, U_n \in U(M)$ and P a non-zero projector in the commutant M' of M :

$$\left\| \sum_{i=1}^n PU_i \otimes \bar{P}\bar{U}_i \right\| = n$$

(where the bar denotes the same operator, but acting on the conjugate Hilbert space).

Proposition 1. *Let π be a representation of a group Γ into $U(M)$. Then any subrepresentation of π is amenable.*

Proof. We may assume that $M = \pi(\Gamma)''$. Let ρ be a subrepresentation of π on a closed subspace $\mathcal{H}_\rho$ which is the range of a projector $p \in M'$. To construct an $\text{Ad } \rho$ -invariant state on $\mathcal{B}(\mathcal{H}_\rho) = p\mathcal{B}(\mathcal{H}_\pi)p$, choose a conditional expectation

Received by the editors October 6, 1995 and, in revised form, December 5, 1995.

1991 *Mathematics Subject Classification.* Primary 22D25; Secondary 46L10.

Key words and phrases. Amenable representations, finite injective von Neumann algebra, Kazhdan's property (T).

$E: \mathcal{B}(\mathcal{H}_\rho) \rightarrow pM$, a trace τ on pM , and set $\phi = \tau \circ E$. Then, for any $T \in \mathcal{B}(\mathcal{H}_\rho)$, $g \in \Gamma$:

$$\phi(\rho(g)T\rho(g^{-1})) = \tau(\rho(g)E(T)\rho(g^{-1})) = \phi(T),$$

which concludes the proof. $\square$

For a representation π of Γ into $U(M)$, the closure of $\pi(\Gamma)$ in $U(M)$ is compact in the L^2 -norm if and only if π decomposes as a direct sum of finite-dimensional representations. Indeed, if $\pi(\Gamma)$ is relatively compact, the identity representation of the compact group $\overline{\pi(\Gamma)}$ decomposes into finite-dimensional representations; conversely, using the fact that the L^2 and strong topologies coincide on $U(M)$, it is easy to see that any unitary representation that decomposes into finite-dimensional ones, has relatively compact range. We then state Robertson's result mentioned above in the following equivalent form.

Proposition 2. *Let Γ be a group with Kazhdan's property (T). Any representation π of Γ into $U(M)$ decomposes as a direct sum of finite-dimensional representations.*

Proof. By Zorn's lemma, find a subrepresentation ρ of π that decomposes as a direct sum of finite-dimensional representations, and maximal with respect to that property. We have to show that $\rho = \pi$. If this is not the case, consider the subrepresentation $\rho^\perp$ on the orthogonal complement of $\mathcal{H}_\rho$. By Proposition 1, $\rho^\perp$ is an amenable representation of Γ . Because Γ has property (T), it follows from Corollary 5.9 of [Be] that any amenable representation of Γ has a (non-zero) finite-dimensional subrepresentation. This, applied to $\rho^\perp$, contradicts maximality of ρ . $\square$

Remarks. (1) Proposition 2 makes precise an earlier result of Kirchberg ([Ki], Corollary 1.2): any Kazhdan group that admits a faithful representation into $U(M)$, must be residually finite. Kirchberg also mentions in the same paper [Ki] that his proof of residual finiteness for certain subgroups of $U(M)$ works for a bigger class of subgroups than just subgroups with property (T) (e.g. it works for non-abelian free groups). However, the proof of Proposition 2 does not extend, in view of Theorem 1 of [BV]: Kazhdan groups are characterized by the fact that any amenable representation has a finite-dimensional subrepresentation.

(2) Robertson proved ([Ro], Lemma 4.2) a result more general than our Proposition 2: any representation of a Kazhdan group into the unitary group of a finite von Neumann algebra with Haagerup's approximation property, decomposes as a direct sum of finite-dimensional representations.

We now turn to Haagerup's result mentioned in the beginning.

Proposition 3. *For any $n \in \mathbb{N}$, $U_1, U_2, \dots, U_n \in U(M)$ and P a non-zero projector in the commutant M' of M :*

$$\left\| \sum_{i=1}^n P U_i \otimes \bar{P} \bar{U}_i \right\| = n.$$

Proof. Let Γ be a finitely generated group, S a finite generating subset, and π a unitary representation of Γ . It follows from Theorem 5.1 of [Be] and remark (b) on p. 74 of [HRV], that π is amenable if and only if 1 belongs to the spectrum of the operator $\frac{1}{|S|} \sum_{s \in S} (\pi \otimes \bar{\pi})(s)$, where $\bar{\pi}$ denotes the contragredient representation of π . If this is the case, then $\| \sum_{s \in S} (\pi \otimes \bar{\pi})(s) \| = |S|$. The result then follows by

considering the U_i 's as generators of a representation of the free group $\mathbb{F}_n$ on n generators, and appealing to Proposition 1. $\square$

Remark. Haagerup has proved that Proposition 3 actually yields a characterization of finite, injective von Neumann algebras. Indeed, it is enough to assume $\|\sum_{i=1}^n PU_i \otimes \bar{P}\bar{U}_i\| = n$ for any n -tuple of unitaries and any non-zero *central* projection, to make sure that M is finite and injective (see [Ha], Lemma 2.2). For a factor acting on a separable Hilbert space, this characterization of the hyperfinite II_1 -factor is due to Connes ([Co], Remark 5.29).

REFERENCES

- [Be] M. E. B. Bekka, *Amenable representations of locally compact groups*, Invent. Math. **100** (1990), 383–401. MR **91g**:22007
- [BV] M. E. B. Bekka & A. Valette, *Kazhdan's property (T) and amenable representations*, Math. Z. **212** (1993), 293–299. MR **94a**:22006
- [Co] A. Connes, *Classification of injective factors*, Ann. of Math. **104** (1976), 585–609.
- [Ha] U. Haagerup, *Injectivity and decomposition of completely bounded maps*, in Operator algebras and their connections with topology and ergodic theory (Proceedings, Busteni, Romania 1983), LNM 1132, Springer 1985, 170–222. MR **87i**:46133
- [HRV] P. de la Harpe, A. G. Robertson & A. Valette, *On the spectrum of the sum of generators for a finitely generated group*, Israel J. Math. **81** (1993), 65–96. MR **94j**:22007
- [Ki] E. Kirchberg, *Discrete groups with Kazhdan's property (T) and factorization property are residually finite*, Math. Ann. **299** (1994), 551–563. MR **95h**:46083
- [Ro] A. G. Robertson, *Property (T) for II_1 -factors and unitary representations of Kazhdan groups*, Math. Ann. **296** (1993), 547–555. MR **94f**:22006

INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ DE NEUCHÂTEL, RUE EMILE ARGAND 11, CH-2007 NEUCHÂTEL, SWITZERLAND

E-mail address: valette@maths.unine.ch