APPROXIMATION OF FIXED POINTS OF STRONGLY PSEUDOCONTRACTIVE MAPS WITHOUT LIPSCHITZ ASSUMPTION

ZHOU HAIYUN AND JIA YUTING

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. In the present paper, the following result is shown: Let X be a real Banach space with a uniformly convex dual X^* , and let K be a nonempty closed convex and bounded subset of X . Assume that $T : K \rightarrow K$ is a continuous strong pseudocontraction. Let $\{\alpha_n\}_{n=1}^\infty$ and $\{\beta_n\}_{n=1}^\infty$ be two real sequences satisfying

- (i) $0 < \alpha_n, \beta_n < 1$ for all $n \geq 1$;
- (ii) $\sum_{n=1}^\infty \alpha_n = \infty$; and
- (iii) $\alpha_n \rightarrow 0, \beta_n \rightarrow 0$ as $n \rightarrow \infty$.

Then the Ishikawa iterative sequence $\{x_n\}_{n=1}^\infty$ generated by

$$(I) \begin{cases} x_1 \in K, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T x_n, n \geq 1, \end{cases}$$

converges strongly to the unique fixed point of T .

1. INTRODUCTION AND PRELIMINARIES

Let X be a normed linear space, $K \subseteq X$. A mapping $T : K \rightarrow K$ is said to be strongly pseudocontractive if there exists $t > 1$ such that the inequality

$$(1) \quad \|x - y\| \leq \|(1 + r)(x - y) - rt(Tx - Ty)\|$$

holds for every $x, y \in K$ and $r > 0$. A mapping U with domain $D(U)$ and range $R(U)$ in X is called accretive if the following inequality

$$(2) \quad \|x - y\| \leq \|x - y + s(Ux - Uy)\|$$

holds for every $x, y \in D(U)$ and for all $s \geq 0$. Browder [1, Prop. 1] proved that T is pseudocontractive if and only if $(I - T)$ is accretive, where I denotes the identity operator.

Let X be a real Banach space with a dual X^* . The normalized duality mapping $J : X \rightarrow 2^{X^*}$ is defined by

$$J(x) = \{f \in X^*; \langle x, f \rangle = \|x\|^2, \|x\| = \|f\|\},$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. It is well known that if X^* is strictly convex, then J is single valued and such that $J(-x) = -J(x)$, $J(tx) =$

Received by the editors December 5, 1995.

1991 *Mathematics Subject Classification*. Primary 47H17.

Key words and phrases. The Ishikawa iteration, strong pseudocontraction, strictly convex Banach space.

$tJ(x)$ for all $t \geq 0$, $x \in X$. If X^* is uniformly convex, then J is uniformly continuous on the bounded sets of X (see, e.g., Barbu [2])

By means of the normalized duality mapping J , we can give the following equivalent definition for accretive mapping.

A mapping $A : K \rightarrow K$ is said to be accretive if for each x, y in K there exists $j \in J(x - y)$ such that

$$(3) \quad \langle Ax - Ay, j \rangle \geq 0.$$

A mapping $A : K \rightarrow K$ is called strongly accretive if for each $x, y \in K$ there exists $j \in J(x - y)$ such that

$$(4) \quad \langle Ax - Ay, j \rangle \geq k\|x - y\|^2$$

for some constant $k > 0$. Without loss of generality we shall assume that $k \in (0, 1)$.

In [3], Bogin proved that U is strongly pseudocontractive if and only if $I - U$ is strongly accretive.

Very recently, Chidume [4] proved the following result: If X is a real Banach space with a uniformly convex dual X^* , and K is a nonempty closed convex and bounded subset of X , and $T : K \rightarrow K$ is a Lipschitz strongly pseudocontractive mapping, the Ishikawa iterative sequence $\{x_n\}_{n=1}^\infty$ generated by (I) converges strongly to the unique fixed point of T . However, when T is continuous strongly pseudocontractive, it is still an open question whether or not the Ishikawa iterative sequence $\{x_n\}_{n=1}^\infty$ defined by (I) converges strongly to a fixed point of T .

It is our purpose in this paper to solve the above question by proving the following much more general result: If X is a real Banach space with a uniformly convex dual X^* , K is a nonempty bounded closed convex subset of X , and $T : K \rightarrow K$ is a continuous strongly pseudocontractive mapping, then the Ishikawa iteration sequence $\{x_n\}_{n=1}^\infty$ defined by (I) converges strongly to the unique fixed point of T .

We need the following lemmas.

Lemma 1.1. *Let X^* be strictly convex; then*

$$(5) \quad \|x + y\|^2 \leq \|x\|^2 + 2\langle y, J(x + y) \rangle$$

for all $x, y \in X$.

Proof. For arbitrary fixed $x, y \in X$, let $\varphi(s) = \|x + sy\|^2$, $s \in [0, +\infty)$; then $\varphi : [0, +\infty) \rightarrow [0, \infty)$ is a convex function.

To see this, for every $s_1, s_2 \in [0, +\infty)$, $t \in [0, 1]$, we have

$$\begin{aligned} \varphi(ts_1 + (1-t)s_2) &= \|t(x + s_1y) + (1-t)(x + s_2y)\|^2 \\ &\leq t^2\|x + s_1y\|^2 + (1-t)^2\|x + s_2y\|^2 + 2t(1-t)\|x + s_1y\|\|x + s_2y\| \\ &\leq (t^2 + t(1-t))\|x + s_1y\|^2 + ((1-t)^2 + t(1-t))\|x + s_2y\|^2 \\ &= t\varphi(s_1) + (1-t)\varphi(s_2). \end{aligned}$$

Since $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ is a convex function, as $s > t > 0$, we have

$$\begin{aligned} \varphi(t) &= \varphi\left(\frac{t}{s} \cdot s + \left(1 - \frac{t}{s}\right) \cdot 0\right) \\ &\leq \frac{t}{s}\varphi(s) + \left(1 - \frac{t}{s}\right)\varphi(0) \\ &= \frac{t}{s}(\varphi(s) - \varphi(0)) + \varphi(0), \end{aligned}$$

and then

$$t^{-1}(\varphi(t) - \varphi(0)) \leq s^{-1}(\varphi(s) - \varphi(0)).$$

Hence, function $t \mapsto t^{-1}(\varphi(t) - \varphi(0))$ is increasing. As a result, $\lim_{t \rightarrow 0^+} t^{-1}(\varphi(t) - \varphi(0))$ exists and

$$\|x + y\|^2 - \|x\|^2 \geq \lim_{t \rightarrow 0^+} t^{-1}(\varphi(t) - \varphi(0)) = \langle y, \text{grad}\|x\|^2 \rangle = 2\langle y, J(x) \rangle,$$

so that, for all $x, y \in X$ we have

$$(6) \quad \|x + y\|^2 \geq \|x\|^2 + 2\langle y, J(x) \rangle.$$

Now by replacing x by $x - y$ in (6) we get

$$(7) \quad \|x\|^2 \geq \|x - y\|^2 + 2\langle y, J(x - y) \rangle.$$

Replacing x by $-x$ in (7), we obtain

$$(8) \quad \|x\|^2 \geq \|x + y\|^2 - 2\langle y, J(x + y) \rangle.$$

Thus, we have

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, J(x + y) \rangle$$

for all $x, y \in X$. □

Lemma 1.2. *Let $\{\rho_n\}_{n=1}^\infty$ be a nonnegative real sequence satisfying*

$$(9) \quad \rho_{n+1} \leq (1 - \lambda_n)\rho_n + \sigma_n,$$

where $\lambda_n \in [0, 1]$, $\sum_{n=1}^\infty \lambda_n = \infty$, and $\sigma_n = o(\lambda_n)$. Then $\rho_n \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Let $\alpha = \inf\{\rho_n : n \geq 1\}$; then $\alpha = 0$. If not, assume that $\alpha > 0$; then for all $n \geq 1$, $\rho_n \geq \alpha > 0$.

Since $\sigma_n = o(\lambda_n)$ we know that

$$\sigma_n < \frac{1}{2}\rho_n\lambda_n$$

for all large n .

From (9), we get

$$\rho_{n+1} < (1 - \lambda_n)\rho_n + \frac{1}{2}\rho_n\lambda_n = (1 - \frac{1}{2}\lambda_n)\rho_n.$$

By a simple iteration we get that

$$0 \leq \rho_{n+1} \leq \exp\left\{-\frac{1}{2}\sum_{j=1}^n \lambda_j\right\}\rho_1 \rightarrow 0 \text{ as } n \rightarrow \infty.$$

i.e., a contradiction. Thus there exists some subsequence of $\{\rho_n\}$, say $\{\rho_{n_j}\}$, such that $\rho_{n_j} \rightarrow 0$ as $j \rightarrow \infty$. Hence for arbitrary given $\epsilon > 0$, there exists large integer $j_0 \geq 1$ such that

$$\sigma_n < \lambda_n\epsilon, \quad \rho_{n_{j_0}} < \epsilon$$

for all large $n \geq n_{j_0}$.

By virtue of (9) we have

$$\rho_{n_{j_0}+1} \leq (1 - \lambda_{n_{j_0}})\epsilon + \lambda_{n_{j_0}}\epsilon = \epsilon;$$

by induction, we have

$$\rho_{n_{j_0}+k} \leq \epsilon, \text{ for all } k \geq 1.$$

This shows $\rho_n \rightarrow 0$ as $n \rightarrow \infty$. □

2. MAIN RESULTS

Now we prove the main results.

Theorem 2.1. *Let X be a real Banach space with a uniformly convex dual X^* , K be a nonempty closed convex bounded subset of X , and $T : K \rightarrow K$ be a continuous strongly pseudocontractive mapping. Then the Ishikawa iterative sequence $\{x_n\}_{n=1}^\infty$ defined by (I) converges strongly to the unique fixed point of T .*

Proof. The existence of a fixed point follows from Deimling [5]. Let q be a fixed point of T . Since $T : K \rightarrow K$ is strongly pseudocontractive, $I - T$ is strongly accretive, and for every $x, y \in K$

$$(10) \quad \langle (I - T)x - (I - T)y, J(x - y) \rangle \geq k\|x - y\|^2,$$

where $k = (t - 1)t^{-1}$ (see, e.g., [3]).

From (10) we obtain

$$(11) \quad \langle Tx - Ty, J(x - y) \rangle \leq (1 - k)\|x - y\|^2$$

for all $x, y \in K$.

Using (5), (11) and (I), we get

$$(12) \quad \begin{aligned} \|x_{n+1} - q\|^2 &= \|(1 - \alpha_n)(x_n - q) + \alpha_n(Ty_n - Tq)\|^2 \\ &\leq (1 - \alpha_n)^2\|x_n - q\|^2 + 2\alpha_n\langle Ty_n - Tq, J(x_{n+1} - q) \rangle \\ &= (1 - \alpha_n)^2\|x_n - q\|^2 + 2\alpha_n\langle Ty_n - Tq, J(x_{n+1} - q) - J(y_n - q) \rangle \\ &\quad + 2\alpha_n\langle Ty_n - Tq, J(y_n - q) \rangle \\ &\leq (1 - \alpha_n)^2\|x_n - q\|^2 + 2\alpha_na_n + 2\alpha_n(1 - k)\|y_n - q\|^2, \end{aligned}$$

where

$$a_n = \langle Ty_n - Tq, J(x_{n+1} - q) - J(y_n - q) \rangle.$$

Now we shall show $a_n \rightarrow 0$ as $n \rightarrow \infty$. Observe that $\{Ty_n - Tq\}$ is bounded; then to show $a_n \rightarrow 0$ as $n \rightarrow \infty$, it suffices to show

$$J(x_{n+1} - q) - J(y_n - q) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Indeed, since X^* is uniformly convex, J is uniformly continuous on any bounded subset of X . Noting that $(x_{n+1} - q) - (y_n - q) = x_{n+1} - y_n = (\beta_n - \alpha_n)x_n + \alpha_nTy_n - \beta_n \rightarrow 0$ as $n \rightarrow \infty$, we see that

$$J(x_{n+1} - q) - J(y_n - q) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Again using (5) and (I), we have

$$(13) \quad \begin{aligned} \|y_n - q\|^2 &= \|(1 - \beta_n)(x_n - q) + \beta_n(Tx_n - Tq)\|^2 \\ &\leq (1 - \beta_n)^2\|x_n - q\|^2 + 2\beta_n\langle Tx_n - Tq, J(y_n - q) \rangle \\ &\leq \|x_n - q\|^2 + M_1\beta_n \end{aligned}$$

for some constant $M_1 > 0$, since $\langle Tx_n - Tq, J(y_n - q) \rangle$ is bounded.

Substituting (13) in (12) yields

$$\begin{aligned}
 (14) \quad \|x_{n+1} - q\|^2 &\leq ((1 - \alpha_n)^2 + 2(1 - k)\alpha_n)\|x_n - q\|^2 \\
 &\quad + 2\alpha_n(a_n + (1 - k)M_1\beta_n) \\
 &= (1 - 2\alpha_n + \alpha_n^2 + 2\alpha_n - 2k\alpha_n)\|x_n - q\|^2 \\
 &\quad + 2\alpha_n(a_n + (1 - k)M_1\beta_n) \\
 &= (1 + \alpha_n^2 - 2k\alpha_n)\|x_n - q\|^2 + \sigma_n,
 \end{aligned}$$

where $\sigma_n = 2\alpha_n(a_n + (1 - k)M_1\beta_n)$.

We choose a large positive integer N such that for all $n > N$

$$\alpha_n < k.$$

Thus, the above inequality yields

$$\|x_{n+1} - q\|^2 \leq (1 - k\alpha_n)\|x_n - q\|^2 + \sigma_n.$$

Set $\rho_n = \|x_n - q\|^2$ and $\lambda_n = k\alpha_n$. Then

$$\rho_{n+1} \leq (1 - \lambda_n)\rho_n + \sigma_n,$$

with $\lambda_n \in [0, 1]$, $\sum_{n=1}^{\infty} \lambda_n = \infty$ and $\sigma_n = o(\lambda_n)$. By Lemma 1.2 we see that $\rho_n \rightarrow 0$ as $n \rightarrow \infty$.

Now we prove the uniqueness. Suppose q_1 is another fixed point of T ; then by (10) we have

$$0 = \langle (q - Tq) - (q_1 - Tq_1), J(q - q_1) \rangle \geq k\|q - q_1\|^2,$$

which implies $q_1 = q$.

The proof of this theorem is completed. $\square$

Remark. Our theorem extends Theorem 2 of Chidume [4] to a larger class of continuous strongly pseudocontractive mappings; thus we have solved an open question put forth by Chidume in [4].

REFERENCES

- [1] F. E. Browder, Nonlinear operators and nonlinear equation of evolution in Banach spaces, Proc. Sympos. Pure. Math., 18(1976). MR **53**:8982
- [2] V. Barbu, Nonlinear Semigroups and Differential Equations in Banach Spaces, Noordhoff, Leyden, 1976. MR **52**:11666
- [3] J. Bogin, On strict pseudo-contractions and a fixed point theorem, Technion Preprint MT-29, Haifa, 1974.
- [4] C. E. Chidume, Approximation of fixed points of strongly pseudocontractive mappings, Proc. Amer. Math. Soc. 120, No.2 (1994), 545-551. MR **94d**:47056
- [5] K. DEIMLING, Zeros of accretive operators, Manuscripta Math. 13 (1974), 283-288. MR **50**:3030

DEPARTMENT OF MATHEMATICS, SHIJIAZHUANG ENGINEERING COLLEGE, SHIJIAZHUANG 050003, PEOPLE'S REPUBLIC OF CHINA

DEPARTMENT OF MATHEMATICS, HEBEI TEACHERS UNIVERSITY, SHIJIAZHUANG 050016, PEOPLE'S REPUBLIC OF CHINA