GROUP HOMOMORPHISMS INDUCING $\text{mod-}p$ COHOMOLOGY MONOMORPHISMS

PHAM ANH MINH

(Communicated by Ronald Solomon)

ABSTRACT. Let $f: G \rightarrow K$ be a homomorphism of p -groups such that $f^{(n)}: H^n(K, \mathbf{Z}/p) \rightarrow H^n(G, \mathbf{Z}/p)$ is injective, for $1 \leq n \leq 2$. We prove that the non-bijectivity of f implies the existence of a quotient L of G containing K as a proper direct factor. This gives a refined proof of a result of Evens, which asserts that f is bijective if $f^{(1)}$ is.

Let p be a prime number and let $\mathbf{Z}/p$ be the prime field of p elements. For every p -group K , let us denote by $H^*(K)$ the $\text{mod-}p$ cohomology of K .

Let $f: G \rightarrow K$ be a homomorphism of p -groups such that $f^{(n)}: H^n(K) \rightarrow H^n(G)$ is injective, for $1 \leq n \leq 2$. We shall give a refined proof of the following result of Evens.

Theorem A (Evens [1, Th. 7.2.4]). *If $f^{(1)}$ is bijective, then so is f .*

Further, we prove

Theorem B. *If f is not bijective, then there exists a quotient L of G containing K as a proper direct factor (i.e., $L = J \times K$, with $J \neq \{1\}$).*

Note that the maps $H^1(K) \xrightarrow{\text{Inf}_1(L, K)} H^1(L)$ and $H^1(L) \xrightarrow{\text{Inf}_1(G, L)} H^1(G)$ are injective, and $\text{Im Inf}_1(L, K)$ is a proper subgroup of $H^1(L)$ (since $J \neq \{1\}$), so, by Theorem B, the non-bijectivity of f implies that

$$\dim_{\mathbf{Z}/p} H^1(K) < \dim_{\mathbf{Z}/p} H^1(L) \leq \dim_{\mathbf{Z}/p} H^1(G),$$

hence $f^{(1)}$ is not bijective. We obtain then an alternative proof of Theorem A.

Proof of Theorem B. We first prove that f is surjective. Set $K' = \text{Im } f$. So f factors through $G \xrightarrow{f} K' \hookrightarrow K$. Since $f^{(1)} = (H^1(K) \xrightarrow{\text{Res}} H^1(K') \xrightarrow{f^*} H^1(G))$ is injective, it follows that $\text{Res}: H^1(K) \rightarrow H^1(K')$ is injective. This implies $K = K'$, so f is surjective.

Consider the extension

$$1 \rightarrow \text{Ker } f \rightarrow G \xrightarrow{f} K \rightarrow 1.$$

Let p^m be the order of $\text{Ker } f$ ($m \geq 1$, since f is not bijective). The above extension

Received by the editors September 7, 1995.

1991 *Mathematics Subject Classification.* Primary 20J05.

can then be obtained by successive central extensions

$$(G_i) \quad 0 \rightarrow \mathbf{Z}/p \rightarrow G_i \xrightarrow{p_i} G_{i-1} \rightarrow 1,$$

$1 \leq i \leq m$, with $G_0 = K$, $G_m = G$, $p_1 \circ \cdots \circ p_m = f$.

Let $z \in H^2(K)$ be the factor set of the extension (G_1) , viewed as a cohomology class of $H^2(K)$; then $z \in \text{Ker Inf}_2(G_1, K)$. Since $f^{(2)} = \text{Inf}_2(G, K) = \text{Inf}_2(G, G_1) \circ \text{Inf}_2(G_1, K)$, it follows from the injectivity of $f^{(2)}$ that $\text{Inf}_2(G_1, K)$ is also injective. Hence $z = 0$, so $G_1 = \mathbf{Z}/p \times K$. The theorem follows.

REFERENCES

1. L. Evens, *Cohomology of groups*, Oxford University Press, 1991. MR **93i**:20059

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF HUE, DAI HOC KHOA HOC, HUE, VIETNAM