FORMULAE AND CONTINUITY FOR THE INDEX OF SUBFACTORS

SERGEY DOROFEEV AND KLAUS THOMSEN

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let $N \subset M$ be an inclusion of II_1 -factors, τ the trace state of M , and $\mathcal{P}(M)$, $\mathcal{P}(N)$ the set of projections in M and N , respectively. We prove that the Jones index for the inclusion is

$$\begin{aligned} [M : N] &= \sup_{e \in \mathcal{P}(M) \setminus \{0\}} \inf_{p \in \mathcal{P}(N) \setminus \{0\}} \frac{\tau(p)}{\tau(ep)} \\ &= \sup_{e \in \mathcal{P}(M) \setminus \{0\}} \inf \left\{ \frac{\tau(p)}{\tau(ep)} : p \in \mathcal{P}(N), e \preceq p \right\}. \end{aligned}$$

This formula is exploited to obtain continuity results for the index. In particular, we obtain a formula for the index which expresses $[M : N]$ in terms of the positions of N_i and M_j , $i, j \in \mathbb{N}$, in M , when $N_1 \subset N_2 \subset N_3 \subset \cdots$ and $M_1 \subset M_2 \subset M_3 \subset \cdots$ are finite-dimensional C^* -subalgebras with dense union in N and M , respectively.

It has been shown by Pimsner and Popa in [PP] that the index $[M : N]$ of an inclusion $N \subset M$ of II_1 factors is the reciprocal of

$$\max \{ \lambda \in [0, 1] : E(x) \geq \lambda x, x \in M^+ \},$$

where $E : M \rightarrow N$ is the trace-invariant conditional expectation. This formula is of crucial importance in the sub-factor theory, and it is the basis for most generalizations of the Jones index to other cases. It can also be used, in the setting of ergodic theory, to define the index of a sub- σ -algebra $\mathcal{B}_0$ of measurable subsets in a measure space $(X, \mathcal{B}, \mu)$ with $\mu(X) = 1$. In this setup it served as a main tool in [DT], where it was shown that

$$\begin{aligned} &\max \{ \lambda \in [0, 1] : E_{\mathcal{B}_0}(f) \geq \lambda f, f \in L^\infty(X, \mathcal{B}, \mu)^+ \} \\ &= \max \{ \lambda \in [0, 1] : \forall A \in \mathcal{B} \text{ there is a } C \in \mathcal{B}_0 \text{ such that} \\ &\quad \mu(A \setminus C) = 0 \text{ and } \lambda \mu(C) \leq \mu(A) \}. \end{aligned}$$

The following result shows that this formula also holds in the non-commutative case of II_1 -factors.

Received by the editors January 16, 1996.

1991 *Mathematics Subject Classification*. Primary 46L37.

Theorem 1. *Let M be a II_1 -factor and $N \subset M$ a subfactor and $\mathcal{P}(N), \mathcal{P}(M)$ the projection lattices in N and M , respectively. Then*

$$\begin{aligned} & [M : N]^{-1} \\ &= \max\{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \exists p \in \mathcal{P}(N) \text{ such that } e \preceq p \text{ and } \lambda p \leq pE_N(e)\} \\ &= \max\{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \exists p \in \mathcal{P}(N) \text{ such that } e \preceq p \text{ and } \lambda\tau(p) \leq \tau(ep)\} \\ &= \max\{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \setminus \{0\} \exists p \in \mathcal{P}(N) \setminus \{0\} \text{ such that } \lambda\tau(p) \leq \tau(ep)\} . \end{aligned}$$

Proof. Set $A = \{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \exists p \in \mathcal{P}(N) \text{ such that } e \preceq p \text{ and } \lambda p \leq pE_N(e)\}$, $B = \{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \exists p \in \mathcal{P}(N) \text{ such that } e \preceq p \text{ and } \lambda\tau(p) \leq \tau(ep)\}$ and $C = \{\lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \setminus \{0\} \exists p \in \mathcal{P}(N) \setminus \{0\} \text{ such that } \lambda\tau(p) \leq \tau(ep)\}$. To handle the least interesting case first, assume that $[M : N] = \infty$. In this case we see from [PP], Theorem 2.2, that for all $\epsilon > 0$ there is a non-zero projection $e \in M$ such that $\|E_N(e)\| < \epsilon$. From this it follows straightforwardly that $A = B = C = \{0\}$. Now we assume that $[M : N]^{-1} > 0$. If $\lambda \in A$ and $e \in \mathcal{P}(M)$ we have a $p \in \mathcal{P}(N)$ such that $e \preceq p$ and $\lambda p \leq pE_N(e)$. Then $\lambda\tau(p) \leq \tau(pE_N(e)) = \tau(ep)$. Thus $A \subset B$. It is trivial that $B \subset C$. Let $\mu \in C$. By [J] and [PP] there is a projection $e \in M$ such that $E_N(e) = [M : N]^{-1}1$. In particular, $e \neq 0$ and we have a non-zero projection $p \in N$ such that $\mu\tau(p) \leq \tau(ep) = \tau(E_N(e)p) = [M : N]^{-1}\tau(p)$, showing that $\mu \leq [M : N]^{-1}$. Thus $\sup A \leq \sup B \leq \sup C \leq [M : N]^{-1}$ and the proof can be completed by showing that $[M : N]^{-1} \in A$.

Set $\lambda = [M : N]^{-1}$. By Proposition 2.1 of [PP] we have that $E_N(x) \geq \lambda x$ for all $x \in M_+$. The proof that $\lambda \in A$ is based on this and the following simple observation:

- (1) When $a \in M_+$, $p \in \mathcal{P}(M) \setminus \{0\}$, $s, t \in [0, \infty[$
such $s < t$ and $tp \leq a$, then the spectral projection of a
corresponding to the interval $[s, \infty[$ is non-zero.

Consider $e \in \mathcal{P}(M) \setminus \{0\}$ and fix $n \in \mathbb{N}$. Since $(\lambda + \frac{1}{n})e \leq E_N(e) + \frac{1}{n}1$ we see from (1) that there is a non-zero projection q in the abelian von Neumann algebra $\mathcal{A}$ generated by $E_N(e)$ and 1 such that $q(E_N(e) + \frac{1}{n}1) \geq \lambda q$. Let $\{q_i\}$ be a maximal family of non-zero mutually orthogonal projections in $\mathcal{A}$ such that $q_i(E_N(e) + \frac{1}{n}1) \geq \lambda q_i$ for all i . Set $f_n = \sum_i q_i \in \mathcal{A}$ and note that $f_n(E_N(e) + \frac{1}{n}1) \geq \lambda f_n$. Note that

$$\begin{aligned} (\lambda + \frac{1}{n})e \wedge (1 - f_n) &\leq \lambda(1 - f_n)e(1 - f_n) + \frac{1}{n}(1 - f_n) \\ &\leq (1 - f_n)(E_N(e) + \frac{1}{n}1)(1 - f_n) \end{aligned}$$

which shows that $e \wedge (1 - f_n) = 0$, since otherwise (1) would give us a non-zero projection q orthogonal to f_n such that $q(E_N(e) + \frac{1}{n}1) \geq \lambda q$, contradicting the maximality of $\{q_i\}$. Now the parallelogram rule shows that $e \sim f_n - (1 - e) \wedge f_n \leq f_n$, so that $\tau(e) \leq \tau(f_n)$. Set $p_n = \bigvee_{i=n}^{\infty} f_i$ and observe that $p_n(E_N(e) + \frac{1}{n}1) \geq \lambda p_n$ and that $\tau(p_n) \geq \tau(f_n) \geq \tau(e)$. Since $\{p_n\}$ is a decreasing sequence of projections the strong limit $p = \lim_{n \rightarrow \infty} p_n$ is a projection. As $\tau(p) = \lim_n \tau(p_n) \geq \tau(e)$ we have that $e \preceq p$. Since $p_n(E_N(e) + \frac{1}{n}1) \geq \lambda p_n$ for all n we find that $pE_N(e) \geq \lambda p$. $\square$

By taking the reciprocal value of the last two expressions of Theorem 1 we get the formulae of the abstract for the index.

One virtue of the alternative expressions for the index obtained in Theorem 1 is that they have nice continuity properties. To illustrate this we prove two results where this is exploited. Consider a type II_1 factor B with trace state τ and two von Neumann subalgebras, N and M , of B . This setting will be fixed throughout the rest of this note. Set

$$\lambda(M, N) = \sup \{ \lambda \in [0, 1] : \forall e \in \mathcal{P}(M) \exists p \in \mathcal{P}(N) \text{ such that } \tau(e) \leq \tau(p) \text{ and } \lambda \tau(p) \leq \tau(ep) \}.$$

By Theorem 1 we have that $\lambda(M, N)^{-1} = [M : N]$ when $M = B$ and N is a factor.

Theorem 2. *Assume that $\overline{\{\tau(e) : e \in \mathcal{P}(M)\}} = \overline{\{\tau(e) : e \in \mathcal{P}(N)\}} = [0, 1]$. Let $M_1 \subset M_2 \subset M_3 \subset \dots$ and $N_1 \subset N_2 \subset N_3 \subset \dots$ be sequences of finite dimensional C^* -subalgebras of M and N such that $\bigcup_k M_k$ and $\bigcup_k N_k$ are strongly dense in M and N , respectively. It follows that*

$$\lambda(M, N) = \lim_{k \rightarrow \infty} \lim_{l \rightarrow \infty} \lambda(M_k, N_l).$$

For the proof we shall need the following lemma.

Lemma 3. *Let $A_1 \subset A_2 \subset A_3 \subset \dots$ be finite dimensional unital C^* -subalgebras of B and let A be the strong closure of $\bigcup_k A_k$. Assume that $\overline{\{\tau(e) : e \in \mathcal{P}(A)\}} = [0, 1]$.*

Let $p \in \mathcal{P}(A)$, $\epsilon > 0$. There are then a $k \in \mathbb{N}$ and a projection $q \in A_k$ such that $\|p - q\|_2 < \epsilon$ and $\tau(p) \leq \tau(q)$.

Proof. Let $0 < \delta$ such that $\delta + \sqrt{3\delta} < \epsilon$. It is well known that there are a $k \in \mathbb{N}$ and a projection $q_0 \in A_k$ such that $\|p - q_0\|_2 < \delta$. By increasing k we may assume that $\{\tau(e) : e \in \mathcal{P}(A_k)\}$ is δ -dense in $[0, 1]$. Let r_i , $i = 1, 2, \dots, m$, be the minimal nonzero central projections in A_k . We can choose a projection $q \in A_k$ such that $|\tau(q) - (\tau(p) + \delta)| \leq \delta$ and for each $i = 1, 2, \dots, m$, $qr_i \leq q_0 r_i$ or $qr_i \geq q_0 r_i$. Then $\tau(q) \geq \tau(p)$ and

$$\begin{aligned} \|q - p\|_2 &\leq \delta + \|q_0 - q\|_2 = \delta + \sqrt{|\tau(q_0 - q)|} \leq \\ &\delta + \sqrt{|\tau(q_0) - \tau(p)| + |\tau(p) - \tau(q)|} \leq \delta + \sqrt{3\delta} < \epsilon. \end{aligned}$$

□

Proof of Theorem 2. It is obvious that $\lambda(M_k, N_l)$ is increasing in l for fixed k and decreasing in k for fixed l . We first prove that

$$(2) \quad \lim_{l \rightarrow \infty} \lambda(M_k, N_l) = \lambda(M_k, N), \quad k \in \mathbb{N}.$$

Since $\lambda(M_k, N_l) \leq \lambda(M_k, N)$ for all l , we have trivially that $\lim_{l \rightarrow \infty} \lambda(M_k, N_l) \leq \lambda(M_k, N)$. Let $\epsilon > 0$. By definition of $\lambda(M_k, N)$ there is for all $e \in \mathcal{P}(M_k) \setminus \{0\}$ a projection $p \in N$ such that $\tau(e) \leq \tau(p)$ and $\tau(ep) > (\lambda(M_k, N) - \epsilon)\tau(p)$. By the density of $\bigcup_l N_l$ in N and Lemma 3 there are then an $i \in \mathbb{N}$ and a projection $p_i \in N_i$ such that $\tau(e) \leq \tau(p_i)$ and $\tau(ep_i) > (\lambda(M_k, N) - \epsilon)\tau(p_i)$. If $\delta < \min\{1, \epsilon\tau(p_i)\}$ and $e' \in \mathcal{P}(M_k)$ such that $\|e - e'\| < \delta$, then $e \sim e'$ and hence $\tau(e) = \tau(e')$ for all e' in this neighbourhood of e . Since

$$\tau(e'p_i) \geq \tau(ep_i) - \delta \geq (\lambda(M_k, N) - \epsilon)\tau(p_i) - \delta \geq (\lambda(M_k, N) - 2\epsilon)\tau(p_i),$$

we can use the compactness of the set of non-zero projections in M_k to find $j \in \mathbb{N}$ such that $\forall e \in \mathcal{P}(M_k) \setminus \{0\}$ there is a $p \in \mathcal{P}(N_j)$ such that $\tau(e) \leq \tau(p)$ and

$\tau(ep) \geq (\lambda(M_k, N) - 2\epsilon)\tau(p)$. It follows that $\lambda(M_k, N_j) \geq \lambda(M_k, N) - 2\epsilon$, proving (2). The next step is to prove that

$$(3) \quad \lim_{k \rightarrow \infty} \lambda(M_k, N) = \lambda(M, N) .$$

Since $\lambda(M_k, N) \geq \lambda(M, N)$ for all k we obviously have that $\lim_{k \rightarrow \infty} \lambda(M_k, N) \geq \lambda(M, N)$. Let $\epsilon > 0$; then there exists an $e \in \mathcal{P}(M) \setminus \{0\}$ such that $\tau(ep) < (\lambda(M, N) + \epsilon)\tau(p)$ for all $p \in \mathcal{P}(N)$ with $\tau(e) \leq \tau(p)$. By Lemma 3 there exist $k \in \mathbb{N}$ and a projection $e_k \in M_k$ such that $\|e - e_k\|_2 \leq \tau(e)\epsilon$ and $\tau(e) \leq \tau(e_k)$. Thus, if $p \in \mathcal{P}(N)$ and $\tau(p) \geq \tau(e_k)$, we have

$$\tau(e_k p) \leq \tau(ep) + \tau(e)\epsilon < (\lambda(M, N) + \epsilon)\tau(p) + \tau(e)\epsilon \leq (\lambda(M, N) + 2\epsilon)\tau(p),$$

proving that $\lambda(M_k, N) < \lambda(M, N) + 2\epsilon$. $\square$

Note that it does not give anything interesting if we exchange the order in which the two limits are taken in Theorem 2. Indeed, the proof of (3) gives that $\lim_{k \rightarrow \infty} \lambda(M_k, N_l) = \lambda(M, N_l)$, $l \in \mathbb{N}$. But $\lambda(M, N_l) = 0$ since M contains projections with arbitrarily small non-zero trace while $\inf\{\tau(e) : e \in \mathcal{P}(N_l) \setminus \{0\}\} > 0$.

For each $\delta \in [0, 1]$ set

$$\begin{aligned} \lambda_\delta(M, N) = \sup\{t \in [0, 1] : \forall e \in \mathcal{P}(M), \tau(e) \geq \delta \exists p \in \mathcal{P}(N) \\ \text{such that } \tau(e) \leq \tau(p) \text{ and } \tau(ep) \geq t\tau(p)\} . \end{aligned}$$

Clearly, $\lim_{\delta \rightarrow 0} \lambda_\delta(M, N) = \lambda_0(M, N) = \lambda(M, N)$. We will use $\lambda_\delta(M, N)$ to investigate to what extent $\lambda(M, N)$ is continuous in M, N . By [MT], Theorem 4, we cannot expect any global continuity of $\lambda(M, N)$. However, λ_δ is always continuous when $\delta > 0$. We define

$$d_0(M, N) = \sup_{e \in \mathcal{P}(M)} \inf_{p \in \mathcal{P}(N)} \|e - p\|_2 ,$$

and

$$D(M, N) = \max\{d_0(N, M), d_0(M, N)\} .$$

D is clearly a metric on the set of von Neumann subalgebras of B ; in fact $D(M, N)$ is the Hausdorff distance between the projection lattices in N and M . We say that a von Neumann subalgebra N of B is τ -continuous when

$$\{\tau(q) : q \in \mathcal{P}(N), q \leq p\} = [0, \tau(p)]$$

for all $p \in \mathcal{P}(N)$. By adopting the arguments from Lemma 3 we get the following

Lemma 4. *Let N_1, N_2 be τ -continuous von Neumann subalgebras of B . For any projection $e_1 \in N_1$ and $\epsilon > 0$ there is a projection $e_2 \in N_2$ such that $\tau(e_1) = \tau(e_2)$ and $\|e_1 - e_2\|_2 \leq D(N_1, N_2) + \sqrt{D(N_1, N_2)} + \epsilon$.*

It is straightforward to use Lemma 4 to prove

Lemma 5. *Let M_1, M_2, N_1, N_2 be τ -continuous von Neumann subalgebras of B . Then*

$$\begin{aligned} |\lambda_\delta(M_1, N_1) - \lambda_\delta(M_2, N_2)| \leq \\ \delta^{-\frac{1}{2}} \left(D(M_1, M_2) + \sqrt{D(M_1, M_2)} + D(N_1, N_2) + \sqrt{D(N_1, N_2)} \right) . \end{aligned}$$

Lemma 5 leads to the following improvement of Theorem 6 of [MT].

Theorem 6. *Assume that M_1, M_2, N_1, N_2 are II_1 -factors contained in a common ambient II_1 -factor such that $N_1 \subset M_1$ and $N_2 \subset M_2$. It follows that*

$$\frac{|[M_1 : N_1]^{-1} - [M_2 : N_2]^{-1}|}{\max\{[M_1 : N_1]^{\frac{1}{2}}, [M_2 : N_2]^{\frac{1}{2}}\}} \leq$$

$$D(M_1, M_2) + \sqrt{D(M_1, M_2)} + D(N_1, N_2) + \sqrt{D(N_1, N_2)} .$$

Proof. The inequality is trivial if $[M_1 : N_1]$ or $[M_2 : N_2]$ is ∞ , so we may assume that $\max\{[M_1 : N_1], [M_2 : N_2]\} < \infty$. By Theorem 1, $[M_i : N_i]^{-1} = \lambda(M_i, N_i)$, $i \in \{1, 2\}$. But there are projections (the Jones projections) $e_i \in M_i$ such that $E_{N_i}(e_i) = \lambda(M_i, N_i)1$, so we see that $\lambda(M_i, N_i) = \lambda_\delta(M_i, N_i)$, $i = 1, 2$, when $\delta = \min\{\lambda(M_1, N_1), \lambda(M_2, N_2)\}$. Apply Lemma 5. $\square$

REFERENCES

- [DT] S. Dorofeev, K. Thomsen, *Factors and subfactors arising from inductive limits of interval algebras*, Preprint, Aarhus, 1996.
- [J] V. Jones, *Index for subfactors*, Invent. Math. **72** (1983), 1 - 25. MR **84d**:46097
- [MT] B. Mashhood, K.F. Taylor, *On the continuity of the Index of Subfactors of a Finite Factor*, J. Functional Analysis **76** (1988), 56-66. MR **89h**:46090
- [PP] M. Pimsner, S. Popa, *Entropy and index for subfactors*, Ann. Sci. Ecole. Norm. Sup. **19** (1986), 57-106. MR **87m**:46120

MATEMATISK INSTITUT, NY MUNKEGADE, 8000 AARHUS C, DENMARK

E-mail address: dorofeev@mi.aau.dk

E-mail address: matkt@mi.aau.dk