HEREDITARY NOETHERIAN CATEGORIES WITH A TILTING COMPLEX

HELMUT LENZING

(Communicated by Eric M. Friedlander)

In memory of Maurice Auslander

ABSTRACT. We are characterizing the categories of coherent sheaves on a weighted projective line as the small hereditary noetherian categories without projectives and admitting a tilting complex. The paper is related to recent work with de la Peña (Math. Z., to appear) characterizing finite dimensional algebras with a sincere separating tubular family, and further gives a partial answer to a question of Happel, Reiten, Smalø (Mem. Amer. Math. Soc. **120** (1996), no. 575) regarding the characterization of hereditary categories with a tilting object.

A CHARACTERIZATION OF WEIGHTED PROJECTIVE LINES

We are going to characterize the categories $\text{coh}(\mathbb{X})$ of coherent sheaves on a weighted projective line $\mathbb{X}$ [3, 4].

Theorem 1. *Let k be an algebraically closed field. For a small connected abelian k -category $\mathcal{H}$ with finite dimensional morphism and extension spaces the following assertions are equivalent:*

- (i) *$\mathcal{H}$ is equivalent to the category of coherent sheaves on a weighted projective line.*
- (ii) *Each object of $\mathcal{H}$ is noetherian. $\mathcal{H}$ is hereditary, has no non-zero projectives, and admits a tilting complex.*
- (iii) *Each object of $\mathcal{H}$ is noetherian, moreover*
 - (a) *There exists an equivalence $\tau: \mathcal{H} \rightarrow \mathcal{H}$ (Auslander-Reiten translation) such that Serre duality $\text{D Ext}^1(A, B) \cong \text{Hom}(B, \tau A)$ holds functorially in $A, B \in \mathcal{H}$.*
 - (b) *The Grothendieck group $K_0(\mathcal{H})$ is finitely generated free, and the Euler form $\langle -, - \rangle: K_0(\mathcal{H}) \times K_0(\mathcal{H}) \rightarrow \mathbb{Z}$ given on classes of objects of $\mathcal{H}$ by $\langle [X], [Y] \rangle = \dim_k \text{Hom}(X, Y) - \dim_k \text{Ext}^1(X, Y)$ is non-degenerate of determinant ± 1 .*
 - (c) *$\mathcal{H}$ has an object without self-extensions which is not of finite length.*

Received by the editors February 9, 1995.

1991 *Mathematics Subject Classification.* Primary 14G14, 16G20; Secondary 18F20, 18E30.

Key words and phrases. Weighted projective line, coherent sheaf, almost-split sequence, Auslander-Reiten theory, finite dimensional algebra, quiver, derived category, exceptional object, tilting complex.

The result includes a characterization of the category $\text{coh}(\mathbb{P}_1(k))$ of coherent sheaves on the projective line adding the requirement: *If F is not of finite length and S is simple in $\mathcal{H}$, then $\text{Hom}(F, S) \neq 0$.* The theorem also gives a partial answer to a question raised by Happel, Reiten, Smalø [6] about the hereditary abelian k -categories admitting a tilting object.

For a finite dimensional k -algebra Σ , $\text{mod}(\Sigma)$ denotes the category of finite dimensional right modules. For a k -vectorspace X we denote its k -dual $\text{Hom}_k(X, k)$ by DX . For an abelian category $\mathcal{H}$ its derived category of bounded complexes is denoted $D^b(\mathcal{H})$, and $X \mapsto X[n]$ denotes the n th iterate of the translation functor for $D^b(\mathcal{H})$. An object X in $\mathcal{H}$ is called *noetherian* if each ascending chain of subobjects of X is stationary. We say that $E \in \mathcal{H}$ is *without self-extensions* if $\text{Ext}^n(E, E) = 0$ holds for each $n \neq 0$. A k -category $\mathcal{H}$ is *connected* if a representation of $\mathcal{H}$ as a coproduct $\mathcal{H} \cong \mathcal{A} \amalg \mathcal{B}$ implies $\mathcal{A} = 0$ or $\mathcal{B} = 0$. We say that $\mathcal{H}$ is *hereditary* if $\text{Ext}_{\mathcal{H}}^2(-, -) = 0$.

A *tilting object* Σ in $\mathcal{H}$ is defined by the fact that the right derived functor of $\text{Hom}(\Sigma, -): \mathcal{H} \rightarrow \text{mod}(\text{End}(\Sigma))$ induces an equivalence of triangulated categories $D^b(\mathcal{H}) \rightarrow D^b(\text{mod}(\text{End}(\Sigma)))$. More generally, we say that $\mathcal{H}$ admits a *tilting complex* if for some finite dimensional algebra Σ there exists an equivalence $\Phi: D^b(\text{mod}(\Sigma)) \rightarrow D^b(\mathcal{H})$ of triangulated categories. Here, Σ has an interpretation as the endomorphism ring of the tilting complex $\Phi(\Sigma)$ of $D^b(\mathcal{H})$. Whenever convenient, we prefer to view a tilting complex as a full subcategory of $D^b(\mathcal{H})$ with finitely many objects, thus avoiding the use of endomorphism rings.

This work has substantially profited from the collaboration with José Antonio de la Peña on finite dimensional algebras with a sincere separating tubular family [9]; we further thank Hagen Meltzer for critical comments on a first draft of this paper.

PROOF OF THE CHARACTERIZATION

Let $\mathcal{H}_0$ be the full subcategory consisting of all objects of $\mathcal{H}$ having finite length.

Lemma 1. *Assume that $\mathcal{H}$ satisfies condition (iii)(a). Then $\mathcal{H}_0$ is uniserial, i.e. each object $U \in \mathcal{H}_0$ has a unique finite composition series. Accordingly, for some index set $\mathbb{X}$ the category $\mathcal{H}_0$ decomposes into a coproduct $\mathcal{H}_0 = \coprod_{x \in \mathbb{X}} \mathcal{U}_x$ of connected uniserial categories $\mathcal{U}_x$, whose indecomposables form stable tubes.*

If moreover $K_0(\mathcal{H})$ is finitely generated, then for each $x \in \mathbb{X}$ the simple objects in $\mathcal{U}_x$ form an Auslander-Reiten orbit of finite cardinality $u(x)$. Moreover $u(x) = 1$ for all but finitely many $x \in \mathbb{X}$.

Proof. In view of (iii)(a) the category $\mathcal{H}$ is hereditary. Hence $\mathcal{H}_0$ is a hereditary abelian category with almost-split sequences where each object has finite length, and the Auslander-Reiten translation $\tau: \mathcal{H}_0 \rightarrow \mathcal{H}_0$ is an equivalence. For simple objects S_i, S_j this implies that $\text{Ext}^1(S_j, S_i) \neq 0$ if and only if $S_i \cong \tau S_j$, moreover that in this case the Ext-space is one-dimensional. Uniseriality of $\mathcal{H}_0$ now follows easily [2].

For each $x \in \mathbb{X}$ we fix a simple object S_x from $\mathcal{U}_x$, and collect a representative system (S_α) , $\alpha \in I$, of the remaining simple objects. I admits a linear ordering such that $\langle [S_\alpha], [S_\beta] \rangle = 1$ for $\alpha = \beta$, but is zero for $\alpha > \beta$. This implies that the classes $[S_\alpha]$, $\alpha \in I$, are linearly independent in $K_0(\mathcal{H})$, and proves the claim on u . $\square$

Proof of (ii) $\Rightarrow$ (iii). In view of $D^b(\mathcal{H}) \cong D^b(\text{mod}(\Sigma))$ the group $K_0(\mathcal{H})$ is isomorphic to $K_0(\text{mod}(\Sigma))$, which is finitely generated free. As a tilting complex for a hereditary category, the algebra Σ has finite global dimension which implies the assertion on the Euler form. Happel's theorem [5] shows that $D^b(\mathcal{H}) = D^b(\text{mod}(\Sigma))$ has Auslander-Reiten triangles, and that further the Auslander-Reiten translation τ is an equivalence for $D^b(\mathcal{H})$. Next we prove that $\tau(\mathcal{H})$ is contained in $\mathcal{H}$. Let X be indecomposable in $\mathcal{H}$, we get $\tau X = Y[n]$ for some $Y \in \mathcal{H}$ and $n \in \mathbb{Z}$. Because

$$\text{DExt}_{\mathcal{H}}^{n+1}(X, Y) \cong \text{Hom}_{D^b(\mathcal{H})}(Y[n+1], \tau X[1]) \neq 0,$$

$n \geq 1$ is impossible for a hereditary category. Assuming $n \leq -1$ implies

$$\text{DExt}_{\mathcal{H}}^1(X, Z) \cong \text{Hom}_{D^b(\mathcal{H})}(Z, Y[n]) = 0$$

for each $Z \in \mathcal{H}$, so leads to a non-zero projective X in $\mathcal{H}$. This proves the assertion on Serre duality for $\mathcal{H}$, which in turn implies the existence of almost-split sequences in $\mathcal{H}$ (cf. [3]).

Finally assume that condition (iii)(c) is violated. Then $D^b(\mathcal{H})$ has a tilting complex lying in $D^b(\mathcal{H}_0)$, which involving Lemma 1 implies that $\mathcal{H} = \mathcal{H}_0$ is connected uniserial, hence the existence of a simple object S of τ -period p such that the classes $[S], [\tau S], \dots, [\tau^{p-1}S]$ form a $\mathbb{Z}$ -basis of $K_0(\mathcal{H})$. Then $\mathbf{a} = \sum_{j=0}^{p-1} [\tau^j S]$ is a non-zero element of $K_0(\mathcal{H})$ with $\langle \mathbf{a}, - \rangle = 0$, contradiction. $\square$

The proof of implication (iii) $\Rightarrow$ (i) will occupy the rest of the paper. Each object X from $\mathcal{H}$ has a largest subobject tX of finite length (its *torsion part*), and X/tX has no simple submodules, i.e. is *torsionfree*. The full subcategory of torsionfree objects is denoted $\mathcal{H}_+$.

Lemma 2. *An indecomposable object X of $\mathcal{H}$ is either torsion ($X \in \mathcal{H}_0$) or torsionfree ($X \in \mathcal{H}_+$).*

Proof. Since $\text{DExt}^1(X/tX, tX) = \text{Hom}(\tau^-(tX), X/tX) = 0$ the sequence $0 \rightarrow tX \rightarrow X \rightarrow X/tX \rightarrow 0$ splits. $\square$

Lemma 3. *There is a surjective linear mapping $\text{rk}: K_0(\mathcal{H}) \rightarrow \mathbb{Z}$, $[X] \mapsto \text{rk } X$, called **rank function**, such that $\text{rk } \tau X = \text{rk } X$ for each $X \in \mathcal{H}$, rk vanishes on $\mathcal{H}_0$, and $\text{rk } F > 0$ for each $0 \neq F \in \mathcal{H}_+$.*

Proof (de la Peña). For $x \in \mathbb{X}$ and $S_x \in \mathcal{U}_x$ simple, we define $w_x = \sum_{j=1}^{\mathcal{U}(x)} [\tau^j S_x]$ in $K_0(\mathcal{H})$. Note that each w_x is stable under the mapping induced by τ on $K_0(\mathcal{H})$. We choose $x_1, \dots, x_s$ from $\mathbb{X}$ such that the subgroup of $K_0(\mathcal{H})$ generated by $w_{x_1}, \dots, w_{x_s}$ contains all w_x ($x \in \mathbb{X}$) and put $w = w_{x_1} + \dots + w_{x_s}$.

By noetherianness each non-zero $F \in \mathcal{H}_+$ has a simple quotient S_x , hence $\langle [F], w_x \rangle > 0$, which implies $\langle [F], w_{x_i} \rangle > 0$ for some $i = 1, \dots, s$, and finally $\langle [F], w \rangle > 0$. If ε denotes the index of $\langle K_0(\mathcal{H}), w \rangle$ in $\mathbb{Z}$ we may take $\text{rk } \mathbf{a} = 1/\varepsilon \cdot \langle \mathbf{a}, w \rangle$. $\square$

Lemma 4. *Assume $B \in \mathcal{H}_+$ and $\text{Ext}^1(\mathcal{U}_x, B) \neq 0$. For each $n \geq 0$ there exists an exact sequence $0 \rightarrow B \rightarrow B_n \rightarrow U_n \rightarrow 0$ with $B_n \in \mathcal{H}_+$, $U_n \in \mathcal{U}_x$ and $[U_n] = n \cdot w_x$.*

Proof. Let S_x be simple in $\mathcal{U}_x$ with $\text{Ext}^1(S_x, B) \neq 0$. Choose non-split exact sequences $0 \rightarrow B'_i \rightarrow B'_{i+1} \rightarrow \tau^{-i} S_x \rightarrow 0$ ($i = 0, 1, 2, \dots$), where $B'_0 = B$, and put $B_n = B'_{nu(x)}$. $\square$

Lemma 5. *For each non-zero F in $\mathcal{H}_+$ and each $x \in \mathbb{X}$ there is a non-zero homomorphism from F into an object from $\mathcal{U}_x$. Accordingly, $[X] = 0$ in $K_0(\mathcal{H})$ implies $X = 0$ in $\mathcal{H}$.*

Proof. We put

$$\begin{aligned}\mathbb{Y} &= \{y \in \mathbb{X} \mid \text{Hom}(F, \mathcal{U}_y) = 0\}, \\ \mathcal{A} &= \{A \in \mathcal{H} \mid \text{Hom}(A, \mathcal{U}_y) = 0 \text{ for all } y \in \mathbb{Y}\}, \\ \mathcal{B} &= \{B \in \mathcal{H} \mid \text{Hom}(\mathcal{A}, B) = 0\}.\end{aligned}$$

Step 1. $\text{Hom}(\mathcal{B}, \mathcal{U}_x) = 0$ for all $x \in \mathbb{X} \setminus \mathbb{Y}$.

Since $\tau \mathcal{U}_x = \mathcal{U}_x$ it is equivalent to prove that $\text{Ext}^1(\mathcal{U}_x, \mathcal{B}) = 0$. Otherwise we obtain some B in $\mathcal{B}$ and exact sequences

$$0 \rightarrow B \rightarrow B_n \rightarrow U_n \rightarrow 0 \quad (n = 1, 2, \dots),$$

having the properties of Lemma 4. We show that each B_n actually belongs to $\mathcal{B}$. Consider a morphism $f: A \rightarrow B_n$ with $A \in \mathcal{A}$, and from the pull-back

$$\begin{array}{ccccccc} 0 & \longrightarrow & A' & \longrightarrow & A & \longrightarrow & U_n \longrightarrow 0 \\ & & \downarrow f' & & \downarrow f & & \parallel \\ 0 & \longrightarrow & B & \longrightarrow & B_n & \longrightarrow & U_n \longrightarrow 0. \end{array}$$

Using $x \notin \mathbb{Y}$ we see that $A' \in \mathcal{A}$, hence $f' = 0$, so f induces a morphism $U_n \rightarrow B_n$ which is zero, implying $\text{Hom}(\mathcal{A}, B_n) = 0$. Since $[B_n] = [B] + nw_x$, we obtain $\langle [F], B_n \rangle > 0$ for large n , thus contradicting $\text{Hom}(\mathcal{A}, \mathcal{B}) = 0$.

Step 2. Each X in $\mathcal{H}$ has the form $X \cong A \oplus B$ with $A \in \mathcal{A}, B \in \mathcal{B}$.

By noetherianness X has a maximal subobject A from $\mathcal{A}$, accordingly X/A has no non-zero subobject from $\mathcal{A}$, hence belongs to $\mathcal{B}$. Obviously, $\mathcal{A}$ is closed under τ^- , therefore $\text{DExt}^1(X/A, A) = \text{Hom}(\tau^- A, X/A) = 0$, and $X \cong A \oplus X/A$.

Step 3. $\text{Hom}(\mathcal{B}, \mathcal{A}) = 0$.

Otherwise we find $A \in \mathcal{A} \cap \mathcal{H}_+$, $B \in \mathcal{B} \cap \mathcal{H}_+$ and a non-zero morphism $f: B \rightarrow A$. Let $A' = \text{im}(f)$. We choose $A' \subseteq A'' \subseteq A$ such that $A''/A' = t(A/A')$. Since $A \in \mathcal{A}$ and $A/A'' \in \mathcal{H}_+$ we see that $A'' \in \mathcal{A}$. Replacing A by A'' we may thus assume that $A/A' \in \mathcal{H}_0$, in fact $A/A' \in \coprod_{x \in \mathbb{X} \setminus \mathbb{Y}} \mathcal{U}_x$. By Step 1 the sequence $0 \rightarrow A' \rightarrow A \rightarrow A/A' \rightarrow 0$ splits, so A' belongs to $\mathcal{A}$. We thus arrive at an epimorphism $B \twoheadrightarrow A' \twoheadrightarrow S_x$ where S_x is simple in $\mathcal{U}_x$ and $x \in \mathbb{X} \setminus \mathbb{Y}$, contradiction, which proves the assertion of Step 3.

Combining Steps 2 and 3 with $\text{Hom}(\mathcal{A}, \mathcal{B}) = 0$ we see that $\mathcal{H} = \mathcal{A} \coprod \mathcal{B}$. Since $\mathcal{H}$ is connected and $0 \neq F \in \mathcal{A}$, it follows that $\mathcal{B} = 0$, hence $\mathbb{Y} = \emptyset$ as claimed. $\square$

By $K'_0(\mathcal{H})$ we denote the image of the natural map $K_0(\mathcal{H}_0) \rightarrow K_0(\mathcal{H})$, i.e. the subgroup of $K_0(\mathcal{H})$ generated by the classes of simple objects.

Lemma 6. *There exists a torsionfree object L of rank one. $K_0(\mathcal{H}) = \mathbb{Z}[L] \oplus K'_0(\mathcal{H})$. Moreover for any $x, y \in \mathbb{X}$, w_x and w_y are proportional.*

Proof. Choose a torsionfree object $L \neq 0$ of minimal rank. Each $\mathbf{a} \in K'_0(\mathcal{H})$ satisfies $\text{rk } \mathbf{a} = 0$, on the other hand $\text{rk } L > 0$, which implies $\mathbb{Z}[L] \cap K'_0(\mathcal{H}) = 0$. We show by induction on $r = \text{rk } A$ that the class $[A]$ of an indecomposable object A belongs

to $\mathbb{Z}[L] + K'_0(\mathcal{H})$. For $r = 0$ this follows from Lemma 3. For $r > 0$, A is torsionfree. We choose $x \in \mathbb{X}$ and a simple object S_x from $\mathcal{U}_x$ with $\text{Ext}^1(S_x, A) \neq 0$. Invoking Lemma 4 we get an exact sequence $0 \rightarrow A \rightarrow \overline{A} \rightarrow U \rightarrow 0$, where $\overline{A}$ is torsionfree, U is in $\mathcal{U}_x$, and a non-zero homomorphism $f: L \rightarrow \overline{A}$. Because L has minimal rank and $\overline{A}$ is torsionfree, f is a monomorphism hence $C = \text{coker}(f)$ has strictly smaller rank than A . By induction $[C]$, hence $[A]$, belongs to $\mathbb{Z}[L] \oplus K'_0(\mathcal{H})$, proving the splitting of $K_0(\mathcal{H})$. Surjectivity of the rank function now implies that L has rank one.

Since the linear forms $\langle -, w_x \rangle$ and $\langle -, w_y \rangle$ vanish on $K'_0(\mathcal{H})$ they are determined by their non-zero value on $[L]$, hence are proportional. Because the Euler form on $K_0(\mathcal{H})$ is non-degenerate, the corresponding assertion holds for w_x and w_y . $\square$

Through the formation of perpendicular categories [4] we are going to reduce the weight function $u: \mathbb{X} \rightarrow \mathbb{N}$. An object E from $\mathcal{H}$ is called *exceptional* if E has a trivial endomorphism ring k and no self-extensions. Invoking Lemma 5 it follows from [7] that each indecomposable object $E \in \mathcal{H}$ without self-extensions is exceptional. Notice that every simple object S in $\mathcal{U}_x$ with $u(x) > 1$ is exceptional.

Proposition 1. *Let $x \in \mathbb{X}$ be exceptional, i.e. $u(x) > 1$, and let S be simple in $\mathcal{U}_x$.*

(a) *The right perpendicular category $\mathcal{H}' = S^\perp$ of S in $\mathcal{H}$, consisting of all $X \in \mathcal{H}$ with $\text{Hom}(S, X) = 0 = \text{Ext}^1(S, X)$, satisfies conditions (iii) (a), (b), (c) again.*

(b) *$\mathbb{X}$ serves naturally also as a parametrizing set for the decomposition of $\mathcal{H}'_0 = S^\perp \cap \mathcal{H}_0$ into connected uniserial categories $\mathcal{U}'_y = S^\perp \cap \mathcal{U}_y$. The number of isomorphism classes of simples from $\mathcal{U}'_y$ is given by $u'(x) = u(x) - 1$ and $u'(y) = u(y)$ for $y \neq x$. Moreover, inclusion $\mathcal{H}' \hookrightarrow \mathcal{H}$ preserves the rank.*

(c) $K_0(\mathcal{H}) = \mathbb{Z}[S] \oplus K_0(\mathcal{H}')$.

Proof. Since $\mathcal{H}$ is hereditary and S is exceptional, inclusion $\mathcal{H}' \hookrightarrow \mathcal{H}$ admits a left adjoint ℓ and a right adjoint r [4]. Since S is simple, the category $\mathcal{H}'_0$ of torsion objects in $\mathcal{H}'$ agrees with $\mathcal{H}' \cap \mathcal{H}_0$, and therefore $\mathcal{H}'_0 = \coprod_{y \in \mathbb{X}} \mathcal{U}'_y$ with $\mathcal{U}'_y = \mathcal{U}_y$ if $y \neq x$; further $\mathcal{U}'_x = S^\perp \cap \mathcal{U}_x$ is connected as well with $u(x) - 1$ isomorphic classes of simple objects. Moreover, $w'_y = w_y$ for each $y \in \mathbb{X}$, so inclusion $\mathcal{H}' \hookrightarrow \mathcal{H}$ preserves the rank.

If E is torsionfree in $\mathcal{H}$ without self-extensions, then $\text{Hom}(S, E) = 0$ and ℓE is given by the middle term of the S -universal extension

$$0 \rightarrow E \rightarrow \ell E \rightarrow \text{Ext}^1(S, E) \otimes_k S \rightarrow 0,$$

in particular ℓE is non-zero torsionfree. From the exactness of

$$0 = \text{Ext}^1(E, E) \rightarrow \text{Ext}^1(E, \ell E) \rightarrow \text{Ext}^1(S, E) \otimes_k \text{Ext}^1(E, S) = 0$$

we get $\text{Ext}^1(E, \ell E) = 0$. By definition $\text{Ext}^1(S, \ell E) = 0$ holds, hence $\text{Ext}^1(\ell E, \ell E) = 0$, which proves the existence of a torsionfree object in $\mathcal{H}'$ without self-extensions.

Finally, assume that A and B are in $\mathcal{H}'$. Since

$$\text{D Ext}^1(A, B) = \text{Hom}(\ell \tau_{\mathcal{H}}^- B, A) = \text{Hom}(B, r \tau_{\mathcal{H}} A),$$

projectivity of A (resp. injectivity of B) in $\mathcal{H}'$ implies $r \tau_{\mathcal{H}} A = 0$ (resp. $\ell \tau_{\mathcal{H}}^- B = 0$) which, by the construction of r and ℓ [4], implies $A \in \mathcal{H}'_0$ (resp. $B \in \mathcal{H}'_0$). But any projective (resp. injective) in $\mathcal{U}'_y$, $y \in \mathbb{X}$, is zero, hence $A = 0$ (resp. $B = 0$). Invoking Auslander-Reiten theory this implies that $\tau_{\mathcal{H}'} = r \tau_{\mathcal{H}}$ and $\tau_{\mathcal{H}'}^- = \ell \tau_{\mathcal{H}}^-$ are inverse equivalences. For assertion (c) we refer to [4]. Further $\langle [S], K_0(\mathcal{H}') \rangle = 0$ implies that the Euler forms on $K_0(\mathcal{H})$ and $K_0(\mathcal{H}')$ do have the same determinant. $\square$

Repeating the process of Proposition 1 $\sum_{x \in \mathbb{X}} (u(x) - 1)$ times leads to an exact abelian subcategory $\mathcal{H}'$ of $\mathcal{H}$ satisfying (iii) (a), (b), (c) with $u'(x) = 1$ for each $x \in \mathbb{X}$.

Proposition 2. *Each torsionfree rank one object L of $\mathcal{H}'$ is exceptional. For each simple object S_x , $x \in \mathbb{X}$, we have $\dim_k \operatorname{Hom}(L, S_x) = 1$ and $[S_x] = w$.*

Moreover there exist torsionfree rank one objects L_1, L_2 such that

$$(1) \quad \operatorname{Hom}(L_2, L_1) = 0, \quad \dim_k \operatorname{Hom}(L_1, L_2) = 2, \quad \operatorname{Ext}^1(L_i, L_j) = 0 \quad \text{for } i, j = 1, 2.$$

Proof. By Lemma 6 the group $K_0(\mathcal{H}')$ is free abelian of rank two. Fix a torsionfree object L of rank one, note that $\operatorname{End}(L) = k$ hence $\langle [L], [L] \rangle = 1 - g$, where $g = \dim_k \operatorname{Ext}^1(L, L)$, and define the *degree function* $\deg: K_0(\mathcal{H}') \rightarrow \mathbb{Z}$ by

$$\deg z = \langle [L], z \rangle - (1 - g) \operatorname{rk} z.$$

We thus obtain a Riemann-Roch formula

$$\langle \mathbf{a}, \mathbf{b} \rangle = (1 - g) \operatorname{rk} \mathbf{a} \operatorname{rk} \mathbf{b} + \begin{vmatrix} \operatorname{rk} \mathbf{a} & \operatorname{rk} \mathbf{b} \\ \deg \mathbf{a} & \deg \mathbf{b} \end{vmatrix} \quad \text{for all } \mathbf{a}, \mathbf{b} \in K_0(\mathcal{H}').$$

The existence of an exceptional object E of $\mathcal{H}'$ implies $0 < \langle [E], [E] \rangle = (1 - g)(\operatorname{rk} E)^2$ hence $g = 0$, and so L is exceptional.

Next we choose $x \in \mathbb{X}$ such that $d = \deg w'_x$ is minimal. Note that $d > 0$. For $y \in \mathbb{X}$ write $\deg w'_y = d \cdot q + r$ with $q, r \in \mathbb{N}$ and $0 \leq r < d$. Extending L by S_y and q times by S_x we obtain torsionfree rank one objects L', L'' with

$$[L'] = [L] + w'_y, \quad [L''] = [L] + q \cdot w'_x,$$

hence

$$\langle [L''], [L'] \rangle = \langle [L], [L] \rangle + \langle [L], w'_y \rangle - q \cdot \langle [L], w_x \rangle \geq 1.$$

We thus arrive at an exact sequence $0 \rightarrow L'' \rightarrow L' \rightarrow U \rightarrow 0$, where $U \in \mathcal{H}'_0$ has degree $< d$. This shows $U = 0$, hence $w'_y = q \cdot w'_x$. In view of Lemma 6 the classes $[L], [S_x]$ thus form a basis of $K_0(\mathcal{H}')$, accordingly the determinant of the Euler form is $\pm 1 = \langle [L], [S_x] \rangle^2$, proving $\dim \operatorname{Hom}(L, S_x) = 1$, therefore $\deg S_x = 1$.

Put $L_1 = L$, and let L_2 be the middle term of a non-split exact sequence $\mu: 0 \rightarrow L_1 \xrightarrow{\xi} L_2 \rightarrow S_x \rightarrow 0$. Application of $\operatorname{Hom}(L_1, -)$ to μ shows $\operatorname{Hom}(L_1, L_2) \cong k^2$ and $\operatorname{Ext}^1(L_1, L_2) = 0$, while application of $\operatorname{Hom}(-, L_1)$ leads to $\operatorname{Hom}(L_2, L_1) = 0$ and $\operatorname{Ext}^1(L_2, L_1) = 0$.

Extend ξ to a k -basis ξ, η of $\operatorname{Hom}(L_1, L_2)$. Invoking the properties of rank and degree, for each $a \in k$ the cokernel T_a of $\eta - a\xi: L_1 \rightarrow L_2$ is seen to be simple. In fact, each simple object S_y ($y \neq x$) is isomorphic to some T_a : Setting $\xi^* = \operatorname{Hom}(\xi, S_y)$, $\eta^* = \operatorname{Hom}(\eta, S_y)$ we see that ξ^* is an isomorphism because $x \neq y$; moreover, since k is algebraically closed there exists an eigenvalue $a \in k$ such that the kernel $\operatorname{Hom}(T_a, S_y)$ of $\eta^* - a\xi^*: \operatorname{Hom}(L_2, S_y) \rightarrow \operatorname{Hom}(L_1, S_y)$ is non-zero. Hence $S_y \cong T_a$, in particular $[S_y] = [L_1] - [L_2] = [S_x]$. $\square$

Proposition 3. $\mathcal{H}$ is a tilting object Δ

$$\begin{array}{ccccc}
 & S_1^{[p_1-1]} & \xrightarrow{\quad} \cdots \xrightarrow{\quad} & S_1^{[1]} \\
 \eta_1 & & & \\
 \\
 L_1 \xrightleftharpoons[\xi_2]{\xi_1} L_2 & \xrightarrow{\eta_2} & S_2^{[p_2-1]} & \xrightarrow{\quad} \cdots \xrightarrow{\quad} & S_2^{[1]} \\
 & & & & \\
 & & \vdots & & \vdots \\
 \eta_t & & S_t^{[p_t-1]} & \xrightarrow{\quad} \cdots \xrightarrow{\quad} & S_t^{[1]}
 \end{array}$$

where L_1, L_2 are torsionfree rank one objects satisfying (1), and $S_i^{[l]}$ is an exceptional torsion object of length l .

Δ is a squid algebra in the sense of [10], i.e. equivalent to the path category of the above quiver subject to the relations

$$(2) \quad \eta_1 \circ \xi_1 = 0, \quad \eta_2 \circ \xi_2 = 0, \quad \eta_i \circ (\xi_2 - \lambda_i \xi_1) = 0, \quad i = 3, \dots, t,$$

where $\lambda_3, \dots, \lambda_t$ are pairwise distinct non-zero elements from k .

Δ also has a realization as a tilting object on the weighted projective line $\mathbb{X}(\mathbf{p}, \boldsymbol{\lambda})$ given by the data $\mathbf{p} = (p_1, \dots, p_t)$, $\boldsymbol{\lambda} = (\lambda_3, \dots, \lambda_t)$.

Proof. Let $x_1, \dots, x_t$ denote the sequence of pairwise distinct exceptional elements from $\mathbb{X}$. We put $p_i = u(x_i)$, $\mathcal{U}_i = \mathcal{U}_{x_i}$. Let S_i be simple in $\mathcal{U}_i$ and $\mathcal{H}'$ denote the right perpendicular category with respect to the system

$$(3) \quad \tau^l S_i, \quad l = 0, \dots, p_i - 2, \quad i = 1, \dots, t,$$

and take a system L_1, L_2 of torsionfree rank one objects from $\mathcal{H}'$ as established in Proposition 2.

By induction on the rank it is easily verified (compare proof of Lemma 6) that each object from $\mathcal{H}$ is contained in the smallest subcategory of $\mathcal{H}$ which is closed under extensions, kernels of epimorphisms, cokernels of monomorphisms and contains

$$(4) \quad L_1, L_2, S_i, \tau S_i, \dots, \tau^{p_i-2} S_i, \quad i = 1, \dots, t;$$

in particular (4) generates $D^b(\mathcal{H})$.

Let $S_i^{[n]}$ denote the indecomposable object of $\mathcal{U}_i$ with top S_i and length n . Then the system

$$(5) \quad L_1, L_2, S_i = S_i^{[1]}, S_i^{[2]}, \dots, S_i^{[p_i-1]}, \quad i = 1, \dots, t,$$

generates $D^b(\mathcal{H})$ as well, moreover satisfies

$$(6) \quad \text{Ext}^1(S_i^{[a]}, S_{i'}^{[a']}) = 0, \quad \text{Ext}^1(S_i^{[a]}, L_j) = 0, \quad \text{Ext}^1(L_j, S_i^{[a]}) = 0$$

for all $i, i' = 1, \dots, t$, $a = 1, \dots, p_i - 1$, $a' = 1, \dots, p_{i'} - 1$, $j = 1, 2$.

Invoking $w_x = w$ for each $x \in \mathbb{X}$, we obtain exact sequences

$$0 \rightarrow L_1 \xrightarrow{\xi_i} L_2 \xrightarrow{\bar{\eta}_i} S_i^{[p_i]} \rightarrow 0 \quad (1 \leq i \leq t),$$

defining t mutually disjoint 1-dimensional subspaces $k\xi_i$ of $\text{Hom}(L_1, L_2)$, thus $\xi_i = \xi_2 - \lambda_i \xi_1$ for $2 \leq i \leq t$, where $0 = \lambda_2, \lambda_3, \dots, \lambda_t$ are pairwise distinct. Setting $\eta_i = [L_2 \xrightarrow{\bar{\eta}_i} S_i^{[p_i]} \rightarrow S_i^{[p_i-1]}]$ and calculating dimensions of Hom-spaces by means of the Euler form further establishes that (2) is a complete set of relations for the subcategory Δ .

It is known (cf. [8]) and not difficult to prove that Δ can be realized as a tilting sheaf on the weighted projective line $\mathbb{X}(\mathbf{p}, \boldsymbol{\lambda})$ arising from the above data $\mathbf{p} = (p_1, \dots, p_t)$ and $\boldsymbol{\lambda} = (\lambda_3, \dots, \lambda_t)$ [3]. Since moreover Δ has finite global dimension (actually $\text{gl. dim } \Delta \leq 2$) we may invoke Beilinson's lemma [1] to conclude that Δ is also a tilting object in $\mathcal{H}$. $\square$

Proof of (iii) $\Rightarrow$ (i). Abbreviating $\mathbb{X}(\mathbf{p}, \boldsymbol{\lambda})$ to $\mathbb{X}$ from now on, the right derived functor of $\text{Hom}_{\mathbb{X}}(\Delta, -): \text{coh}(\mathbb{X}) \rightarrow \text{mod}(\Delta)$ induces an equivalence of triangulated categories $\alpha: D^b(\text{coh}(\mathbb{X})) \rightarrow D^b(\text{mod}(\Delta))$, similarly the right derived functor of $\text{Hom}_{\mathcal{H}}(\Delta, -)$ induces an equivalence $\beta: D^b(\mathcal{H}) \rightarrow D^b(\text{mod}(\Delta))$. Since α and β preserve Δ the composition $\phi = \alpha^{-1} \circ \beta: D^b(\mathcal{H}) \rightarrow D^b(\text{coh}(\mathbb{X}))$ preserves the rank and maps $\mathcal{H}_0$ onto the category $\text{coh}_0(\mathbb{X})$ of finite length sheaves.

An indecomposable object X from $D^b(\mathcal{H})$ lies in $\mathcal{H}$ if and only if X is either in $\mathcal{H}_0$ or has rank > 0 and satisfies $\text{Hom}(X, \mathcal{H}_0) \neq 0$. Similarly an indecomposable object Y from $D^b(\text{coh}(\mathbb{X}))$ belongs to $\text{coh}(\mathbb{X})$ if and only if either Y is in $\text{coh}_0(\mathbb{X})$ or Y has rank > 0 and satisfies $\text{Hom}(Y, \text{coh}_0(\mathbb{X})) \neq 0$. Because $\phi(\mathcal{H}_0) = \text{coh}_0(\mathbb{X})$, and ϕ preserves the rank, it hence follows that ϕ induces an equivalence $\mathcal{H} \rightarrow \text{coh}(\mathbb{X})$, thus concluding the proof of the theorem. $\square$

Given $\mathcal{H}$ satisfying (ii) or (iii), the theorem implies the existence of a “natural” bijection between the set $\mathbb{X}$, parametrizing the decomposition of $\mathcal{H}_0$ into connected components, and the projective line $\mathbb{P}_1(k)$, the point set underlying $\mathbb{X}$. To determine the weighted projective line corresponding to $\mathcal{H}$, one fixes a simple object S with $\tau S \cong S$ further a rank one object L of $\mathcal{H}$, and forms a non-split extension $0 \rightarrow L \rightarrow \bar{L} \rightarrow S \rightarrow 0$. The k -space $\text{Hom}(L, \bar{L})$ has dimension two; each non-zero $v: L \rightarrow \bar{L}$ is a monomorphism whose cokernel is indecomposable of finite length, therefore belongs to $\mathcal{U}_{\psi(v)}$ for a unique $\psi(v) \in \mathbb{X}$. The induced mapping $\mathbb{P}_1(k) = \mathbb{P}(\text{Hom}(L, \bar{L})) \rightarrow \mathbb{X}$, $kv \mapsto \psi(v)$, is a bijection turning the function u from Lemma 1 into a weight function on $\mathbb{P}_1(k)$, determining thus completely the data for the weighted projective line.

REFERENCES

1. A. A. Beilinson, *Coherent sheaves on P_n and problems of linear algebra*, Funct. Anal. Appl. **12** (1979), 214–216. MR **80c**:14010b
2. P. Gabriel, *Indecomposable representations II*, Symposia Mat. Inst. Naz. Alta Mat. **11** (1973), 81–104. MR **49**:5132
3. W. Geigle and H. Lenzing, *A class of weighted projective curves arising in representation theory of finite dimensional algebras*. In *Singularities, representations of algebras, and vector bundles*. Lect. Notes Math. 1273 (1987), 265–297, Springer-Verlag, Berlin-Heidelberg-New York. MR **89b**:14049
4. ———, *Perpendicular categories with applications to representations and sheaves*, J. Algebra **144** (1991), 273–343. MR **93b**:16011

5. D. Happel, *Triangulated categories in the representation theory of finite dimensional algebras*, London Math. Soc. Lect. Notes **119**, Cambridge University Press, 1988. MR **89e**:16035
6. D. Happel, I. Reiten and S. Smalø, *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. **120** (1996), no. 575. CMP 96:08
7. D. Happel and C. M. Ringel, *Tilted Algebras*, Trans. Amer. Math. Soc. **274** (1982), 399–443. MR **84d**:16027
8. H. Lenzing and H. Meltzer, *Tilting sheaves and concealed-canonical algebras*, CMS Conf. Proc. **18** (1996), 455–473. CMP 96:12
9. H. Lenzing and J. A. de la Peña, *Algebras with a separating tubular family*, Math. Z., to appear.
10. C. M. Ringel, *The canonical algebras* with an appendix by William Crawley-Boevey. In *Topics in Algebra*, Banach Center Publ. **26** (1990), 407–432. MR **93e**:16022

UNIVERSITÄT-GH PADERBORN, FACHBEREICH MATHEMATIK-INFORMATIK, D-33095 PADERBORN, GERMANY

E-mail address: `helmut@uni-paderborn.de`