A REMARKABLE REARRANGEMENT OF THE HAAR SYSTEM IN L_p

PAUL F. X. MÜLLER AND GIDEON SCHECHTMAN

(Communicated by Dale Alspach)

ABSTRACT. We introduce a non-standard but, to our opinion natural, order on the initial segments of the Haar system and investigate the isomorphic classification of the linear span, in L_p , of block bases, with respect to this order.

0. INTRODUCTION

In [DS] it was proved that every unconditional basic sequence $\{x_i\}_{i=1}^\infty$ in L_p , $2 < p < \infty$, which is not equivalent to the natural basis of ℓ_p has the property that for some $K \geq 1$ and every positive integer n there are n vectors of the form $y_i = \sum_{j \in \sigma_i} a_j x_j$, $i = 1, \dots, n$, where the sets σ_i are pairwise disjoint and the sequence $\{y_i\}$ is equivalent, with constant K , to the unit vector basis in ℓ_2^n . See also [JMST] for a generalization of this fact for more general lattices. It was left open in these two papers (and specifically asked in Problem 3.A of [DS]) whether $\{y_i\}$ can be chosen to be a block basis, i.e., whether the sets σ_i can be chosen to be successive, that is, $\max\{j; j \in \sigma_k\} < \min\{j; j \in \sigma_l\}$ for all $k < l$.

The initial motivation of this paper was to solve this question in the negative. As is well known the Haar basis, $\{h_{n,i}\}_{n=0, i=1}^\infty$, in $L_p(0, 1)$, $1 < p < \infty$, in its common order, has a block basis (i.e., the Rademacher functions) equivalent to the unit vector basis of ℓ_2 . It follows from the main results of this paper (Theorems 2.2 and 3.1) that the initial segments of the Haar system can be rearranged so that they will not have (the finite version of) this property anymore. We denote the new order by $\preceq$.

Note that the space $(\sum_{N=1}^\infty \oplus [h_{n,i}]_{n=0, i=1}^{N, 2^n})_p$ is isomorphic to ℓ_p . (Here $[x_i]_{i \in A}$ denotes the closed linear span of $\{x_i\}_{i \in A}$.) Let $\{b_i\}_{i=1}^\infty$ be the natural (relative to the order $\preceq$) basis of $(\sum_{N=1}^\infty \oplus [h_{n,i}]_{n=0, i=1}^{N, 2^n})_p$, i.e., the basis composed of the successive conjunction of the bases $\{h_{n,i}\}_{n=0, i=1}^{N, 2^n}$, each in its $\preceq$ -order. Clearly, $\{b_i\}_{i=1}^\infty$ is not equivalent to the natural basis of ℓ_p as long as $p \neq 2$, and it easily follows from the previous paragraph that

---

Received by the editors September 5, 1995 and, in revised form, February 23, 1996.

1991 *Mathematics Subject Classification*. Primary 46B07, 60G42.

The first author was supported by the Austrian Academy of Sciences (APART Program).

The second author was supported in part by BSF.

Both authors participated in the Workshop in Linear Analysis and Probability, Texas A&M University.

• For each p , $1 < p \neq 2 < \infty$, and for each K , $1 \leq K < \infty$, there is an n such that no block basis of $\{b_i\}_{i=1}^\infty$ of length n is K -equivalent to the unit vector basis of ℓ_2^n .

Thus, in the case $2 < p < \infty$, the sequence $\{b_i\}_{i=1}^\infty$ gives the required counterexample to the problem above.

In Section 1 below we introduce the new order on the initial segments of the Haar system and determine some of its properties. In Section 2 we prove the main theorem (2.2) which states that, for $2 < p < \infty$, every block basis of such an initial segment of the Haar system is well equivalent to a diagonal of the unit vector basis of ℓ_p^k and a subsequence of a permutation of the Haar system. In Section 3 we deal with the case $1 < p < 2$ and prove a somewhat weaker property of such block bases in Theorem 3.1 but which is also enough to deduce • above.

1. THE ORDER

An interval of the form $I = [\frac{i-1}{2^n}, \frac{i}{2^n})$, $n = 0, 1, \dots$, $i = 1, 2, \dots, 2^n$, will be called a dyadic interval. The left half of the interval $I = [\frac{i-1}{2^n}, \frac{i}{2^n})$ is the interval $[\frac{2i-2}{2^{n+1}}, \frac{2i-1}{2^{n+1}})$ and the right half of I is $[\frac{2i-1}{2^{n+1}}, \frac{2i}{2^{n+1}})$. The Haar function $h_I = h_{n,i}$ associated with I is the function

$$h_I(t) = \begin{cases} 1, & \text{if } t \text{ is in the left half of } I, \\ -1, & \text{if } t \text{ is in the right half of } I. \end{cases}$$

The common order on $\{h_I\}$ is the lexicographic order on $\{(n, i)\}$. Fixing N we would like to introduce a different order, $\preceq$, on the set $T = T_N$ of dyadic intervals of length larger than or equal to 2^{-N} .

Definition. $I \preceq J$ if either I and J are disjoint and I is to the left of J , or, I is contained in J .

In terms of the natural dyadic tree structure of T , Figure 1 describes the order, for $N = 4$.

FIGURE 1

Proposition 1.1 below describes the order intervals with respect to the order $\preceq$. We first need two definitions.

Definition. Given two dyadic intervals $I \subseteq J$

1. The *cone* $\mathcal{C} = \mathcal{C}(I, J)$ of dyadic intervals between I and J is the unique collection of dyadic intervals, $\mathcal{C} = \{C_1, \dots, C_k\}$, satisfying
  - a) $k = \log_2 \frac{|J|}{|I|} + 1$,
  - b) $C_1 = I, C_k = J$,
  - c) $|C_s| = \frac{1}{2}|C_{s+1}|, C_s \subset C_{s+1}$, for $s \in \{1, \dots, k-1\}$.
2. The *right fill-up* of the cone $\mathcal{C}$ is the collection of dyadic intervals, $\mathcal{R} = \mathcal{R}(I, J) = \bigcup_{s=1}^{k-1} \mathcal{U}_{s+1}$, where $\mathcal{U}_{s+1} = \emptyset$ if C_s is the right half of C_{s+1} and

$$\mathcal{U}_{s+1} = \{J : |J| \geq 2^{-N}, J \subseteq C_{s+1} \setminus C_s\}$$

if C_s is the left half of C_{s+1} .

Proposition 1.1. Let $J_1, J_2 \in T_N$ satisfy $J_1 \preceq J_2$, then there exists a (unique) collection $\mathcal{L} = \{L_1, \dots, L_m\}$ of pairwise disjoint dyadic intervals satisfying:

- 1) $|L_i| \leq \frac{1}{2}|L_{i-1}|$ if $1 \leq i \leq m-1$;
- 2) $|L_{m-1}| = |L_m|$ (if $m \geq 2$);
- 3a) $\overline{L_i}$ and $\overline{L_{i+1}}$ intersect in exactly one point (the left endpoint of $\overline{L_{i+1}}$);
- 3b) L_{i+1} lies right of L_i ;
- 4) $J_1 \subseteq L_1, J_2 = L_m$ such that

$$(1) \quad \{I \in T_N : J_1 \preceq I \preceq J_2\} = \mathcal{C}(J_1, L_1) \cup \mathcal{R}(J_1, L_1) \cup_{i=2}^m \mathcal{M}_i$$

where $\mathcal{M}_i = \{I \in T_N : I \subseteq L_i\}$.

Proof. Assume first that $J_1 \subseteq J_2$. Then the assertion is that a dyadic interval I satisfies $J_1 \preceq I \preceq J_2$ if and only if $I \in \mathcal{C}(J_1, J_2) \cup \mathcal{R}(J_1, J_2)$.

The if part is clear. Assume now $J_1 \preceq I \preceq J_2$. Clearly, I cannot be disjoint from J_2 : otherwise it would have to be to the left of J_2 and thus to the left of J_1 contradicting $J_1 \preceq I$. So $I \subseteq J_2$. If $J_1 \subseteq I$ then by the definition of a cone, $I \in \mathcal{C}(J_1, J_2)$. If not, then I lies to the right of J_1 . Then there exists $C_{i_0} \in \mathcal{C}(J_1, J_2)$, so that $I \subset C_{i_0}$ and $I \cap C_{i_0-1} = \emptyset$. We then claim that $C_{i_0} \setminus C_{i_0-1}$ is the right half of C_{i_0} , and then $I \in \mathcal{R}(J_1, J_2)$.

Indeed, if $C_{i_0} \setminus C_{i_0-1}$ is the left half of C_{i_0} , then I lies to the left of C_{i_0-1} and thus to the left of J_1 , a contradiction. This concludes the proof of this special case and we turn to the general case.

Let D_1 be the minimal dyadic interval which contains both J_1 and J_2 . If $D_1 = J_2$ put $m = 1$ and $L_1 = D_1$.

If D_1 strictly contains J_2 then take the left half of D_1 and call it L_1 . Observe that $J_2 \subseteq D_1 \setminus L_1 = R_1$ where R_1 is the right half of D_1 .

Let I be the dyadic interval of length 2^{-N} which has the same left endpoint as R_1 . Let $D_2 \subseteq R_1$ be the minimal dyadic interval which contains both I and J_2 .

If $D_2 = J_2$ put $m = 2, L_2 = D_2$. If D_2 strictly contains J_2 then take the left half of D_2 and call it L_2 .

Observe that $J_2 \subseteq D_2 \setminus L_2 = R_2$ where R_2 is the right half of D_2 .

Continuing in that manner (let I be the dyadic interval of length 2^{-N} which has the same left endpoint as R_2 . Let $D_3 \subseteq R_2$ be the minimal dyadic interval which contains both I and J_2 ...), the process stops after finitely many steps and we get a collection $\mathcal{L} = \{L_1, \dots, L_m\}$ of pairwise disjoint dyadic intervals satisfying 1

through 4. We are left with the task of proving (1). Clearly, the right-hand side in (1) is contained in the left-hand side. To prove the other containment, note that if $J_1 \preceq I \preceq J_2$ then none of the $L_i, i \leq m$ is strictly contained in I . Suppose not; then for some i , $D_i \subseteq I$ and from $J_2 \subseteq D_i \subseteq I$ we get $J_2 \prec I$. A contradiction.

It follows that there exists i_0 , so that $I \subseteq L_{i_0}$. Suppose not; then using that $\bigcup_{i=1}^m L_i$ is an interval (not necessarily dyadic) I is either to the right of L_m or to the left of L_1 . Hence $I \prec J_1$ or $J_2 \prec I$. A contradiction.

If $i_0 \geq 2$ then $I \in \mathcal{M}_{i_0}$ and we are done, so, we may suppose $J_1 \preceq I \preceq L_1$. This is exactly the special situation described in the beginning of this proof and we conclude that in this case $I \in \mathcal{C}(J_1, L_1) \cup \mathcal{R}(J_1, L_1)$. $\square$

Next we would like to describe the interaction between two successive order intervals. Let $I_1 \prec I_2 \prec J_1 \prec J_2$ and put

$$\mathcal{B}_1 = \{I \in T_N ; I_1 \preceq I \preceq I_2\}, \quad \mathcal{B}_2 = \{I \in T_N ; J_1 \preceq I \preceq J_2\}.$$

Let $\mathcal{L}_1 = \{L_1^1, \dots, L_{m(1)}^1\}$ be the maximal (with respect to inclusion) elements of $\mathcal{B}_1$ as given in Proposition 1.1 and let $\mathcal{L}_2 = \{L_1^2, \dots, L_{m(2)}^2\}$ be the maximal elements of $\mathcal{B}_2$. Let $B_i = \bigcup_{I \in \mathcal{B}_i}$. Clearly B_1 and B_2 may intersect.

Lemma 1.2. a) B_1 may have intersection only with L_1^2 ; i.e.,

$$(B_2 \setminus L_1^2) \cap B_1 = \emptyset.$$

b) If $B_1 \cap B_2 \neq \emptyset$ then $L_1^2 \supseteq L_{m(1)}^1$.

Proof. Note that if $I \preceq J_1$ then $I \cap L_j^2 = \emptyset$ for $j \geq 2$. Indeed, if $I \cap L_j^2 \neq \emptyset$ for some $j \geq 2$ then either

Case 1. $I \subseteq L_j^2$ and then I lies to the right of L_1^2 and L_1^2 lies above J_1 . Hence by the definition of the order $J_1 \preceq I$, a contradiction. Or,

Case 2. $I \supset L_j^2$ and then $J_1 \prec L_j^2 \prec I$. A contradiction again.

This proves a.

To prove b, note that $L_{m(1)}^1 = I_2 \prec J_1 \preceq L_1^2$. If $L_{m(1)}^1 \not\subseteq L_1^2$ then necessarily $L_{m(1)}^1$ is to the left of L_1^2 . But then L_j^1 is to the left of L_1^2 for all j and $B_1 \cap L_1^2 = \emptyset$. Since also $B_1 \cap (B_2 \setminus L_1^2) = \emptyset$ we get a contradiction to $B_1 \cap B_2 \neq \emptyset$. $\square$

It follows from the lemma that if an interval I from $\mathcal{B}_2$ intersects B_1 then it belongs to either $\mathcal{C}(J_1, L_1^2)$ or to $\mathcal{R}(J_1, L_1^2)$. The second possibility is easily ruled out since in this case I lies to the right of J_1 while every interval in $\mathcal{B}_1$ lies to the left or below J_1 , and one gets (using part b of the lemma for the ‘moreover’ part)

Proposition 1.3. If $B_1 \cap B_2 \neq \emptyset$ then every interval in $\mathcal{B}_2$ which intersects B_1 belongs to $\mathcal{C}(J_1, L_1^2)$. Moreover,

$$\{I \in \mathcal{B}_2 ; I \cap B_1 \neq \emptyset\} = \mathcal{C}(J_1, L_1^2) \cap \mathcal{C}(I_2, L_1^2).$$

2. BLOCK BASES OF $\{h_I\}_{I \in T_N}$ WITH RESPECT TO THE ORDER $\preceq$

Given $M \in \mathbf{N}$ and elements of T_N ,

$$I_1 \prec J_1 \prec I_2 \prec J_2 \prec \dots \prec I_M \prec J_M,$$

consider the order intervals $\mathcal{B}_i = \{I \in T_N ; I_i \preceq I \preceq J_i\}$. Let $\mathcal{L}_i = \{L_1^i, \dots, L_{m(i)}^i\}$ be the maximal (with respect to inclusion) elements of $\mathcal{B}_i$ as given in Proposition

1.1. Put also $\mathcal{C}_i = \mathcal{C}(I_i, L_1^i)$ —the cone of intervals between I_i and L_1^i , and let $\mathcal{G}_i = \mathcal{B}_i \setminus \mathcal{C}_i$. Note that, by Proposition 1.3,

- a) The sets $\bigcup_{I \in \mathcal{G}_i} I$, $i = 1, 2, \dots, M$ are pairwise disjoint, and
- b) the sets $L_1^i = \bigcup_{I \in \mathcal{C}_i} I$, $i = 1, 2, \dots, M$ are nested; i.e., for $i < j$, either $L_1^i \cap L_1^j = \emptyset$ or $L_1^i \subset L_1^j$. In the latter case $|L_1^i| \leq |L_1^j|/2$.

Now let $\{Z_i\}_{i=1}^M$ be a block basis of $\{h_I\}$ with respect to the blocking $\{\mathcal{B}_i\}$; i.e.,

$$Z_i = \sum_{I \in \mathcal{B}_i} \alpha_I h_I, \quad i \leq M.$$

Consider

$$X_i = \sum_{I \in \mathcal{C}_i} \alpha_I h_I, \quad Y_i = \sum_{I \in \mathcal{G}_i} \alpha_I h_I.$$

Then $X_i + Y_i = Z_i$. Clearly, the Y_i 's have disjoint supports (with respect to the underlying measure space—the interval $[0, 1]$) and thus are equivalent to multiples of the canonical basis elements of ℓ_p^M . To analyze the $\{X_i\}$ we must now study more carefully the cones $\{\mathcal{C}_i\}$ and their interaction. We start with the following observation which is self evident.

Observation. Let $\mathcal{C}_1 = \{I_1, \dots, I_k\}$ with $I_1 \subset I_2 \subset \dots \subset I_k$, $|I_i| = \frac{1}{2}|I_{i+1}|$, and $\mathcal{C}_2 = \{J_1, \dots, J_l\}$ with $J_1 \subset J_2 \subset \dots \subset J_l$, $|J_i| = \frac{1}{2}|J_{i+1}|$ be two cones of intervals consisting of distinct intervals (i.e., $I_i \neq J_j$ for all i, j). Suppose that $J_l \subset I_k$ and put, for convenience of notation, $I_0 =$ the right half of I_1 and $I_{-1} = \emptyset$; then there exists $0 \leq i \leq k$ such that $J_l \subseteq I_i \setminus I_{i-1}$.

Let $\mathcal{C} = \{I_1, \dots, I_k\}$ be a cone of intervals and put, for $i = 1, \dots, k$, $D_i = I_i \setminus I_{i-1}$. Then the D_i , $i = 1, \dots, k$, are disjoint dyadic intervals such that $|D_i| = \frac{1}{2}|I_i|$ and $D_i \subset I_i$. Also, $|I_k| \leq 2|\bigcup_{i=1}^k D_i| \leq |I_k|$.

Returning to the X_i 's, denoting $\mathcal{C}_i = \{I_1^i, \dots, I_{k(i)}^i\}$ and by $\mathcal{D}_i = \{D_1^i, \dots, D_{k(i)}^i\}$ the collection of intervals obtained from $\mathcal{C}_i$ by the process above, put

$$\phi_i = \sum_{j=1}^{k(i)} \alpha_{I_j^i} h_{D_j^i}.$$

(To obtain ϕ_i from X_i we just replace $h_{I_j^i}$ by $h_{D_j^i}$.)

Lemma 2.1. The map $X_i \mapsto \phi_i$, $i = 1, \dots, M$, extends to an isomorphism from $(\text{span } \{X_i\}, \|\cdot\|_{L_p})$ to $(\text{span } \{\phi_i\}, \|\cdot\|_{L_p})$ where the isomorphism constant depends on p alone, $1 < p < \infty$.

Proof. The assertion follows easily from a theorem of Semenov [S]. We prefer to give an ad hoc proof.

Extend the map $X_i \mapsto \phi_i$, $i = 1, \dots, M$, to the map given by $T_p : h_{I_j^i} \mapsto h_{D_j^i}$ from $(\text{span } \{h_{I_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_p})$ to $(\text{span } \{h_{D_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_p})$. Note that the square function $\left(\sum (a_j^i)^2 (h_{I_j^i})^2\right)^{1/2}$ pointwise dominates $\left(\sum (a_j^i)^2 (h_{D_j^i})^2\right)^{1/2}$ and consequently T_p is bounded in L_p , $1 < p < \infty$, with bound depending on p alone.

The unconditionality of the Haar system also implies that the dual space to $(\text{span } \{h_{I_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_p})$ can be naturally identified with $(\text{span } \{h_{I_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_q})$, $\frac{1}{p} + \frac{1}{q} = 1$, with equivalent norms, and a similar identification can be done

between the dual of $(\text{span } \{h_{D_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_p})$ and $(\text{span } \{h_{D_j^i}\}_{j=1, i=1}^{k(i), M}, \|\cdot\|_{L_q})$. Since the conjugate operator to T_p is T_q^{-1} , we are done. $\square$

We shall show in Proposition 2.3 below that, for $2 < p < \infty$, $\{X_i\}$ is well equivalent to an appropriately normalized subsequence of a permutation of the Haar system and thus we'll get the following theorem.

Theorem 2.2. *For every $2 < p < \infty$ there exists a constant K_p , $1 \leq K_p < \infty$, such that, for all N , every block basis of $\{h_I\}_{I \in T_N}$ (with respect to the order $\preceq$) is equivalent, with constant at most K_p , to a sequence of the kind $\{a_i x_i \oplus b_i e_i\}_{i=1}^n$, where $\{x_i\}_{i=1}^n$ is a subsequence of a permutation of $\{h_I\}_{I \in T_N}$ and $\{e_i\}_{i=1}^n$ is the unit vector basis in ℓ_p^n .*

It may be instructive to recall here the local version of the Gamlen-Gaudet Theorem proved by the first named author in [M].

Theorem. *For every $1 < p < \infty$ there exists a constant K_p , $1 \leq K_p < \infty$, such that, for all N , every subsequence of $\{h_I\}_{I \in T_N}$ (of length k , say) spans a space isomorphic, with constant at most K_p , to ℓ_p^k .*

Problem. Can Theorem 2.2 (and Theorem 3.1 below) be strengthened to show that for every $1 < p < \infty$ there exists a constant K_p , $1 \leq K_p < \infty$, such that, for all N , every block basis of $\{h_I\}_{I \in T_N}$ (with respect to the order $\preceq$) of length k , say, spans a space isomorphic, with constant at most K_p , to ℓ_p^k ?

An easy consequence of Theorem 2.2 (see the proof of Theorem 3.1) is that there exists a subset $A \subseteq \{1, \dots, M\}$ with $|A| \geq M/2$ and such that $\{Z_i\}_{i \in A}$ is well equivalent to the unit vector basis of $\ell_p^{|A|}$, $2 < p < \infty$. This turns out to be true also for $1 < p < 2$ and is easier to deduce than Theorem 2.2. We shall do it in the next section. Of course this is enough for the solution of problem 3.A in [DS]. We continue now with the preparations to the proof of Theorem 2.2.

Let $\mathcal{E}$ be the collection of the largest intervals in the cones $\mathcal{C}_i$, $i = 1, \dots, M$, i.e., $\mathcal{E} = \{I_{k(i)}^i\}$. Let $G_1(\mathcal{E})$ be the subcollection of maximal (with respect to inclusion) intervals out of $\mathcal{E}$. Note that any two distinct intervals in $G_1(\mathcal{E})$ are disjoint. Define, by induction, $G_k(\mathcal{E})$ to be the subcollection of maximal intervals out of $\mathcal{E} \setminus \bigcup_{i=1}^{k-1} G_i(\mathcal{E})$.

Given $C = I_{k(i)}^i \in \mathcal{E}$ denote $\mathcal{D}(C) = \mathcal{D}_i$ (that is the collection of intervals obtained as successive differences of the intervals of the cone that C is the maximal element of). Denote also by $\mathcal{D}^{1/2}(C)$ the collection of the left and right halves of the intervals in $\mathcal{D}(C)$ and put

$$\mathcal{H}_k = \bigcup_{C \in G_k(\mathcal{E})} \mathcal{D}(C), \quad \mathcal{H}_k^{1/2} = \bigcup_{C \in G_k(\mathcal{E})} \mathcal{D}^{1/2}(C), \quad k = 1, 2, \dots$$

We now define an increasing sequence of σ -fields by

$$\mathcal{E}_1 = \mathcal{A}\{\mathcal{H}_1\}, \quad \mathcal{E}_k = \mathcal{A}\{\mathcal{E}_{k-1}, \mathcal{H}_k, \mathcal{H}_{k-1}^{1/2}\}$$

where $\mathcal{A}\{\mathcal{H}\}$ denotes the σ -field generated by $\mathcal{H}$. We are now ready to prove

Proposition 2.3. *Let $\mathcal{C}_i = \{I_1^i, \dots, I_{k(i)}^i\}$, $i = 1, \dots, M$, be pairwise disjoint cones of intervals. Using the notation above (for the $\mathcal{D}$'s), let $\phi_i = \sum_{j=1}^{k(i)} \alpha_{I_j^i} h_{D_j^i}$, $i = 1, \dots, M$, be a normalized, in L_p , sequence. Then $\{\phi_i\}_{i=1}^M$ is equivalent, with*

constant depending only on p , $2 < p < \infty$, to a permutation of a subsequence of the normalized Haar system.

Proof. Reindex the sequence $\{\phi_i\}$, using $\mathcal{E}$ as an index set, by putting $\phi_i = \phi_C$ if $C = I_{k(i)}^i$. Then for $C \in G_k(\mathcal{E})$ we have by the **Observation** above

- 1) ϕ_C is $\mathcal{E}_{k+1}$ measurable,
- 2) $\mathbf{E}(\phi_C | \mathcal{E}_k) = 0$,
- 3) $C \neq C' \in G_k(\mathcal{E})$ implies $\text{supp } \phi_C \cap \text{supp } \phi_{C'} = \emptyset$.

By Burkholder's theorem (see [B, (21.5)]), we have for all scalars $\{a_C, C \in \mathcal{E}\}$

$$\left\| \sum_k \sum_{C \in G_k(\mathcal{E})} a_C \phi_C \right\|_{L_p}^{C_p} \lesssim \max \left\{ \left\| \left(\sum_k \sum_{C \in G_k(\mathcal{E})} a_C^2 \mathbf{E}(\phi_C^2 | \mathcal{E}_k) \right)^{1/2} \right\|_{L_p}, \left(\sum_{C \in \mathcal{E}} |a_C|^p \right)^{1/2} \right\}.$$

To prove the proposition it is thus enough to prove that, for every k and every $C \in G_k(\mathcal{E})$, there exists an $I \in \mathcal{D}(C)$ such that ratio of the two functions $\mathbf{E}(\phi_C^2 | \mathcal{E}_k)$ and $\mathbf{E}(h_I^2 | \mathcal{E}_k)$ is bounded and bounded away from 0 by constants depending only on p . This follows easily, by rearranging and rescaling, from the following lemma. $\square$

Lemma 2.4. *Let $p > 2$, let $D_m = [2^{-m}, 2^{-(m-1)})$ and let $\beta_m > 0$ so that $\sum_{m=1}^n |\beta_m|^p = 1$. Let h_n be the L_p -normalized Haar function having D_m as support. Then for $\phi = \sum_{m=1}^n \beta_m h_m$ we have*

$$2^{-n(1-\frac{2}{p})} \leq \int_0^1 |\phi|^2 dt \leq 1.$$

Consequently, there exists an $1 \leq m \leq n$ satisfying

$$1 \leq \int_0^1 |\phi|^2 dt \bigg/ \int_0^1 |\bar{h}_m|^2 dt \leq 2^{(1-\frac{2}{p})}.$$

Proof of Lemma 2.4. The right-hand side inequality is clear. To prove the left-hand side, notice that this is the same problem as finding the (square of the) minimal L_2 norm of a function which is constant on each of the D_m 's and has L_p norm equal to 1. The minimum is attained for the function with minimal support, i.e., for the characteristic function of D_n normalized in L_p and thus is equal to $2^{-n(1-\frac{2}{p})}$.

Let $1 \leq m \leq n$ be such that

$$2^{-m(1-\frac{2}{p})} \leq \int_0^1 |\phi|^2 dt \leq 2^{-(m-1)(1-\frac{2}{p})};$$

then

$$\begin{aligned} 1 &\leq \int_0^1 |\phi|^2 dt \bigg/ 2^{-m(1-\frac{2}{p})} = \int_0^1 |\phi|^2 dt \bigg/ \int_0^1 |\bar{h}_m|^2 dt \\ &\leq 2^{(1-\frac{2}{p})} \int_0^1 |\phi|^2 dt \bigg/ 2^{-(m-1)(1-\frac{2}{p})} \leq 2^{(1-\frac{2}{p})}. \end{aligned}$$

This ends the proof of the lemma and thus of Proposition 2.3 and Theorem 2.2. $\square$

3. THE CASE $1 < p < 2$

Here we prove:

Theorem 3.1. *For every $1 < p < 2$ there exists a constant K_p , $1 \leq K_p < \infty$, such that, for all N , every normalized block basis $\{Z_i\}_{i=1}^M$ of $\{h_I\}_{I \in T_N}$ (with respect to the order $\preceq$) contains a subsequence $\{Z_i\}_{i \in A}$ with $|A| \geq M/2$ which is equivalent, with constant at most K_p , to the unit vector basis of $\ell_p^{|A|}$.*

An identical statement holds for $2 < p < \infty$ and its proof follows that of the proof below using Theorem 2.2. With a bit more effort one can also avoid referring to Theorem 2.2. We do not reproduce this case here.

For the proof of Theorem 3.1 we shall need only the first two paragraphs in Section 2, that is up to, and not including, the Observation. We start with a lemma similar to one used in ([F, p. 156]).

Lemma 3.2. *Let T be a finite tree with partial order $<$. Put*

$$A_0 = \{t \in T ; t \text{ is a minimal element}\}$$

and

$$A_1 = \{t \in T ; t \notin A_0 \text{ and there is exactly one maximal } s \text{ with } s < t\}.$$

Then $|T| < 2|A_0| + |A_1|$.

Proof. Assume first $A_1 \neq \emptyset$. Let $t \in A_1$ and let s be the unique maximal element with $s < t$. Put $T' = T \setminus \{s\}$ and let A'_0 and A'_1 be the corresponding “ A_0 ” and “ A_1 ” associated to this new tree. Note that $|A'_0| = |A_0|$. (If $s \in A_0$ then $t \in A'_0$ while if $s \notin A_0$, $A'_0 = A_0$.) Note also that $|A'_1| = |A_1| - 1$. (If $s \notin A_1$ then $A'_1 = A_1 \setminus \{t\}$ while if $s \in A_1$ then $A'_1 = A_1 \setminus \{s\}$.) We thus reduce the problem to proving $|T'| < 2|A'_0| + |A'_1|$.

Continuing reducing the tree in this fashion we finally reach a tree for which $A_1 = \emptyset$, i.e., it branches everywhere : every node has at least two offsprings. For such a tree the conclusion is well known and easy. $\square$

Proof of Theorem 3.1. We follow the notation of the beginning of Section 2. Since the sequence $\{Y_i/\|Y_i\|\}_{i=1}^M$ is isometrically equivalent to the unit vector basis of ℓ_p^M , it is enough to prove that there exists a subset $A \subseteq \{1, \dots, M\}$ with $|A| \geq M/2$ for which $\{X_i/\|X_i\|\}_{i \in A}$ is equivalent, with constant depending only on p , to the unit vector basis of $\ell_p^{|A|}$.

Recall that the supports (with respect to the underlying measure space) of the X_i 's are the sets L_1^i , $i = 1, \dots, M$, and they are nested; i.e., any two of them are either disjoint or one is contained in the other. Let

$$A_0 = \{i; L_1^i \text{ contains no other } L_1^j\}$$

and

$$A_1 = \{i; L_1^i \text{ contains exactly one other maximal } L_1^j\}$$

(i.e., $\exists j \neq i$ with $L_j \subset L_i$ but no two disjoint L_j 's with this property). By Lemma 3.2 $|A_0| + |A_1| \geq M/2$. The set $A = A_0 \cup A_1$ can be decomposed into sets B_k , $k = 1, \dots, K$, with $L_1^i \cap L_1^j = \emptyset$ if i and j are in different B_k 's and $L_1^i \subset L_1^j$ or $L_1^j \subset L_1^i$ if they are in the same one. It is thus enough to prove that $\{X_i/\|X_i\|\}_{i \in B_k}$ is equivalent, with constant depending only on p , to the unit vector basis of $\ell_p^{|B_k|}$ and

thus, without loss of generality that, if $L_1^1 \subset L_1^2 \subset \cdots \subset L_1^m$, then $\{X_i/\|X_i\|\}_{i=1}^m$ is well equivalent to the unit vector basis of ℓ_p^m .

Let $1 < i \leq m$ and let $\mathcal{C}_i = \{I_1, \dots, I_r\}$ be the cone of intervals associated with X_i . Note that there exists a $1 \leq s \leq r$ with $L_{i-1} \subseteq I_s \setminus I_{s-1}$ (I_0 is one of the two halves of I_1). Clearly $\|X_i|_{[0,1] \setminus L_{i-1}}\| \geq c\|X_i\|$ for some constant $c > 0$ depending only on p . Now a simple square function argument shows that the sequence $\{X_i\}$ well dominates the sequence $\{X_i|_{[0,1] \setminus L_{i-1}}\}$. This finishes the proof since the functions $X_i|_{[0,1] \setminus L_{i-1}}$, $i = 1, \dots, m$, are disjoint. $\square$

REFERENCES

- [B] D.L. Burkholder, *Distribution function inequalities for martingales*, Ann. Prob. **1** (1973), 19-43. MR **51**:1944
- [DS] L.E. Dor and T. Starbird, *Projections of L_p onto subspaces spanned by independent random variables*, Compositio Math. **39** (1979), 141-175. MR **82e**:46043
- [F] C.L. Fefferman, *The uncertainty principle*, Bull. AMS **9** (1983), 129-206. MR **85f**:35001
- [JMST] W.B. Johnson, B. Maurey, G. Schechtman, and L. Tzafriri, *Symmetric structures in Banach spaces*, Mem. AMS **217** (1979). MR **82j**:46025
- [M] P.F.X. Müller, *A local version of a result of Gamlen-Gaudet*, Israel J. Math. **63** (1988), 212-222. MR **89m**:46016
- [S] E.M. Semenov, *On the equivalence in L_p of rearrangements of the Haar system*, English translation in Soviet Math. Dokl. **19** (1978), 1292-1294.

INSTITUT FÜR MATHEMATIK, J. KEPLER UNIVERSITÄT, 4040 LINZ, AUSTRIA

E-mail address: paul.mueller@jk.uni-linz.ac.at

Current address: Department of Mathematics, Yale University, 10 Hill House Avenue, New Haven, Connecticut 06520

E-mail address: muller@math.yale.edu

DEPARTMENT OF THEORETICAL MATHEMATICS, THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: mtschec@weizmann.weizmann.ac.il