KEEPING ADDITIVITY OF THE NULL IDEAL SMALL

JINDŘICH ZAPLETAL

(Communicated by Andreas R. Blass)

ABSTRACT. We shall show that various statements are consistent with additivity of the null ideal equal to $\aleph_1$; for example, “all branchless trees of size $\aleph_1$ are special”, (S) conjecture and “there are only five cofinal types of directed posets of size $\aleph_1$ ”.

0. INTRODUCTION

In this paper we provide machinery for proving that a certain large class of forcings has a certain regularity property. The class in question includes posets used for

- (1) specializing branchless trees [S],
- (2) (S) conjecture [T2],
- (3) embedding the poset for adding $\aleph_1$ Cohen reals into a given poset of uniform density $\aleph_1$ [SZ],
- (4) classification of directed posets [T1] or transitive relations [T3] of size $\aleph_1$,
- (5) other “side condition” combinatorics on $\aleph_1$; e.g. shooting an uncountable set through a coherent sequence on ω_1 [T2].

The regularity property we obtain implies preservation of additivity of the ideal of Lebesgue null sets. As a corollary, it is consistent with ZFC set theory that additivity of the null ideal is $\aleph_1$ and all statements obtainable through (1)–(5) above hold, that is, all branchless trees of size $\aleph_1$ are special, (S) conjecture holds etc. Thus a definite limitation on a canonical variation of the powerful “side condition” method has been exacted for the first time.

Our notation follows the set-theoretical standard as set forth in [J]. In a forcing notion, $p \leq q$ means “ p is more informative than q ”. A tree T of height ω_1 is special if there is a function $f : T \rightarrow \omega$ with $s < t$ in T implying $f(s) \neq f(t)$. Trees grow upwards. (S) conjecture is the statement “every hereditarily separable Hausdorff space is hereditarily Lindelöf”. The symbol $\mathcal{N}$ denotes the collection of Borel null sets, often confused with their Borel codes. H_κ is the set of all sets of hereditary cardinality $< \kappa$. If N is an elementary submodel of H_κ and $P \in N$ is a forcing, a condition $p \in P$ is called N -master if for every dense set $D \subset P$ in N , the set $D \cap N$ is predense below p .

Received by the editors November 8, 1995 and, in revised form, February 26, 1996.

1991 *Mathematics Subject Classification*. Primary 03E35, 03E50.

Research at MSRI partially supported by NSF grant # DMS 9022140. The author wishes to thank Itay Neeman for asking the original inspiring question.

1. LOCALIZATION

Definition 1. Let $\mathfrak{F} \subset {}^\omega\omega$ and let e be a positive integer. The family $\mathfrak{F}$ is said to be e -localized if there exists a function $h : \omega \rightarrow [\omega]^{<\aleph_0}$ such that

- (1) $|h(n)| \leq n^e$ for every integer n ,
- (2) for every $f \in \mathfrak{F}$ there is an integer $n \in \omega$ so that for every $m > n$, $f(m) \in h(m)$ holds.

If $e = 1$ then the family $\mathfrak{F}$ is said to be localized.

The relevance of the above definition is revealed in the result of Bartoszyński [B, BJ Section 2.3.A] saying that for a transitive model M of ZFC the following are equivalent:

- (1) $M \cap {}^\omega\omega$ is localized;
- (2) $M \cap {}^\omega\omega$ is e -localized for some positive integer e ;
- (3) the union of all Lebesgue measure zero Borel sets coded in M has measure zero.

There is a natural c.c.c. forcing for making the set of ground model reals localized [Tr] and there are some preservation theorems for “the set of ground model reals is not localized” [JS], [BJ]. We shall prove that a large class of forcings preserves unlocalized families in a strong sense.

Definition 2. (1) [JS] Let κ be a large regular cardinal and $N \prec H_\kappa$. We say that a function $f \in {}^\omega\omega$ is N -big if for every $h : \omega \rightarrow [\omega]^{<\aleph_0}$ in the model N with $|h(n)| \leq n$ the set $\{m \in \omega : f(m) \notin h(m)\}$ is infinite.

(2) Let P be a forcing. We say that P is friendly if for every $p \in P$, every large enough regular cardinal κ , every countable elementary submodel $N \prec H_\kappa$ with p, P in N and every N -big function $f \in {}^\omega\omega$ there is an N -master condition $q \leq p$ such that $q \Vdash \dot{f}$ is $N[G]$ -big”.

The important point is that it is possible to iterate friendly forcings preserving the statement “the family of ground model reals is not localized” or equivalently, “ $\bigcup(\mathcal{N} \cap V) \notin \mathcal{N}$ ” –Lemma 13. Obviously, a finite iteration of friendly forcings is friendly and friendliness is inherited by regular subposets. In [JS] it is proved that the random algebra as well as every σ -centered forcing is friendly. We considerably extend these results.

2. SPECIALIZING TREES

The purpose of this section is to prove that the usual specializing forcing for a branchless tree of height ω_1 is friendly. The technique will be of great use in the next section. For now, fix a tree T of height ω_1 and no branches of length ω_1 . There is no restriction on the size of levels of T .

Definition 3. (1) If a, b are disjoint finite subsets of T then we say that a and b fit together if for every $s \in a$ and $t \in b$, s and t are incompatible as elements of T .

(2) The specialization forcing is $P = \{p : p \text{ is a finite function from } T \text{ to } \omega \text{ such that } s <_T t \text{ in } \text{dom}(p) \text{ implies } p(s) \neq p(t)\}$ ordered by reverse inclusion.

Lemma 4. Let $\{a_\alpha : \alpha \in \omega_1\}$ be a family of pairwise disjoint finite subsets of T . Then there is an infinite set $Y \subset \omega_1$ such that $a_\alpha : \alpha \in Y$ pairwise fit together.

Proof. The usual proof of c.c.c.-ness of P [S, p. 103] shows that there are $\alpha \neq \beta$ with a_α, a_β fitting together. The lemma follows from the Erdős-Dushnik-Miller partition relation $\omega_1 \rightarrow (\omega_1, \omega)$ applied to the function $f : [\omega_1]^2 \rightarrow 2$ defined by $f(\alpha, \beta) = 0$ iff a_α, a_β fit together. $\square$

Now assume that κ is a large regular cardinal, $N \prec H_\kappa$ is a countable submodel with $T \in N$ and f is N -big. We shall prove that $P \Vdash \check{f}$ is $N[G]$ -big. Then, since the forcing P is c.c.c., any condition in it is N -master and it witnesses the friendliness of P as desired.

For contradiction, let $p \in P, n \in \omega$ and $\dot{h} \in N$ be such that

- (1) $p \Vdash$ “for all $m \in \omega$, $|\dot{h}(m)| \leq m$ ”,
- (2) $p \Vdash$ “for every $m > n$, $\check{f}(m) \in \dot{h}(m)$ ”.

Let $p_0 = p \cap N \in N$. By c.c.c.-ness of P , by strengthening the condition p if necessary one can arrange that $p_0 \Vdash$ “for all $m \in \omega$, $|\dot{h}(m)| \leq m$ ”.

Work in N . Fix an integer $m > n$ and by a tree induction construct a tree $X \subset {}^{<\omega}(\omega + 1)$, a partition $X = X_0 \cup X_1$ and a function $F : X \rightarrow P$ so that

- (1) the empty sequence $\emptyset$ is in X and $F(\emptyset) = p_0$,
- (2) $s \subset t$ in X implies $F(s) \geq F(t)$ in P ,

and for each $s \in X$ with $lth(s) = i$ exactly one of the following holds:

- (3) either, there is a sequence $\langle q_j : j \in \omega \rangle$ such that each $q_j \leq F(s)$ forces in P that $i \in \dot{h}(m)$, and moreover, the sets $dom(q_j \restriction F(s)) : j \in \omega$ pairwise fit together. In this case, $s \in X_0$, the set of successors of s in X is exactly $\{s^\frown \langle j \rangle : j \in \omega\}$ and $F(s^\frown \langle j \rangle) = q_j$,
- (4) or, no such sequence exists. Then $s \in X_1$, the only successor of s in the tree X is $s^\frown \langle \omega \rangle$ and $F(s^\frown \langle \omega \rangle) = F(s)$.

A set $o(X) \subset \omega$ is defined by $i \in o(X)$ iff there exists a function $G : X \rightarrow \omega$ such that for every sequence $s \in X$ of length i , if $\forall j \in i$ $s(j) > G(s \restriction j)$ then $s \in X_0$.

Claim 5. $|o(X)| \leq m$.

Proof. Suppose for a contradiction that there are $m + 1$ elements of $|o(X)|$, enumerated in the increasing order as i_0 through i_m . Pick witnesses $G_k : k \leq m$ for $i_k \in o(X)$. Then for every sequence $s \in X$ of length $i_m + 1$ such that $\forall j \leq i_m$ $\forall k \leq m$ $s(j) > G_k(s \restriction j)$ (and there are plenty of these) the value $F(s)$ as an element of P forces each one of the $m + 1$ distinct integers $i_k : k \leq m$ into the set $\dot{h}(m)$. But this is absurd, since $p \geq F(s)$ and $p \Vdash |\dot{h}(m)| \leq m$. $\square$

Claim 6. $f(m) \in o(X)$.

Proof. The proof of this fact takes place outside of the model N . To define the witness $G : X \rightarrow \omega$ for $f(m) \in o(X)$, consider two cases:

- (1) $s \in X_1$. Then let $G(s) = 0$.
- (2) $s \in X_0$. By (3) above, $\{a_j = dom(F(s^\frown \langle j \rangle) \restriction F(s)) : j \in \omega\}$ is a family of pairwise disjoint fitting finite subsets of the tree T . There is an integer j_0 such that for every $j > j_0$, the sets a_j and $dom(p \restriction p_0)$ fit together; set $G(s) = j_0$.

The existence of an integer j_0 as in (2) above can be demonstrated as follows. By elementarity of the model N , if $u \in N \cap T$ and $t \in dom(p \restriction p_0)$ are compatible as elements of the tree T , then necessarily $u <_T t$, for if $t <_T u$ then $t \in N$, contradicting the definition of the condition p_0 . By the mutual fitting of the a_j 's, only finitely many of the sets $a_j \subset N \cap T$ can have nonempty intersection with the

finitely many linearly ordered subsets $\{u \in N \cap T : u <_T t\} : t \in \text{dom}(p \setminus p_0)$ of the tree T . Any j_0 larger than the indexes of these sets will do.

Why does the function G have the desired properties? Well, choose an arbitrary sequence $s \in X$ of length $f(m)$ with $\forall j \in f(m) \ s(j) > G(s \restriction j)$. It is necessary to verify that $s \in X_0$. By the definition of G and X , the condition $F(s)$ is compatible with p . By induction on $\alpha \in \omega_1$ construct a sequence $\langle a_\alpha : \alpha \in \omega_1 \rangle$ so that

- (1) a_α are finite pairwise disjoint subsets of the tree T ,
- (2) for every $\alpha \in \omega_1$ there is $q \leq F(s)$ such that $q \Vdash "f(m) \in \dot{h}(m)"$ and $a_\alpha = \text{dom}(q \setminus F(s))$.

Note that any sequence $\langle a_\alpha : \alpha \in \beta \rangle$ of countable length β satisfying (1,2) above for $\alpha \in \beta$ can be prolonged further to some $\langle a_\alpha : \alpha \in \beta+1 \rangle$ such that (1,2) continue to hold. For if this were not the case, by elementarity there would be a witness $\langle a_\alpha : \alpha \in \beta \rangle \in N$ which cannot be prolonged. However, such a sequence from the model N can be prolonged using $a_\beta = \text{dom}(p \setminus p_0)$, with the condition $q = F(s) \cup p$ witnessing the property (2) at β .

By elementarity of the model N , there is a sequence $\langle a_\alpha : \alpha \in \omega_1 \rangle$ satisfying (1,2) above in N . It follows from Lemma 4 that the case (3) of definition of the tree X is valid at s and $s \in X_0$. $\square$

Now within the model N , for each integer $m > n$ construct a tree X_m and a set $o(X_m)$ as above. The function $g : \omega \setminus n \rightarrow [\omega]^{<\aleph_0}$ defined by $g(m) = o(X_m)$ is in the model N and contradicts the assumption of f being N -big. Therefore, we have proved

Theorem 6. *Let T be a tree of height ω_1 without branches of height ω_1 . Then the standard T -specialization forcing is friendly.*

Remark. The result becomes rather trivial if the tree T is supposed to have countable levels. In such a case, it is easy to prove that every real added by the specialization forcing comes from a Cohen-generic extension. Thus the specialization forcing for T must necessarily be friendly by results of [JS].

Remark. The same technology can be used to demonstrate friendliness of a number of finite condition forcings, whose c.c.c. is proved in a certain canonical manner. For example, let $\langle f_\alpha : \alpha \in \omega_1 \rangle \subset {}^\omega\omega$ be a modulo finite increasing unbounded sequence of increasing functions. Let the partition $H : [\omega_1]^2 \rightarrow 2$ be defined as $H(\alpha, \beta) = 0$ if $\alpha < \beta$ and there is an integer $n \in \omega$ with $f_\alpha(n) > f_\beta(n)$. It is known [T2] that the poset P of finite 0-homogeneous sets ordered by reverse inclusion is c.c.c. and destroys the unboundedness of the sequence. Using the same method as above, it is possible to show that P is friendly. The following is open:

Question 7. Is OCA [T2] consistent with additivity of the null ideal equal to $\aleph_1$?

3. SIDE CONDITION FORCINGS

The purpose of this section is to define the class of ideal-based forcings and to prove friendliness of elements of this class. Our scheme is designed to comprehend many side condition forcings as used in the work of S. Todorcevic [T1], [T2], [T3] and others. Let A be a set of finite subsets of ω_1 ordered by $\sqsubseteq$ and let $\mathcal{I}$ be an ideal on ω_1 such that the following axioms are satisfied:

- (A) $\sqsubseteq$ refines the inclusion, for each $a \in A$ and $\beta \in \omega_1$ $a \cap \beta \sqsubseteq a$ holds and if a, b are both in A and $\sqsubseteq$ -compatible then $a \cup b \in A$ is their $\sqsubseteq$ -upper bound.

- (B) $\mathfrak{I}$ contains singletons, every $\mathfrak{I}$ -positive set has a countable $\mathfrak{I}$ -positive subset and the σ -ideal $\sigma\mathfrak{I}$ generated by $\mathfrak{I}$ is proper.

Moreover, for each $a \in A$ there are

- (C) a $\sigma\mathfrak{I}$ positive set $Z \subset \omega_1$ such that $a \sqsubseteq a \cup \{\beta\} \in A$ holds for every $\beta \in Z$,
 (D) an $\mathfrak{I}$ -large set $Y \subset \omega_1$ such that for every $\beta \in Y$ the implication $a \cap \beta \sqsubseteq (a \cap \beta) \cup \{\beta\} \in A \rightarrow a \sqsubseteq a \cup \{\beta\} \in A$ holds.

The pair $\langle A, \sqsubseteq \rangle$ is to be understood as a problematic finite-condition forcing construction for which c.c.c. cannot be proved, or which collapses $\aleph_1$ outright. The existence of the ideal ensures that there is a way to add a $\sqsubseteq$ -filter which meets many dense subsets of $\langle A, \sqsubseteq \rangle$. Fix a large regular cardinal κ . The ideal-based forcing P derived from $A, \sqsubseteq, \mathfrak{I}$ has the following form:

$P = \{f : f \text{ is a finite function from } \omega_1 \text{ to } H_\kappa, \text{ for } \alpha \in \text{dom}(f) \ f(\alpha) = \langle M_\alpha, \xi_\alpha \rangle\}$
 and

- (E) the set $\text{body}(f) = \{\xi_\alpha : \alpha \in \text{dom}(f)\}$ is in A ,
 (F) every M_α is a countable elementary submodel of H_κ containing $A, \sqsubseteq, \mathfrak{I}, f \restriction \alpha$,
 (G) $\xi_\alpha \notin \bigcup (M_\alpha \cap \mathfrak{I})$.

The order on P is defined by $f \leq g$ if $g \subset f$ and $\text{body}(g) \sqsubseteq \text{body}(f)$.

The forcing P adds a $\sqsubseteq$ -filter $\{a \in A : a = \text{body}(p) \text{ for some } p \in G\}$ which meets all dense subsets of $\langle A, \sqsubseteq \rangle$ which are in some sense large as measured by $\mathfrak{I}$.

Frequently, for the sake of preservation of $\aleph_2$ one needs to consider an amended variation of P which has matrices of models as side conditions instead of just an $\in$ -chain of models as above [T2]. We call such forcings *amended ideal-based*; since our proofs carry over to the class of amended ideal-based forcings with only more complicated notation, we concentrate on the class of ideal-based forcings proper. The point of course is that this class is reasonably wide; indeed, our scheme includes many of the side condition posets used in the literature. The following fact provides a by no means complete list.

Fact 8. The forcings for the following problems are (amended) ideal-based:

- (1) (S) conjecture [T2],
- (2) making a poset of uniform density $\aleph_1$ add $\aleph_1$ Cohen reals [SZ],
- (3) classification of transitive relations on $\aleph_1$ [T1], [T3],
- (4) shooting an uncountable set through a coherent sequence on ω_1 [T2].

Proof. We consider the case of (S) conjecture. As in [T2], it is only necessary to cope with the following problem. Let ω_1 be equipped with topology $\mathfrak{T}$ so that the space $(\omega_1, \mathfrak{T})$

- (1) is hereditarily separable, that is, for every $X \subset \omega_1$ there is a countable subset $Y \subset X$ with the same closure,
- (2) is not hereditarily Lindelöf, and it is even right separated, that is, for each $\alpha \in \omega_1$ there is an open set O_α such that $\alpha \in O_\alpha$ and the closure of O_α is a subset of $\alpha + 1$.

We wish to violate the hereditary separability of the space $(\omega_1, \mathfrak{T})$ by introducing an uncountable discrete subset to it. The forcing for doing that [T2] can be cast as an ideal-based forcing derived from $A = [\omega_1]^{<\aleph_0}$, $a \sqsubseteq b$ just in case $a \subset b$ and for every $\xi \in (b \setminus a)$ and every $\zeta \in a$ $\xi \notin O_\zeta$; furthermore, $\mathfrak{I} = \{X \subset \omega_1 : \text{the closure of } X \text{ is countable}\}$. It is not difficult to check the axioms (A) through (D) in the definition of ideal-based. (B) follows from hereditary separability and (D) from $\mathfrak{I}$ -smallness of every O_α .

The intended uncountable discrete set will be $\bigcup\{body(f) : f \in G\}$, where $G \subset P$ is a generic filter. $\square$

Theorem 9. *Any ideal-based forcing P is proper and friendly.*

Proof. Let $A, \sqsubseteq, \mathfrak{I}, \kappa$ be the parameters from which P is defined. To prove the properness, let $p_0 \in P$, λ be a large regular cardinal, let $N \prec H_\lambda$ be a countable elementary submodel with $p_0, A, \sqsubseteq, \mathfrak{I}, \kappa \in N$, and let $\delta = N \cap \omega_1$. By (C) there is a countable ordinal ξ such that $\xi \notin \bigcup(\mathfrak{I} \cap N)$ and $body(p_0) \sqsubseteq body(p_0) \cup \{\xi\} \in A$. Let $p_1 = p_0 \cup \{\langle \delta, \langle N \cap H_\kappa, \xi \rangle \rangle\}$. Obviously, $p_1 \leq p_0$ and we shall show that p_1 is a master condition for the model N . Thus, for any dense set D of P which happens to be in N , the set $D \cap N$ must be proved predense below p_1 . Fix $p_2 \leq p_1$ and a dense set $D \in N$; we shall produce conditions $p_5 \leq p_2$ and $q \in D \cap N$ with $p_5 \leq q$, completing the proof of properness. By strengthening p_2 if necessary, it can be assumed that there is an element of D above p_2 .

Let $p_3 = p_2 \cap N$. Obviously $p_3 \in P \cap N$ and $p_2 \leq p_3$, by (A). The whole point of the proof is to find a way of carefully extending p_3 within N while preserving compatibility with p_2 . Let $k = |p_2 \setminus p_3|$ and $\xi_0 \dots \xi_{k-1}$ enumerate $body(p_2) \setminus body(p_3)$ in the increasing order. By induction on $l < k$ define sets $S(t)(l) \subset \omega_1$ for all $t \in {}^{<\omega}\omega_1$ simultaneously by

- (1) $S(t)(0) = \{\zeta \in \omega_1 \setminus \max(body(p_3) \cup rng(t)) : \exists p_4 \leq p_3 \text{ such that } p_4 \text{ has an element of } D \text{ above it and } body(p_3) \cup rng(t) \sqsubseteq body(p_4) = body(p_3) \cup rng(t) \cup \{\zeta\} \in A\}$.
- (2) $S(t)(l+1) = \{\zeta \in \omega_1 \setminus \max(body(p_3) \cup rng(t)) : body(p_3) \cup rng(t) \sqsubseteq body(p_3) \cup rng(t) \cup \{\zeta\} \in A \text{ and the set } S(t \smallfrown \langle \zeta \rangle)(l) \text{ is } \mathfrak{I}\text{-positive}\}$.

Claim 10. The set $S(\langle \rangle)(k-1)$ is $\mathfrak{I}$ -positive.

Proof. Note that the system $\{S(t)(l) : t \in {}^{<\omega}\omega_1, l < k\}$ belongs to all the models mentioned in p_2 above δ since it is in $N \cap H_{\aleph_2}$. The claim will be proved by contradiction. If $S(\langle \rangle)(k-1)$ were an element of $\mathfrak{I}$, by induction on $l < k$ one could show that $\xi_l \notin S(\langle \xi_{l'} : l' < l \rangle)(k-1-l)$. But this is a contradiction to the case (1) of the definition of the system $\{S(t)(l) : t \in {}^{<\omega}\omega_1, l < k\}$, since $\xi_{k-1} \in S(\langle \xi_l : l < k-1 \rangle)(0)$ as witnessed by the condition p_2 . $\square$

Now by induction on $l < k$ build ζ_l, T_l, X_l so that

- (1) for $l \leq k$, $a_l = \langle \zeta_{l'} : l' < l \rangle$ is an increasing sequence of countable ordinals larger than $\max(body(p_3))$ in N ,
- (2) $T_l \in N$ is an $\mathfrak{I}$ -positive countable subset of the $\mathfrak{I}$ -positive set $S(a_l)(k-l-1)$, by (B),
- (3) $body(p_2)$ and $body(p_3) \cup rng(a_l)$ are $\sqsubseteq$ -compatible and an $\mathfrak{I}$ -large set $X_l \subset \omega_1$ is a witness to (D) for $body(p_2) \cup rng(a_l)$,
- (4) $\zeta_l \in T_l \cap X_l$.

By the construction, $a_k \in N$, $body(p_2)$ and $body(p_3) \cup rng(a_k)$ are $\sqsubseteq$ -compatible and moreover, there is a condition $p_4 \leq p_3$ such that there is an element of D above it and $body(p_4) = body(p_3) \cup rng(a_k)$. By the elementarity of the model N , there are such p_4 and $q \in D$ above it already in N . By the definition of the forcing P , $p_5 = p_4 \cup p_2$ is a lower bound of p_4 and p_2 and has $q \in D \cap N$ above it as desired.

The friendliness of P is proved by a trick similar to the one in Section 2. Let us adopt the framework from the proof of properness of P , in particular, choose

$p_0, N \dots$ and the master condition $p_1 \leq p_0$ for the model N as constructed above. Let $f \in {}^\omega\omega$ be an N -big function. We shall show that $p_1 \Vdash \check{f}$ is $N[G]$ -big".

For contradiction, let $p_2 \leq p_1, \dot{h} \in N$ and $n \in \omega$ be such that

- (1) $p_2 \Vdash$ "for all integers $m, |\dot{h}(m)| \leq m$ ",
- (2) $p_2 \Vdash$ "for all integers $m > n, \dot{f}(m) \in \dot{h}(m)$ ".

Let $p_3 = p_2 \cap N$. Then $p_3 \geq p_2$ is in N and by strengthening the condition p_2 if necessary we may assume that $p_3 \Vdash$ "for all integers $m, |\dot{h}(m)| \leq m$ ". Let $k = |p_2 \setminus p_3| \geq 1$.

Work in N . Fix an integer $m > n$. By a tree induction construct a tree $X \subset {}^{<\omega}\omega$, its subset X_0 and functions $F : X \rightarrow P, T : X_0 \rightarrow V$ so that:

- (1) the empty sequence 0 is in X and $F(0) = p_3$,
- (2) for every $s \subset t$ both in X , $F(t) \leq F(s)$ holds in P .

Moreover, at each $s \in X$, exactly one of the two following cases will hold.

Case 1. There is a tree $T(s)$ on ${}^{\leq k}\omega_1$ such that

- (a) the empty sequence is in $T(s)$ and T consists of increasing sequences of ordinals above $\max(\text{body}(F(s)))$,
- (b) for every $t \in T(s)$ of length $< k$ the set $\{\zeta \in \omega_1 : t \smallfrown \langle \zeta \rangle \in T(s)\}$ is countable and $\mathfrak{I}$ -positive; moreover for each ζ in this set, $\text{body}(F(s)) \cup \text{rng}(t) \sqsubseteq \text{body}(F(s)) \cup \text{rng}(t) \cup \{\zeta\} \in A$,
- (c) for every $t \in T(s)$ of length k (i.e. a terminal node) there is a condition $q_t \leq F(s)$ in P such that $\text{body}(q_t) = \text{body}(F(s)) \cup \text{rng}(t)$ and $q_t \Vdash$ "length(s) $\in \dot{h}(m)$ ".

In this case, let $s \in X_0$, $T(s)$ will be the value of T at s and $s \smallfrown \langle l \rangle \in X$ for all $l \in \omega$. Also, $F(s \smallfrown \langle l \rangle) : n \in \omega$ enumerates the set $\{q_t : t \in T(s) \text{ is a terminal node}\}$.

Case 2. No such tree exists. Then $s \notin X_0$, the only successor of the sequence s in X is $s \smallfrown \langle 0 \rangle$ and $F(s) = F(s \smallfrown \langle 0 \rangle)$.

This completes the inductive definition of the tree X in N . Define a set $o(X) \subset \omega$ by $i \in o(X)$ if there exists a collection $\{A(s, t) : s \in X_0, t \in T(s) \text{ is not a terminal node}\}$ of $\mathfrak{I}$ -large sets so that for each sequence $u \in X$ of length i if (*) below holds for all $j < i$ then $u \in X_0$.

- (*) If $u \restriction j \in X_0$ then $F(u \restriction (j+1)) = q_t$ for some terminal node $t \in T(u \restriction j)$ such that $\forall l < k \ t(l+1) \in A(u \restriction j, t \restriction l)$.

Claim 11. $|o(X)| \leq m$.

Proof. For contradiction, suppose that there are $m+1$ elements of $o(X)$ enumerated in the increasing order as i_0 through i_m . Pick witnesses $\{A_l(s, t) : s \in X_0, t \in T(s) \text{ is not a terminal node}\}$ for $i_l \in o(X)$ and define $\{A(s, t) : s \in X_0, t \in T(s) \text{ is not a terminal node}\}$ by $A(s, t) = \bigcap_{l \leq m} A_l(s, t)$. Now each $A(s, t)$ is still an $\mathfrak{I}$ -large set and so it is possible to find a sequence $u \in X$ of length $i_m + 1$ such that (*) holds for each $j < \text{length}(u)$. But then, from the construction of X and F it follows that $F(u) \Vdash$ " $\{i_0 \dots i_m\} \subset \dot{h}(m)$ ", which is absurd since $F(u) \leq p_3$ and $p_3 \Vdash$ " $|\dot{h}(m)| \leq m$ ". $\square$

Claim 12. $f(m) \in o(X)$.

Proof. It is necessary to define the witness collection; our candidate lies outside of N . For $s \in X_0$ and $t \in T(s)$ there will be two cases:

- (1) Either $\text{body}(p_2)$ and $\text{body}(F(s)) \cup \text{rng}(t)$ are $\sqsubseteq$ -compatible. In such a case let $A(s, t)$ be an $\mathfrak{I}$ -large witness to (D) for $\text{body}(p_2) \cup \text{body}(F(s)) \cup \text{rng}(t)$.
- (2) Otherwise let $A(s, t) = \omega_1$.

Now suppose that a sequence $u \in X$ of length $f(m)$ satisfies $(*)$ for all $j < f(m)$. By the definition of $A(s, t)$ and the tree X , necessarily $\text{body}(F(u))$ and $\text{body}(p_2)$ are $\sqsubseteq$ -compatible and therefore p_2 and $F(u)$ are compatible conditions in P . The proof of $u \in X_0$ is essentially a repetition of the proof of the properness of P with the condition p_2 replaced with $F(u) \cup p_2$ and the phrase “element of D above it” replaced with “forces $f(m)$ into $\dot{h}(m)$ ”. $\square$

Now the definition of $X, o(X)$ was uniform for integers $m > n$. Thus within the model N there is a sequence $X_m, o(X_m) : m > n$ such that $|o(X_m)| \leq m$ and $f(m) \in o(X_m)$ for every $m > n$. But then the sequence $o(X_m) : m > n$ in the model N , understood as a function of m , contradicts N -bigness of the function f . $\square$

It should be remarked that while ideal-based forcings preserve $\text{add}(\mathcal{N})$, they can add dominating functions. If $\langle f_\alpha : \alpha \in \omega_1 \rangle \subset {}^\omega \omega$ is a modulo finite increasing unbounded sequence of increasing functions then it is possible to derive an S-space from it [T2]. Then the ideal-based forcing killing that space adds a function which modulo finite dominates all $f_\alpha : \alpha \in \omega_1$.

4. CONCLUSION

At last, we are in a position to construct some interesting models of set theory with the additivity of the null ideal equal to $\aleph_1$. The classical iteration vehicle gives

Lemma 13. *Let $\langle P_\alpha : \alpha \leq \theta, \dot{Q}_\alpha : \alpha < \theta \rangle$ be a countable support iteration of forcings such that $P_\alpha \Vdash \dot{Q}_\alpha$ is friendly” for each $\alpha < \theta$. Then $P_\theta \Vdash$ “the union of the null sets coded in the ground model is not null”.*

It seems likely that in fact P_θ is a friendly forcing, but we have no argument for that.

Proof. By induction on $\beta \leq \theta$ we shall demonstrate that $P_\beta \Vdash$ “the union of the null sets coded in the ground model is not null”. The limit step is handled by [BJ, Theorem 6.3.41]. For the successor step, assume that $P_\beta \Vdash$ “the union of the null sets coded in the ground model is not null”; we shall prove the same statement for $P_{\beta+1}$. Choose a generic filter $G \subset P_\beta$ and work in $V[G]$. It is enough to show that $Q_\beta \Vdash “V \cap {}^\omega \omega$ is not localized”. For contradiction, suppose $q \in Q_\beta, \dot{h}$ are such that $q \Vdash_{Q_\beta}$ “for all $n \in \omega$, $|\dot{h}(n)| \leq n$ and $\dot{h}$ localizes $V \cap {}^\omega \omega$ ”. Choose a large regular cardinal κ and a countable elementary submodel $N \prec H_\kappa$ with $q, Q_\beta, \dot{h} \in N$.

Claim 14. There is an N -big function $f \in V \cap {}^\omega \omega$.

Proof. By an easy bookkeeping argument, there is a function $k : \omega \rightarrow [\omega]^{<\aleph_0}$ such that $\forall n \in \omega$ $|k(n)| \leq n^2$ and for every function $l \in N$ with $l : \omega \rightarrow [\omega]^{<\aleph_0}$, $\forall n \in \omega$ $|l(n)| \leq n$ there is an integer m_0 such that for every $m > m_0$, $l(m) \subset k(m)$. By the induction hypothesis, $V \cap {}^\omega \omega$ is not 2-localized and therefore there is a function $f \in V \cap {}^\omega \omega$ such that the set $\{m \in \omega : f(m) \notin k(m)\}$ is infinite. Obviously, the function f is N -big. $\square$

The postulated friendly master condition $r \leq q$ for N, f contradicts the assumption $q \Vdash “\exists n \forall m > n f(m) \in \dot{h}(m)”$. $\square$

Therefore, starting from a model of the Continuum Hypothesis, any sufficiently generic iteration of length ω_2 of proper, $\aleph_2$ -p.i.c. [S, pg. 262] and friendly forcings will provide for $\Vdash$ “ $\text{add}(\mathcal{N}) = \aleph_1, \mathfrak{c} = \aleph_2$, all branchless trees of size $\aleph_1$ are special, (S) conjecture holds, every poset of uniform density $\aleph_1$ adds $\aleph_1$ Cohen reals etc.” In the construction, it is necessary to use amended ideal-based forcings in order to ensure $\aleph_2$ -c.c. of the resulting iteration. The standard bookkeeping arguments are left to the reader.

REFERENCES

- [B] T. Bartoszyński, *Additivity of measure implies additivity for category*, Trans. Amer. Math. Soc. **281** (1984), 209–213. MR **85b**:03083
- [BJ] T. Bartoszyński and H. Judah, *Set theory. On the structure of the real line*, A K Peters, Wellesley, Massachusetts, 1995. MR **96k**:03002
- [J] T. Jech, *Set Theory*, Academic Press, New York, 1978. MR **80a**:03062
- [JS] H. Judah and S. Shelah, *The Kunen-Miller chart*, Jour. Symbolic Logic **55** (1990), 909–927. MR **91g**:03097
- [RS] J. Raisonier and J. Stern, *The strength of measurability hypothesis*, Israel Jour. Math. **50** (1985), 337–349. MR **87d**:03129
- [Tr] J. Truss, *Connection between different amoeba algebras*, Fund. Math. **130** (1988), 137–155. MR **90h**:03036
- [S] S. Shelah, *Proper Forcing*, Springer-Verlag, New York, 1982. MR **84h**:03002
- [SZ] S. Shelah and J. Zapletal, *Embeddings of Cohen algebras*, Adv. Math. (to appear).
- [T1] S. Todorcevic, *Directed sets and cofinal types*, Trans. Amer. Math. Soc. **290** (1985), 711–723. MR **87a**:03084
- [T2] ———, *Partition problems in topology*, Amer. Math. Soc., Providence, 1989. MR **90d**:04001
- [T3] ———, *A classification of transitive relations on ω_1* , preprint, 1995.

M. S. R. I., 1000 CENTENNIAL DRIVE, BERKELEY, CALIFORNIA 94720

E-mail address: `jindra@msri.org`

Current address: Mailcode 253-37, California Institute of Technology, Pasadena, California 91125

E-mail address: `jindra@cco.caltech.edu`