ON THE GANEA CONJECTURE FOR MANIFOLDS

YU. B. RUDYAK

(Communicated by Thomas Goodwillie)

ABSTRACT. Using a result of Singhof, we prove that $\text{cat}(M \times S^m) = \text{cat } M + 1$ provided M is a connected closed PL manifold with $\dim M \leq 2 \text{cat } M - 3$ and S^m is the m -sphere, $m > 0$.

Let $\text{cat } X$ denote the Lusternik–Schnirelmann category of X (normalized, i.e., $\text{cat } S^m = 1$). There is a long standing Ganea conjecture that $\text{cat}(X \times S^m) = \text{cat } X + 1$ for every connected finite CW -complex X and every $m > 0$, see [1, Problem 2]. In [2, Corollary 6.7] Singhof proved the following

Singhof’s Theorem. (i) *Let M be a connected closed PL manifold such that*

$$\text{cat } M \geq \frac{\dim M + m + 2}{2}.$$

Then $\text{cat}(M \times S^m) = \text{cat } M + 1$ provided $m > 0$.

(ii) *In particular, if*

$$\text{cat } M \geq \frac{\dim M + 3}{2}$$

then $\text{cat}(M \times S^1) = \text{cat } M + 1$.

However, it seems that nobody noted that Singhof’s Theorem implies a stronger result. Namely, the following theorem is valid:

Theorem. *Let M be a connected closed PL manifold such that*

$$\text{cat } M \geq \frac{\dim M + 3}{2}.$$

Then $\text{cat}(M \times S^m) = \text{cat } M + 1$ for every $m > 0$.

Proof. First, we prove that $\text{cat}(M \times T^r) = \text{cat } M + r$ where T^r is the r -torus. We prove this by induction. For $r = 1$ this follows from (ii). Now, suppose that $\text{cat}(M \times T^r) = \text{cat } M + r$. Then

$$\begin{aligned} \text{cat}(M \times T^r) &= \text{cat } M + r \geq \frac{\dim M + 3}{2} + r \\ &\geq \frac{\dim M + r + 3}{2} = \frac{\dim(M \times T^r) + 3}{2}. \end{aligned}$$

Received by the editors March 7, 1996.

1991 *Mathematics Subject Classification.* Primary 55M30; Secondary 57Q99, 57R19.

The author was partially supported by Deutsche Forschungsgemeinschaft.

So, by (ii), $\text{cat}(M \times T^{r+1}) = \text{cat } M + r + 1$.

Now, given $m > 0$, we prove that $\text{cat}(M \times S^m) = \text{cat } M + 1$. Choose r such that

$$\text{cat}(M \times T^r) = \text{cat } M + r \geq \frac{\dim(M \times T^r) + m + 2}{2}$$

(for example, $r \gg m$). Then, by (i),

$$\text{cat}(M \times T^r \times S^m) = \text{cat}(M \times T^r) + 1 = \text{cat } M + r + 1.$$

Now, if $\text{cat}(M \times S^m) \neq \text{cat } M + 1$ then $\text{cat}(M \times S^m) \leq \text{cat } M$. But then

$$\text{cat}(M \times S^m \times T^r) \leq \text{cat}(M \times S^m) + \text{cat } T^r \leq \text{cat } M + r.$$

This is a contradiction. $\square$

Corollary. *For every connected closed PL manifold M there exists a natural number k such that $\text{cat}(M \times T^k \times S^m) = \text{cat}(M \times T^k) + 1$ for every $m > 0$.*

Proof. Indeed, you can find k such that

$$\text{cat}(M \times T^k) \geq k \geq \frac{\dim(M \times T^k) + 3}{2}.$$

$\square$

REFERENCES

1. T. Ganea, *Some problems on numerical homotopy invariants*, Symposium in Algebraic Topology, Seattle 1971, Edited by P. Hilton (Lect. Notes in Math., vol. 249), Berlin Heidelberg New York: Springer, 1971, pp. 23–30. MR **49**:3910
2. W. Singhof, *Minimal coverings of manifolds with balls*, Manuscripta Math. **29** (1979), 385–415. MR **80k**:55012

MATHEMATISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 288, D-69120 HEIDELBERG, GERMANY

E-mail address: july@mathi.uni-heidelberg.de