ISOMORPHICALLY EXPANSIVE MAPPINGS IN ℓ_2

J. GARCÍA-FALSET, A. JIMÉNEZ-MELADO, AND E. LLORÉNS-FUSTER

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We show that for any renorming $\|\cdot\|$ of ℓ_2 , the well known fixed point free mappings by Kakutani, Baillon and others are not nonexpansive.

1. INTRODUCTION

Let C be a subset of a Banach space $(X, \|\cdot\|)$. A mapping $T : C \rightarrow C$ is called *k-lipschitzian* ($k > 0$) if $\|T(x) - T(y)\| \leq k \|x - y\|$ for all $x, y \in C$.

Nonexpansive mappings are those which have Lipschitz constant $k = 1$. A mapping $T : C \rightarrow C$ is said to be *uniformly k-lipschitzian* if every iteration T^n is *k-lipschitzian*, i.e. if for every positive integer n , and $x, y \in C$

$$\|T^n(x) - T^n(y)\| \leq k \|x - y\|.$$

We say that X has the fixed point property (FPP) if every nonexpansive mapping $T : C \rightarrow C$ defined on a nonempty convex and weakly compact subset C of X has a fixed point.

If $T : C \rightarrow C$ is a nonexpansive mapping with respect to some norm $|\cdot|$ on X equivalent to the norm $\|\cdot\|$, then it is straightforward to see that T is also uniformly lipschitzian with respect to the norm $\|\cdot\|$. Hence, in order to look for a fixed point free $|\cdot|$ -nonexpansive mapping we must seek this mapping among those which are uniformly lipschitzian with respect to the norm $\|\cdot\|$.

It is known that bounded closed convex subsets of ℓ_2 have the fixed point property for nonexpansive mappings, but it is unknown whether the same is true for bounded closed convex subsets of a Banach space X where X is ℓ_2 with an equivalent norm. In this paper it is shown that certain uniformly lipschitzian mappings which are known to be fixed point free in the unit ball of ℓ_2 are also not nonexpansive relative to any renorming of ℓ_2 . This shows that one strategy for answering the above question is precluded. At the same time it shows that the class of mappings which are uniformly lipschitzian on ℓ_2 is strictly larger than the class of mappings which are nonexpansive with respect to some equivalent norm on ℓ_2 . Theorems 1 and 2 serve mainly as vehicles for reaching this conclusion.

Received by the editors November 28, 1995 and, in revised form, March 18, 1996.

1991 *Mathematics Subject Classification*. Primary 47H09, 47H10.

Key words and phrases. Nonexpansive mappings, uniformly lipschitzian mappings, fixed points.

This research has been partially supported by D.G.I.C.Y.T. PB93-1177-C02-02 and D.G.I.C.Y.T. PB94-1496.

2. PRELIMINARIES

We recall now some famous examples of uniformly lipschitzian mappings in ℓ_2 .

Let $\|\cdot\|_2$ be the usual euclidean norm on the sequence Hilbert space ℓ_2 , let B be the closed $\|\cdot\|_2$ -unit ball and let $S : \ell_2 \rightarrow \ell_2$ be the right shift operator defined by

$$S(x_1, x_2, \dots) = (0, x_1, x_2, \dots).$$

Then the mapping $K : B \rightarrow B$ defined by

$$K(x) = \frac{(1 - \|x\|_2)e + S(x)}{\|(1 - \|x\|_2)e + S(x)\|_2},$$

where $e = (1, 0, \dots)$, is *uniformly* 2-lipschitzian and has no fixed point in B ([GKT]).

Later on, Baillon ([B]) found another example of a uniformly $\frac{\pi}{2}$ -lipschitzian mapping without fixed point in B . This is the mapping $G : B \rightarrow B$ defined by

$$G(x) = \begin{cases} \cos\left(\frac{\pi}{2}\|x\|_2\right)e + \sin\left(\frac{\pi}{2}\|x\|_2\right)\frac{S(x)}{\|x\|_2}, & x \neq 0, \\ e, & x = 0 \end{cases}$$

(for details about this map, see [T]).

Both examples are modifications of an earlier one due to Kakutani (see [GK]): for $0 < r < 1$, define $K_r : B \rightarrow B$ by

$$K_r(x) = r(1 - \|x\|_2)e + S(x).$$

Then K_r have the (non uniform) Lipschitz constant $\sqrt{1+r^2}$ and has no fixed point in B .

3. THE RESULTS

The announced results will follow easily from the following lemma.

Lemma 1. *Let $(X, |\cdot|)$ be a Banach space and let B_X, S_X be the closed unit ball and the unit sphere of X respectively. Suppose that $T : B_X \rightarrow X$ is a mapping which is nonexpansive with respect to some norm $\|\cdot\|$ on X , equivalent to $|\cdot|$. Then we have that*

$$d = \inf_{y \in S_X} |T(0) - T(y)| \leq 1.$$

Proof. There exist positive constants α, β such that, for every $v \in X$,

$$\alpha|v| \leq \|v\| \leq \beta|v|.$$

Since T is $\|\cdot\|$ -nonexpansive, for any $y \in S_X$ we have that

$$\alpha|T(0) - T(y)| \leq \|T(0) - T(y)\| \leq \|y\| \leq \beta$$

and then

$$\alpha d \leq \|y\| \leq \beta$$

for all $y \in S_X$. Thus, for every $v \in X$ we have that

$$\alpha d|v| \leq \|v\| \leq \beta|v|.$$

Indeed, an induction argument shows that

$$\alpha d^n|v| \leq \|v\| \leq \beta|v|$$

for every positive integer n and every $v \in X$, which is impossible unless $d \leq 1$. $\square$

The above lemma will allow us to give a fixed point result. Recall that a sequence (X_n) of finite dimensional subspaces of a Banach space X is called a *Schauder finite dimensional decomposition* (FDD) of X , if every $x \in X$ has a unique representation of the form $x = \sum x_i$ with $x_i \in X_i$ for every $i \in \mathbb{N}$. If X is a Banach space with a FDD (X_n) , and $x \in X$, $\text{supp}(x)$ denotes, as usual, the set of positive integers i such that $x_i \neq 0$. M. A. Khamsi ([K]) defined, for a Banach space X with a FDD, the coefficient $\beta_p(X)$ in the following way: For $p \in [1, \infty)$, $\beta_p(X)$ is the infimum of the set of numbers λ such that

$$(\|x\|^p + \|y\|^p)^{1/p} \leq \lambda \|x + y\|$$

for every $x, y \in X$ which verify $\text{supp}(x) < \text{supp}(y)$, that is, $\max[\text{supp}(x)] < \min[\text{supp}(y)]$.

For example, $\beta_p(l_p) = 1$ ($1 \leq p < \infty$), and $\beta_2(J) = 1$, where J is the James space which consists of all sequences $x = (x_n) \in c_0$ such that

$$\|x\| := \sup \{ (x_{p_1} - x_{p_2})^2 + \dots + (x_{p_{n-1}} - x_{p_n})^2 \}^{1/2}$$

is finite (the supremum is taken for every n and for every finite increasing sequence of positive integers (p_i)).

Now, for such Banach spaces we can state the following

Theorem 1. *Let $(X, |\cdot|)$ be a Banach space with a FDD such that $\beta_p(X) = 1$ for some $p \in [1, \infty)$. Let B_X, S_X be the closed unit ball and the unit sphere of X , respectively, and let $T : B_X \rightarrow X$ be a mapping such that*

- a) $T(S_X) \subset S_X$ and
- b) $\text{supp}[T(0)] < \text{supp}(v)$ for every $v \in T(S_X)$.

If T is nonexpansive with respect to some equivalent norm on X then T has a fixed point. Specifically, $T(0) = 0$.

Proof. Since $\beta_p(X) = 1$, conditions a) and b) imply that

$$|T(0)|^p + |T(y)|^p \leq |T(0) - T(y)|^p$$

for all $y \in S_X$, and then we have that

$$d = \inf_{y \in S_X} |T(0) - T(y)| \geq [|T(0)|^p + 1]^{1/p}.$$

This inequality, when combined with Lemma 1, gives that

$$1 \geq [|T(0)|^p + 1]^{1/p}$$

and consequently $T(0) = 0$. □

A similar result can be stated in the setting of Hilbert space if we replace condition b) by: $T(0)$ is orthogonal to the set $T(S_X)$. Under this assumption we would obtain that

$$\|T(0)\|_2^2 + \|T(y)\|_2^2 = \|T(0) - T(y)\|_2^2$$

for every $y \in S_X$, from which the same conclusion follows.

Theorem 2. *Let $(X, \|\cdot\|_2)$ be a Hilbert space. Let B_X, S_X be the closed unit ball and the unit sphere of X , respectively, and let $T : B_X \rightarrow X$ be a mapping such that*

- a) $T(S_X) \subset S_X$.
- b) $T(0)$ is orthogonal to the set $T(S_X)$.

If T is nonexpansive with respect to some equivalent norm on X then T has a fixed point. Specifically, $T(0) = 0$.

Suppose now that $T : B \rightarrow B$ is any of the mappings K , G or K_r which we refer to in the introduction. It is clear that T verifies the conditions (a) and (b) of Theorem 2 and that $T(0) \neq 0$. This proves the following corollary.

Corollary 1. *Let $T : B \rightarrow B$ be any of the mappings K , G or K_r and let $\|\cdot\|$ be any norm on ℓ_2 , equivalent to $\|\cdot\|_2$. Then T is not $\|\cdot\|$ -nonexpansive.*

We wish to thank the referee for his valuable comments.

REFERENCES

- [B] Baillon, J.B. *Quelques aspects de la théorie des points fixes dans les espaces de Banach*, I, Séminaire d'Analyse Fonctionnelle de l'Ecole Polytechnique, no. VII, 1978-79.
- [GK] K. Goebel and W.A. Kirk *Topics in metric fixed point theory*. Cambridge Univ. Press 1990. MR **92c**:47070
- [GKT] K. Goebel, W.A. Kirk and R.L. Thele. *Uniformly Lipschitzian semigroups in Hilbert space*. Canad. J. Math. **26** (1974), 1245-1256. MR **50**:10919
- [K] Khamsi, M.A. *Normal structure for Banach spaces with Schauder decomposition*. Canad. Math. Bull. **33** (3), (1989), 1-8. MR **90i**:46032
- [T] D. Tingley *Noncontractive uniformly Lipschitzian semigroups in Hilbert space*, Proc. of Am. Math. Soc. **92**, 3 (1984) (355-361). MR **85k**:47105

(J. García-Falset and E. Lloréns-Fuster) DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICAS, UNIVERSITAT DE VALÈNCIA, 46100 BURJASSOT, VALENCIA, SPAIN

E-mail address: `jesus.garcia@uv.es`

E-mail address: `enrique.llorens@uv.es`

(A. Jiménez-Melado) DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE MÁLAGA, 29071 MÁLAGA, SPAIN

E-mail address: `melado@ccuma.sci.uma.es`