A CONVERSE OF THE GELFAND THEOREM

YIPING MAO

(Communicated by Christopher Croke)

ABSTRACT. In this short note we obtain a converse to the Gelfand theorem: a Riemannian manifold is homogeneous if the isometrically invariant operators on the manifold form a commutative algebra.

1. INTRODUCTION

A remarkable property of a symmetric space is that the isometrically invariant differential operators on it form a commutative algebra. This is known as the Gelfand theorem ([4]). The converse of the theorem is not true in general. There are many non-symmetric homogeneous Riemannian manifolds on which the algebra of isometrically invariant differential operators is commutative ([6], [11]). However, there still are many results relating to the converse of the Gelfand theorem (See [1], [7], [8], etc.). Among them is Kowalski and Vanhecke's theorem which asserts that a homogeneous Kähler manifold is locally Hermitian symmetric if all isometrically invariant differential operators on the manifold commute each other.

In this short note, we prove a converse of the Gelfand theorem:

Theorem. *If the algebra of isometrically invariant differential operators on a complete connected Riemannian manifold M is commutative, M is a Riemannian homogeneous manifold.*

A Riemannian homogeneous manifold with commutative invariant differential operators is called a commutative space ([10]). Our result says that the “homogeneous” assumption can be removed from the above definition. Commutative spaces are contained in the class of Riemannian manifolds with volume-preserving local geodesic symmetries ([9]). It is known that a naturally reductive homogeneous space of dimension ≤ 5 is necessarily a commutative space ([10]). This is no longer true for dimension ≥ 6 ([5]). On the other hand, a commutative space is not necessarily a naturally reductive homogeneous space due to the results of Kaplan and Ricci's on Heisenberg type Lie groups ([6], [11]). It is an interesting problem to characterize commutative spaces.

The author wishes to thank Carolyn Gordon, David Webb and Steve Zelditch for their help and encouragement.

Received by the editors April 4, 1995 and, in revised form, April 30, 1996.

1991 *Mathematics Subject Classification.* Primary 53C25, 53C30.

Key words and phrases. Invariant differential operator, homogeneous manifold.

2. PROOF OF THE THEOREM

In this section, M denotes a complete connected Riemannian manifold. Denote by $C^\infty(M)$ the Fréchet space of smooth functions on M with the topology under which a sequence of functions is convergent if for any integer $k \geq 0$, the sequence of the k -th derivatives of the functions is uniformly convergent in any compact subset of M . As usual, a differential operator on M is a continuous linear operator $A : C^\infty(M) \rightarrow C^\infty(M)$ which has local form $A = \sum_{|\alpha| \leq k} f_\alpha D^\alpha$. A differential operator A is called isometrically invariant if $g^*A = A$ holds for any isometry g of M , where g^*A is defined by $g^*A(u)(x) = A(u \circ g)(g^{-1}x)$. It is easy to see that the Laplacian Δ on M is an isometrically invariant differential operator. To list other examples of the invariant operators, we need the following definition of the spherical mean operator.

The spherical mean operator $L_r : C^\infty(M) \rightarrow C^\infty(M)$ is defined by

$$L_r f(x) = \int_{S_x(M)} f(\exp_x r\xi) d\mu_x(\xi)$$

where $d\mu_x$ stands for the normalized canonical measure on the unit tangent sphere $S_x(M) = \{\xi \in T_x M \mid \|\xi\| = 1\}$. We refer to [3] as a reference for the operator L_r . It is easy to see that the spherical mean operator L_r is an isometrically invariant operator. We notice that L_r is smooth in r . Let Z be the vector field on $S(M)$ generating the geodesic flow G^t . A direct calculation shows that for $k = 1, 2, \dots$,

$$\begin{aligned} \frac{d^k}{dr^k} \Big|_{r=0} L_r f(x) &= \int_{S_x(M)} (Z^k \pi^* f)(\xi) d\mu_x(\xi), \\ Z^k(\pi^* f)(\xi) &= \nabla_{\xi \dots \xi}^k f = \sum_{\alpha_1 \dots \alpha_k} \xi^{\alpha_1} \dots \xi^{\alpha_k} \nabla_{\alpha_k} \dots \nabla_{\alpha_1} f, \end{aligned}$$

where $\pi : S(M) \rightarrow M$ is the natural projection, $\xi = \sum \xi^i \frac{\partial}{\partial x^i} \in S_x(M)$, and ∇ is the covariant differential on M . Applying the above identity to the Taylor expansion of L_r , we get that, for each integer $n \geq 0$,

$$L_r = \sum_{k \leq n} \frac{1}{(2k)!} P_{2k} r^{2k} + \frac{d^{2n+1}}{dr^{2n+1}} \Big|_{r=\bar{r}} (L_r) \frac{r^{2n+1}}{(2n+1)!}$$

where

$$P_{2k} f(x) = \int_{S_x(M)} \sum \xi^{\alpha_1} \dots \xi^{\alpha_{2k}} \nabla_{\alpha_{2k}} \dots \nabla_{\alpha_1} (f)(x) d\mu_x(\xi).$$

Since L_r is an isometrically invariant operator, so is each coefficient P_{2k} .

If the algebra of isometrically invariant differential operators on M is commutative, M has cyclically parallel Ricci tensor by a theorem of Z. Szabo's ([12]). Therefore, M is real analytic (see [2], [12]). In this case, for a local analytic function f ,

$$L_r f(x) = \sum_k \frac{1}{(2k)!} P_{2k}(f)(x) r^{2k}$$

holds for small enough $r > 0$ such that $\exp_x(r\xi)$ is contained in the analytic domain of f . Since analytic functions are locally dense in $C^\infty(M)$, we get the following lemma:

Lemma 1. *If M is a Riemannian manifold with commutative algebra of isometrically invariant differential operators, then the spherical mean operator L_r commutes with all isometrically invariant differential operators.*

In the rest of the paper, we denote by G the isometry group of M . To prove the theorem, we need the following lemma:

Lemma 2. *Let $x \in M$ and $U \subseteq M$ be an open set with compact closure $\bar{U}$. The set $K_x = \{g \in G \mid gx \in \bar{U}\}$ is pre-compact in G with regard to the compact-open topology.*

Proof. Let $\{g_n\}_1^\infty \subseteq K_x$ be a sequence. According to the definition of K_x , we see that $\{g_n x\} \subseteq \bar{U}$. There is a subsequence $\{g_{n_i}\}$ of $\{g_n\}$, due to the compactness of $\bar{U}$, such that $\{g_{n_i}(x)\}$ converges to a point $x_1 \in M$. Let $\{e_1, \dots, e_n\}$ be an orthonormal frame of M at x . Since $(g_{n_i})_*(e_j)$ are contained in a compact subset of the sphere tangent bundle $S(M)$, by passing to a subsequence if necessary, we may assume that $\lim_{i \rightarrow \infty} (g_{n_i})_*(e_j)$ exists for all $j = 1, \dots, n$. Therefore, $\{(g_{n_i})_*(e)\}$ is uniformly convergent in $S_x M$.

We will show that $\{g_{n_i}(p)\}$ is uniformly convergent for p in any compact set of M . To this end, choose a geodesic $\gamma : [0, l] \rightarrow M$ connecting x with p . Since g_{n_i} are isometries, $\varphi_{n_i}(t) = g_{n_i}(\gamma(t))$ are also geodesics with initial data $\varphi_{n_i}(0) = g_{n_i}(x)$, $\varphi'_{n_i}(0) = (g_{n_i})_*\gamma'(0)$. Hence, by the convergence of $g_{n_i}(x)$ and $(g_{n_i})_*\gamma'(0)$ and the continuous dependence of ordinary differential equations on the initial data, we see that $g_{n_i}(p) = \varphi_{n_i}(l)$ is convergent; more precisely, $g_{n_i}(p)$ is uniformly convergent about p in any compact set.

Define $g(p) = \lim_{i \rightarrow \infty} g_{n_i}(p)$. It is easy to see that g is an isometry of M and $\lim g_{n_i} = g$ under the compact-open topology of G . This proves our lemma. $\square$

Proof of Theorem. For each $y \in M$, the orbit $G(y)$ is a smooth submanifold of M . Since $G(y)$ is a closed subset of M (see [13]), it suffices to prove $\dim(G(y)) = \dim(M)$ for each $y \in M$.

Assume that $\dim(G(y)) < \dim(M)$. Take an open neighborhood U of y such that $\text{dist}^2(x, G(y))$ is smooth in U and U has compact closure $\bar{U}$. Let $\lambda(x)$ be a smooth cut-off function of U and define

$$h(x) = \lambda(x) \text{dist}^2(x, G(y)).$$

It is easy to see that $h(x)$ is a global smooth function satisfying $h(x) > 0$ for each $x \in U \setminus G(y)$ and $\text{supp}(h) \subseteq \bar{U}$. Furthermore, we define

$$h_1(x) = \int_G h(gx) d\mu(g).$$

Here $d\mu$ is the Harr measure on G . Since $h(gx) \neq 0$ implies that $g \in K_x = \{g \in G \mid gx \in \bar{U}\}$, according to Lemma 2, we see that the integrand $g \rightarrow h(gx)$ has compact support in G . Therefore, the function h_1 is well-defined. Moreover, h_1 is a non-negative G -invariant function and satisfies $h_1 > 0$ in $U \setminus G(y)$.

Obviously, $h_1 \Delta$ is a G -invariant differential operator. Here Δ is the Laplacian on M . According to Lemma 1, we get $[h_1 \Delta, L_r] = 0$. On the other hand, we still have $[\Delta, L_r] = 0$. Therefore, it holds that $h_1 L_r \Delta = L_r(h_1 \Delta)$. Take $f \in C^\infty(M)$ such that $\Delta f = 1$ in U . Then for $r > 0$ small enough such that $\exp_y(r\xi) \in U$ for

$\xi \in S_y(M)$, we have $h_1(y)L_r(1) = L_r(h_1)$, i.e.,

$$h_1(y) = \int_{S_y(M)} h_1(\exp_y(r\xi)) d\mu(\xi).$$

However $h_1(y) = 0$ and $\int_{S_y(M)} h_1(\exp_y(r\xi)) d\mu(\xi) > 0$. This is a contradiction. $\square$

REFERENCES

1. J. D'Atri, J. Dorfmeister and Y. Zhao, *The isotropy representation for homogeneous Siegel domains*, Pacific J. Math. **120** (1985), 295-326. MR **87a**:32026
2. D. DeTurck and J. Kazdan, *Some regularity theorems in Riemannian geometry*, Ann. Scient. Ec. Norm. Sup. 4⁰ série **14** (1981), 249-260. MR **83f**:53018
3. A. Gray and T. J. Willmore, *Mean-value Theorems for Riemannian Manifolds*, Proc. Royal Soc. Edinburgh **92A** (1982), 343-364. MR **84f**:53038
4. S. Helgason, *Differential geometry and symmetric spaces*, Academic Press, New York, 1962. MR **26**:2986
5. J. Jiménez, *Existence of Hermitian n -symmetric spaces and non-commutative naturally reductive spaces*, Math. Z. **196** (1987), 133-139. MR **89b**:53088a
6. A. Kaplan, *On the geometry of groups of Heisenberg type*, Bull. Lond. Math. Soc. **15** (1983), 35-42. MR **84h**:53063
7. O. Kowalski and L. Vanhecke, *The Gelfand Theorem and Its Converse for Kähler manifolds*, Proc. Amer. Math. Soc. **102** (1988), 150-152. MR **89d**:53096
8. O. Kowalski and L. Vanhecke, *The Gelfand Theorem and Its Converse in Sasakian geometry*, Kodai Math. J. **11** (1988), 70-77. MR **89d**:53094
9. O. Kowalski and L. Vanhecke, *Opérateurs différentiels invariants et symétries géodésiques préservant le volume*, C. R. Acad. Sci., Paris Sér. I **296** (1983), 1001-1003. MR **86f**:53057
10. O. Kowalski and L. Vanhecke, *Classification of five-dimensional naturally reductive spaces*, Math. Proc. Camb. Philos. Soc. **97** (1985), 445-463. MR **86d**:53033
11. F. Ricci, *Commutative algebras of invariant functions on groups of Heisenberg type*, J. London Math. Soc. **32** (1985), 265-271. MR **87c**:22015
12. Z. Szabo, *Spectral theory for operator families on Riemannian manifolds*, Differential geometry: Riemannian geometry (Los Angeles, CA, 1990), 615-665, Proc. Sympos. Pure Math., 54, Part 3, Amer. Math. Soc., Providence, RI, 1993. MR **94k**:58152
13. S. T. Yau, *Remarks on the Groups of Isometries*, Topology **16** (1977), 239-247. MR **56**:6686

DEPARTMENT OF MATHEMATICS, TEXAS TECH UNIVERSITY, LUBBOCK, TEXAS 79409

Current address: Department of Mathematics, University of South Carolina, Columbia, South Carolina 29208

E-mail address: mao@math.sc.edu