THE RANK STABLE TOPOLOGY OF INSTANTONS ON $\overline{\mathbf{CP}}^2$

JIM BRYAN AND MARC SANDERS

(Communicated by Ronald A. Fintushel)

ABSTRACT. Let $\mathcal{M}_k^n$ be the moduli space of based (anti-self-dual) instantons on $\overline{\mathbf{CP}}^2$ of charge k and rank n . There is a natural inclusion $\mathcal{M}_k^n \hookrightarrow \mathcal{M}_k^{n+1}$. We show that the direct limit space $\mathcal{M}_k^\infty$ is homotopy equivalent to $BU(k) \times BU(k)$. Let ℓ_∞ be a line in the complex projective plane and let $\widetilde{\mathbf{CP}}^2$ be the blow-up at a point away from ℓ_∞ . $\mathcal{M}_k^n$ can be alternatively described as the moduli space of rank n holomorphic bundles on $\widetilde{\mathbf{CP}}^2$ with $c_1 = 0$ and $c_2 = k$ and with a fixed holomorphic trivialization on ℓ_∞ .

1. INTRODUCTION

In his 1989 paper [Ta], Taubes studied the stable topology of the based instanton moduli spaces. He showed that if $\mathcal{M}_k^n(X)$ denotes the moduli space of based $SU(n)$ -instantons of charge k on X , then there is a map $\mathcal{M}_k^n(X) \rightarrow \mathcal{M}_{k+1}^n(X)$ and, in the direct limit topology, $\mathcal{M}_\infty^n(X)$ has the homotopy type of $\text{Map}_0(X, BSU(n))$.

There is also a map $\mathcal{M}_k^n(X) \hookrightarrow \mathcal{M}_k^{n+1}(X)$ given by the direct sum of a connection with the trivial connection on a trivial line bundle and one can consider the direct limit $\mathcal{M}_k^\infty(X)$. For the case of $X = S^4$ with the round metric, it was shown by Kirwan and also by Sanders ([Kir],[Sa]) that the direct limit has the homotopy type of $BU(k)$.

In this note we consider the case of $X = \overline{\mathbf{CP}}^2$ where $\overline{\mathbf{CP}}^2$ denotes the complex projective plane with the Fubini-Study metric and the opposite orientation of the one induced by the complex structure. Our result is:

Theorem 1.1. $\mathcal{M}_k^\infty(\overline{\mathbf{CP}}^2)$ has the homotopy type of $BU(k) \times BU(k)$.

The main tool in the proof of the theorem is a construction of the moduli spaces $\mathcal{M}_k^n(\overline{\mathbf{CP}}^2)$ due to King [Ki]. In general, Buchdahl [Bu] has shown that, for appropriate metrics on the N -fold connected sum $\#_N \overline{\mathbf{CP}}^2$, the moduli spaces $\mathcal{M}_k^n(\#_N \overline{\mathbf{CP}}^2)$ are diffeomorphic to certain spaces of equivalence classes of holomorphic bundles on $\mathbf{CP}^2$ blown-up at N points. The universal $U(k) \times U(k)$ bundle that appears giving the homotopy equivalence of Theorem 1.1 can be constructed as higher direct image bundles (see section 3).

Remark 1.1. The cofibration $S^2 \rightarrow \overline{\mathbf{CP}}^2 \rightarrow S^4$ gives rise to the fibration of mapping spaces $\Omega^4 BSU(n) \rightarrow \text{Map}_*(\overline{\mathbf{CP}}^2, BSU(n)) \rightarrow \Omega^2 BSU(n)$ which for K-theoretic

---

Received by the editors August 2, 1996.

1991 *Mathematics Subject Classification.* Primary 58D27, 53C07, 55R45, 14Dxx.

reasons is a trivial fibration in the limit over n . The total space of this fibration is homotopy equivalent to the space of based gauge equivalence classes of all connections on $\overline{\mathbf{CP}}^2$. Thus, from Taubes' result, $\mathcal{M}_k^\infty$ must have the property that taking the limit over k gives $BU \times BU$. For S^4 , similar remarks imply that $\lim_{k \rightarrow \infty} \mathcal{M}_k^\infty(S^4) \simeq BU$ and the inclusion of $\mathcal{M}_k^\infty(S^4)$ into this limit has been shown to be (up to homotopy) the natural inclusion $BU(k) \hookrightarrow BU$ ([Sa]). Theorem 1.1 and these results for S^4 suggest a general conjecture which is supported by the fact that the higher direct image bundle giving our homotopy equivalence generalizes in an appropriate way.

Conjecture 1.1. *For appropriate metrics on $\#_N \overline{\mathbf{CP}}^2$, $\mathcal{M}_k^\infty(\#_N \overline{\mathbf{CP}}^2)$ has the homotopy type of a product $BU(k) \times \cdots \times BU(k)$ with $N + 1$ factors.*

Remark 1.2. Combining Theorem 1.1 with Taubes' stabilization result leads to an alternate proof of Bott periodicity for the unitary group. There is a natural map from instantons on S^4 to those on $\overline{\mathbf{CP}}^2$ (given by pull-back) which in the limit as $n \rightarrow \infty$ is homotopy equivalent to the diagonal $BU(k) \rightarrow BU(k) \times BU(k)$. Taking the limit as $k \rightarrow \infty$ and applying Taubes' result, the diagonal map appears as the inclusion of fibers in the fibration $BU \simeq \Omega_k^4 BU \rightarrow BU \times BU \rightarrow \Omega^2(BSU)$ (see Remark 1.1). However, the map $BU \rightarrow BU \times BU$ has homotopy fiber $U \times U / U \simeq U$, and therefore, given the above fibration we must have $U \simeq \Omega^3 BSU \simeq \Omega^2 SU \simeq \Omega^2 U$.

Tian ([Ti]) noticed that results for limits of instantons on S^4 (see [Kir] or [Sa]) already imply the four-fold periodicity $\Omega^4 BU \simeq Z \times BU$. The $\overline{\mathbf{CP}}^2$ case thus gives the finer two-fold periodicity $\Omega^2 BU \simeq Z \times BU$, as one might expect due to the nontrivial $S^2 \hookrightarrow \overline{\mathbf{CP}}^2$.

In a future paper we will study limits of $Sp(n)$ and $SO(n)$ instantons on $\overline{\mathbf{CP}}^2$ and their relationships to those on S^4 . As an amusing corollary, we will be able to rederive many of the Bott periodicity relationships among Sp , U , SO , and their homogeneous spaces.

2. THE CONSTRUCTION OF $\mathcal{M}_k^n(\overline{\mathbf{CP}}^2)$

Let $x_0 \in \overline{\mathbf{CP}}^2$ be the base point. Since $\overline{\mathbf{CP}}^2 - \{x_0\}$ is conformally equivalent to $\tilde{\mathbf{C}}^2$, the complex plane blown-up at the origin, $\mathcal{M}_k^n(\overline{\mathbf{CP}}^2)$, can be regarded as instantons on $\tilde{\mathbf{C}}^2$ based “at infinity”. Buchdahl [Bu] proved an analogue in this non-compact setting of Donaldson's theorem relating instantons to holomorphic bundles: Let $\tilde{\mathbf{C}}_N^2$ be the complex plane blown-up at N points with a Kähler metric. Then $\tilde{\mathbf{C}}_N^2$ has a “conformal compactification” to $\#_N \overline{\mathbf{CP}}^2$ and a “complex compactification” to $\widetilde{\mathbf{CP}}_N^2$ (the projective plane blown-up at N points). We have added a point x_0 in the former case and a complex projective line ℓ_∞ in the latter.

Define $\mathcal{M}_{\text{alg},k}^n(\widetilde{\mathbf{CP}}_N^2)$ to be the moduli space consisting of pairs $(\mathcal{E}, \tau)$ where $\mathcal{E}$ is a rank n holomorphic bundle on $\widetilde{\mathbf{CP}}_N^2$ with $c_1(\mathcal{E}) = 0$, $c_2(\mathcal{E}) = k$, and where $\tau : \mathcal{E}|_{\ell_\infty} \rightarrow \mathbf{C}^n \otimes \mathcal{O}_{\ell_\infty}$ is a holomorphic trivialization of $\mathcal{E}$ on ℓ_∞ .

There is a natural map $\Phi : \mathcal{M}_k^n(\#_N \overline{\mathbf{CP}}^2) \rightarrow \mathcal{M}_{\text{alg},k}^n(\widetilde{\mathbf{CP}}_N^2)$ defined as follows. Let $p : \widetilde{\mathbf{CP}}_N^2 \rightarrow \#_N \overline{\mathbf{CP}}^2$ be the map that collapses $\ell_\infty \mapsto x_0$. If $[A] \in \mathcal{M}_k^n$

then the $\bar{\partial}$ operator that defines the holomorphic bundle $\mathcal{V} = \Phi(A)$ is taken to be $(d_{p^*(A)})^{(0,1)}$, the anti-holomorphic part of the covariant derivative defined by the pullback of the connection. The anti-self-duality of A implies that the curvature of $p^*(A)$ is a $(1,1)$ -form and so $\bar{\partial}^2 = 0$.

Buchdahl's theorem is then

Theorem 2.1. *The map $\Phi : \mathcal{M}_k^n(\#_N \overline{\mathbf{CP}}^2) \longrightarrow \mathcal{M}_{\text{alg},k}^n(\widetilde{\mathbf{CP}}_N^2)$ is a diffeomorphism.*

The case $N = 1$ was first proved by King [Ki]. We now restrict ourselves to that case and simply write $\mathcal{M}_k^n$ for $\mathcal{M}_k^n(\overline{\mathbf{CP}}^2)$ and $\mathcal{M}_{\text{alg},k}^n(\widetilde{\mathbf{CP}}^2)$.

King constructed $\mathcal{M}_k^n$ explicitly in terms of linear algebra data. We recall his construction. Consider configurations of linear maps:

$$\begin{array}{ccc} W_0 & \xrightleftharpoons[a_2]{a_1} & W_1 \\ & \searrow b & \swarrow c \\ & V_\infty & \end{array}$$

where W_0 , W_1 and V_∞ are complex vector spaces of dimensions k , k , and n respectively.

A configuration (a_1, a_2, b, c, x) is called *integrable* if it satisfies the equation

$$a_1 x a_2 - a_2 x a_1 + b c = 0.$$

A configuration (a_1, a_2, b, c, x) is *non-degenerate* if it satisfies the following conditions:

$$\forall (\lambda_1, \lambda_2), (\mu_1, \mu_2) \in \mathbf{C}^2 \text{ such that } \lambda_1 \mu_1 + \lambda_2 \mu_2 = 0 \text{ and } (\mu_1, \mu_2) \neq (0, 0),$$

$$\begin{aligned} \nexists v \in W_1 \text{ such that } & \begin{cases} x a_1 v = \lambda_1 v, & (\mu_1 a_1 + \mu_2 a_2) v = 0, \\ x a_2 v = \lambda_2 v, & c v = 0, \end{cases} \\ \text{and } \nexists w \in W_0^* \text{ such that } & \begin{cases} x^* a_1^* w = \lambda_1 w, & (\mu_1 a_1^* + \mu_2 a_2^*) w = 0, \\ x^* a_2^* w = \lambda_2 w, & b^* w = 0. \end{cases} \end{aligned}$$

Let A_k^n be the space of all integrable non-degenerate configurations. $G = Gl(W_0) \times Gl(W_1)$ acts canonically on A_k^n . The action is explicitly given by

$$(g_0, g_1) \cdot (a_1, a_2, b, c, x) = (g_0 a_1 g_1^{-1}, g_0 a_2 g_1^{-1}, g_0 b, c g_1^{-1}, g_1 x g_0^{-1}).$$

Theorem 2.2. *The moduli space $\mathcal{M}_k^n$ is isomorphic to A_k^n/G .*

Proof. King uses such configurations to determine monads that in turn determine holomorphic bundles. Configurations in the same G orbit determine the same bundle. For the sake of brevity we refer the reader to [Ki] or [Br] for details. The construction identifies the vector spaces W_0 and W_1 canonically as $H^1(\mathcal{E}(-\ell_\infty))$ and $H^1(\mathcal{E}(-\ell_\infty + E))$ respectively, where $E \subset \widetilde{\mathbf{CP}}^2$ is the exceptional divisor. The vector space V_∞ is identified with the fiber over ℓ_∞ .

3. PROOF OF THEOREM 1.1

We prove the theorem in two steps: We first show that the space of monad data A_k^n forms a principal $G = Gl(k) \times Gl(k)$ bundle over $\mathcal{M}_k^n$. We then show that the induced G -equivariant inclusion $A_k^n \hookrightarrow A_k^{n+2k}$ is null-homotopic so that we can conclude that A_k^∞ is contractible.

Lemma 3.1. *G acts freely on the space of monad data A_k^n .*

Proof. This is essentially proved in [Ki] where it is implicitly shown that the non-degeneracy conditions are precisely the conditions that guarantee freeness. We point out that this also follows more conceptually from the existence of a universal family $\mathbb{E} \rightarrow \mathcal{M}_k^n \times \widetilde{\mathbf{CP}}^2$ and the cohomological interpretation of W_0 and W_1 :

First, the existence of a universal family can be shown via the gauge theoretic construction: Let V be a smooth hermitian vector bundle on $\widetilde{\mathbf{CP}}^2$ with $c_1(V) = 0$ and $c_2(V) = k$. Let $\mathcal{A}_0^{1,1}$ denote unitary connections on V with curvature of pure type $(1, 1)$ and that restrict to the trivial connection on ℓ_∞ and let $\mathcal{G}_0^C$ denote the complex gauge transformations of V that are the identity restricted to ℓ_∞ . Then $\mathcal{M}_k^n = \mathcal{A}_0^{1,1} / \mathcal{G}_0^C$. The quotient

$$(\mathcal{A}_0^{1,1} \times V) / \mathcal{G}_0^C \rightarrow \mathcal{M}_k^n \times \widetilde{\mathbf{CP}}^2$$

will form a universal bundle if the moduli space is smooth and no $\mathcal{E} \in \mathcal{M}_k^n$ has non-trivial automorphisms (cf. [Fr-Mo] Chapt. IV):

Lemma 3.2. *$\mathcal{M}_k^n$ is smooth and any $\mathcal{E} \in \mathcal{M}_k^n$ has no non-trivial automorphisms preserving $\tau : \mathcal{E}|_{\ell_\infty} \rightarrow \mathbf{C}^n \otimes \mathcal{O}_{\ell_\infty}$.*

By Serre duality $H^2(\mathcal{E} \otimes \mathcal{E}^*) = H^0(\mathcal{E} \otimes \mathcal{E}^* \otimes K)^*$. Since $\mathcal{E} \otimes \mathcal{E}^*$ is trivial on ℓ_∞ , it is also trivial on nearby lines. Any section of $\mathcal{E} \otimes \mathcal{E}^* \otimes K$ restricts to a section of $\mathbf{C}^{n^2} \otimes \mathcal{O}_{\ell_\infty}(-3)$ and so must vanish on ℓ_∞ . Likewise, it must vanish on nearby lines and so it is 0 on an open set and must be identically 0. Thus $H^2(\mathcal{E} \otimes \mathcal{E}^*) = 0$ and smoothness follows once we show there are no automorphisms.

Suppose that there exists an automorphism $\phi \in H^0(\mathcal{E} \otimes \mathcal{E}^*)$ such that $\phi \neq 1$ and ϕ preserves τ so that $\phi|_{\ell_\infty} = 1|_{\ell_\infty}$. Then $\phi - 1$ is a non-zero section of $\mathcal{E} \otimes \mathcal{E}^*$ vanishing on ℓ_∞ . We then get an injection $0 \rightarrow \mathcal{O}(\ell_\infty) \rightarrow \mathcal{E} \otimes \mathcal{E}^*$. Restricting this sequence to ℓ_∞ we get an injection $0 \rightarrow \mathcal{O}_{\ell_\infty}(1) \rightarrow \mathcal{O}_{\ell_\infty} \otimes \mathbf{C}^{n^2}$ which is a contradiction.

Let $\pi : \mathcal{M}_k^n \times \widetilde{\mathbf{CP}}^2 \rightarrow \mathcal{M}_k^n$. The higher direct image sheaves $R^1\pi_*(\mathbb{E}(-\ell_\infty))$ and $R^1\pi_*(\mathbb{E}(-\ell_\infty + E))$ are locally free and rank k . This follows from the index theorem and the vanishing of the H^0 and H^2 cohomology of $\mathcal{E}(-\ell_\infty)$ and $\mathcal{E}(-\ell_\infty + E)$. The H^0 vanishing follows by again considering the restriction of a section of the bundles to lines nearby to ℓ_∞ . Using Serre duality and the same argument, one gets the vanishing for H^2 .

Consequently the vector spaces W_0 and W_1 are the fibers of the vector bundles $R^1\pi_*(\mathbb{E}(-\ell_\infty))$ and $R^1\pi_*(\mathbb{E}(-\ell_\infty + E))$. The G -orbit of a configuration giving a bundle $\mathcal{E}$ can be identified with the group of isomorphisms $g_0 : H^1(\mathcal{E}(-\ell_\infty)) \rightarrow \mathbf{C}^k$ and $g_1 : H^1(\mathcal{E}(-\ell_\infty + E)) \rightarrow \mathbf{C}^k$. Thus A_k^n is realized precisely as the total space of

the principal $Gl(k) \times Gl(k)$ bundle associated to

$$R^1\pi_*(\mathbb{E}(-\ell_\infty)) \oplus R^1\pi_*(\mathbb{E}(-\ell_\infty + E)).$$

Recall that the map $\mathcal{M}_k^n \hookrightarrow \mathcal{M}_k^{n+1}$ is defined by the direct sum with the trivial connection: $[A] \mapsto [A \oplus \theta]$. In terms of holomorphic bundles this is $\mathcal{E} \mapsto \mathcal{E} \oplus \mathcal{O}$. Tracing through the monad construction, it is easy to see that the inclusion induces the G -equivariant map $A_k^n \hookrightarrow A_k^{n+1}$ given by $(a_1, a_2, x, b, c) \mapsto (a_1, a_2, x, b', c')$ where b' is b with an extra first column of zeroes and c' is c with an extra first row of zeroes. Define A_k^∞ to be the direct limit $\lim_{n \rightarrow \infty} A_k^n$ so that there is a homeomorphism between $\mathcal{M}_k^\infty$ and A_k^∞/G .

Lemma 3.3. A_k^∞ is a contractible space.

Proof. Since the A_k^n 's are algebraic varieties and the maps $A_k^n \rightarrow A_k^{n+1}$ are algebraic, they admit triangulations compatible with the maps. Thus A_k^∞ inherits the structure of a CW-complex and so it is sufficient to show that all of its homotopy groups are zero. To this end we prove that for any k and l there is an $r > l$ such that the natural inclusion from $A_k^n \hookrightarrow A_k^r$ is homotopically trivial.

Consider the homotopy $H_t : A_k^n \rightarrow A_k^{2k+n}$ defined as follows:

$$H_t((a_1, a_2, x, b, c)) = ((1-t)a_1, (1-t)a_2, (1-t)x, b_t, c_t)$$

where

$$c_t = \begin{pmatrix} tI_k \\ 0_{k,k} \\ (1-t)c \end{pmatrix}, \quad b_t = (0_{k,k}, tI_k, (1-t)^2b),$$

I_k is the $k \times k$ identity matrix and $0_{k,k}$ is the $k \times k$ zero matrix. To see that $H_t(v) \in A_k^{n+2k}$ for any $v \in A_k^n$, we check that the integrability and non-degeneracy conditions are satisfied for all $0 \leq t \leq 1$. Integrability holds because $b_t c_t = (1-t)^3 bc$. Non-degeneracy is satisfied for all $t \neq 0$ because there is a full rank $k \times k$ block, tI_k , in both c_t and b_t . Furthermore, H_0 is just the inclusion $A_k^n \hookrightarrow A_k^{n+2k}$, so non-degeneracy also holds when $t = 0$. Finally, note that H_1 is a constant map.

These lemmas show that A_k^∞ is a contractible space acted on freely by $G = Gl(k) \times Gl(k)$ and $A_k^\infty/G = \mathcal{M}_k^\infty$. Thus $\mathcal{M}_k^\infty$ is homotopic to BG which in turn has the homotopy type of $BU(k) \times BU(k)$. We end by remarking that the proof shows that the universal $U(k) \times U(k)$ bundle is the bundle that restricts to any of the finite $\mathcal{M}_k^n$'s as $R^1\pi_*(\mathbb{E}(-\ell_\infty)) \oplus R^1\pi_*(\mathbb{E}(-\ell_\infty + E))$.

REFERENCES

- [Br] Bryan, J. *Symplectic Geometry and the Relative Donaldson Invariants of $\overline{\mathbb{CP}}^2$* , to appear in Forum Math.
- [Bu] Buchdahl, N. *Instantons on $n\mathbb{CP}^2$* . J. Diff. Geo. 37(1993), 669–687. MR **94e**:53025
- [Fr-Mo] Robert Friedman and John Morgan. *The Differential Topology of Complex Surfaces*. Springer-Verlag, 1994.
- [Ki] King, A. *Instantons and Holomorphic Bundles on the Blown Up Plane*, Ph.D. Thesis, Oxford, 1989.
- [Kir] Kirwan, F. *Geometric invariant theory and the Atiyah-Jones conjecture*, Proceedings of the Sophus Lie Memorial Conference (Oslo, 1992), editors O. A. Laudal and B. Jahren, Scandinavian University Press, 1994, 161–188.
- [Sa] Sanders, M., *Classifying spaces and dirac operators coupled to instantons*, Trans. of the A.M.S. Vol. 347, No. 10 **1995**. MR **96m**:58030

- [Ta] Taubes, C. *The stable topology of self-dual moduli spaces* J. Diff. Geo. 29 (**1989**), 163–230.
MR **90f**:58023
- [Ti] Tian, Y. *The Atiyah-Jones Conjecture for classical groups and Bott periodicity*, to appear
in J. Diff. Geo.

MATHEMATICAL SCIENCES RESEARCH INSTITUTE, 1000 CENTENNIAL DRIVE, BERKELEY, CALIFORNIA 94720-5070

E-mail address: `jbryan@msri.org`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, DICKINSON COLLEGE, CARLISLE, PENNSYLVANIA 17013

E-mail address: `sandersm@dickinson.edu`