MICROLOCAL ANALYSIS OF ULTRADISTRIBUTIONS

STEVAN PILIPOVIĆ

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. The ultradistributional wave front sets of an ultradistribution u are characterized by the behaviour of $K*u$ on the boundary of the tube domain $D\mathbf{R}^n$, where K is the kernel analysed by Hörmander.

1. INTRODUCTION

We follow the results and ideas of Chapter 8 in [2] and characterize the ultradistributional wave front sets of type WF^* and WF_* for a tempered ultradistribution u by using its convolution with the kernel K (cf. [2]). We give the assertions which are analogous to the corresponding ones for distributions.

The paper is organized as follows. In Section 2 we recall the definitions of ultradistributional wave front sets WF^* and WF_* . These notions are introduced by Eida [1] and Komatsu [4] and called singular spectrums SS^* and SS_* . In Section 3 we recall the definition of tempered ultradistribution spaces $\mathcal{S}'^*$ (cf. [5], [7]) and a theorem from [8] which is needed for later use. In Section 4 we give the microlocal analysis of a $u \in \mathcal{S}'^*$ (Theorems 2, 3 and Corollary 1). Necessary and sufficient conditions for an analytic function in the tube domain $D\mathbf{R}^n$ which determine its boundary value in $\mathcal{S}'^*$ are given in Theorem 2.

2. NOTATION AND NOTIONS

By M_p , $p \in \mathbf{N}_0$, we denote a sequence of positive numbers with $M_0 = 1$. We refer to [3] and [6] for the meaning of conditions (M.1), (M.2)', (M.2), (M.3)' and (M.3). We also use the following condition ([6]):

$$(M.1)^* \quad M_{p-1}^* M_{p+1}^* \leq M_p^*, p \in \mathbf{N}, \text{ where } M_0^* = 1, M_p^* = M_p/p!, p \in \mathbf{N}.$$

Let M_p satisfy (M.1) and (M.3)'. The associated function $M(\rho)$ and the growth function $\tilde{M}(\rho)$ related to M_p are defined by

$$M(\rho) = \sup_{p \in \mathbf{N}_0} \ln \frac{\rho^p}{M_p}, \quad \tilde{M}(\rho) = \sup_{p \in \mathbf{N}_0} \ln \frac{\rho^p}{M_p^*}, \quad \rho > 0.$$

Note, for a given $L > 0$ there is $L_1 > 0$ such that ([8])

$$(1) \quad M(L|\xi|) - |\eta||\xi| \leq \tilde{M}(L_1/|\eta|), \quad \xi, \eta \in \mathbf{R}^n.$$

Received by the editors February 27, 1996.

1991 *Mathematics Subject Classification*. Primary 32A40, 46F15; Secondary 58G07.

Key words and phrases. Ultradistributions, wave front.

We denote by Ω an open set in $\mathbf{R}^n$, and $K \subset\subset \Omega$ means that K is a compact subset of Ω . Recall,

$$\|\varphi\|_{K,h,M_p} = \sup_{x \in K, \alpha \in \mathbf{N}_0^n} \frac{|\varphi^{(\alpha)}(x)|}{h^{|\alpha|} M_{|\alpha|}}, \varphi \in C^\infty(\Omega).$$

The symbol $*$ is used for both (M_p) and $\{M_p\}$. We refer to [3] for the definitions of $\mathcal{D}'^*(\Omega)$, $\mathcal{D}_K^*(\Omega)$ and $\mathcal{D}'_K^*(\Omega)$. Throughout the paper we will assume that (M.1), (M.2)' and (M.3)' hold. Eida [1] and Komatsu [4] have defined SS_* - and SS^* -singular support of a hyperfunction f . We will recall the definitions related to ultradistributions and call them wave front sets.

Let N_p be a sequence of positive numbers which satisfies (M.1), (M.2)', (M.3)' and $N_0 = 1$. Then ([3]):

$$(N_p) \leq (M_p) \quad (\text{resp., } \{N_p\} \leq \{M_p\})$$

if there are constants $L > 0$ and $C > 0$ (resp., for every $\epsilon > 0$ there is $C_\epsilon > 0$) such that

$$N_p \leq CL^p M_p \quad (\text{resp., } N_p \leq C_\epsilon \epsilon^p M_p), \quad p \in \mathbf{N}_0.$$

Let $f \in \mathcal{D}'^\dagger$, where $\dagger = N_p \leq * = M_p$. Then $(x, \omega) \in S^* \Omega = \Omega \times S^{n-1}$ is not in $WF_* f$ (resp., not in $WF^* f$) iff there exist a neighborhood $U \subset \Omega$ of x and a conic neighborhood Γ of ω of the form $\Gamma = \{\xi \neq 0; |\xi/|\xi| - \omega| < \eta\}$ such that for every $\phi \in \mathcal{D}^*(U)$ the following hold:

In the (M_p) case, for every $\epsilon > 0$ there is $C_\epsilon > 0$ such that $|\widehat{\phi f}(\xi)| \leq C_\epsilon e^{-M(\epsilon|\xi|)}$, $\xi \in \Gamma$ (resp., there are $k > 0$ and $C > 0$ such that $|\widehat{\phi f}(\xi)| \leq C e^{M(k|\xi|)}$, $\xi \in \Gamma$).

In the $\{M_p\}$ case, there exist $k > 0$ and $C > 0$ such that $|\widehat{\phi f}(\xi)| \leq C e^{-M(k|\xi|)}$, $\xi \in \Gamma$ (resp., for every $\epsilon > 0$, there is $C_\epsilon > 0$ such that $|\widehat{\phi f}(\xi)| \leq C_\epsilon e^{M(\epsilon|\xi|)}$, $\xi \in \Gamma$). Note, the notion $WF_{\{M_p\}}$ is equal to Hörmander's notion WF_L .

The definition of the singular spectrum SSf , where $f \in \mathcal{B}(\Omega)$, is given by Sato (cf. [10]). For an $f \in \mathcal{D}'^*(\Omega)$, $(x, \omega) \in S^* \Omega$ is not in SSf if this point is not in $SS\{f\}$, where $\{f\}$ denotes the corresponding hyperfunction. This notion is equal to Hörmander's $WF_A f$, the analytic wave front set of f ([2], Definition 9.3.2, Theorem 9.6.3) and we will use this notation.

We also use the definition according to which $(x, \xi) \in \Omega \times (\mathbf{R}^n \setminus \{0\})$ is an element of the corresponding singular spectrum defined above if this holds for $(x, \xi/|\xi|)$.

Let $D\mathbf{R}^n = \{z \in \mathbf{C}^n; |\text{Im } z| < 1\}$ and $S^*\mathbf{R}^n = \partial D\mathbf{R}^n$. Recall ([4]), $O^*|_{D\mathbf{R}^n}$ (resp., $O_*|_{D\mathbf{R}^n}$) is a sheaf over $\mathbf{C}^n$ of holomorphic functions in $D\mathbf{R}^n$ which satisfy the following growth condition near $S^*\mathbf{R}^n$:

Let U be an open set in $\mathbf{C}^n$. Then a function $F(z)$ is in $O^*|_{D\mathbf{R}^n}(U)$ (resp., $O_*|_{D\mathbf{R}^n}(U)$) if F is holomorphic in $D\mathbf{R}^n \cap U$ such that for every compact set $K \subset\subset U$,

in the (M_p) case, there are $C > 0$ and $k > 0$,

in the $\{M_p\}$ case, for every $k > 0$ there is $C > 0$ such that

$$|F(z)| \leq C e^{\tilde{M}(\frac{k}{1-|y|})}, \quad z = x + \sqrt{-1}y \in K \cap D\mathbf{R}^n$$

(resp., for every compact set $K \subset\subset U$,

in the (M_p) case, for every ultradifferential operator $P(\partial)$ of class (M_p) ,

in the $\{M_p\}$ case, for every ultradifferential operator $P(\partial)$ of class $\{M_p\}$

$$P(\partial_z)F \quad \text{is bounded in } K \cap D\mathbf{R}^n.)$$

3. TEMPERED ULTRADISTRIBUTIONS

We will recall the definitions and the basic structural properties of tempered ultradistributions (cf. [5] and [7]). Let $m > 0$. The space of smooth functions φ on $\mathbf{R}^n$ which satisfy

$$\sigma_{m,2}(\varphi) = \left(\sum_{\alpha, \beta \in \mathbf{N}_0^n} \int_{\mathbf{R}^n} \left| \frac{m^{|\alpha+\beta|}}{M_{|\alpha|} M_{|\beta|}} (1 + |x|^2)^{|\beta|/2} \varphi^{(\alpha)}(x) \right|^2 dx \right)^{1/2} < \infty,$$

equipped with the topology induced by the norm $\sigma_{m,2}$, is denoted by $\mathcal{S}_2^{M_p, m}$. The strong duals of $\mathcal{S}^{(M_p)} = \text{proj} \lim_{m \rightarrow \infty} \mathcal{S}_2^{M_p, m}$ and $\mathcal{S}^{\{M_p\}} = \text{ind} \lim_{m \rightarrow 0} \mathcal{S}_2^{M_p, m}$ are called spaces of tempered ultradistributions of Beurling and Roumieu type. For every fixed $p \in [1, \infty]$, the family of norms $\{\sigma_{m,2}; m > 0\}$ is equivalent to the family of norms $\{\sigma_{m,p}; m > 0\}$ where instead of the L^2 norm we put the L^p norm. In fact, in the sequel we will use the family of norms

$$s_h(\phi) = \sup \left\{ \frac{h^{|\alpha|}}{M_{|\alpha|}} \mid \phi^{(\alpha)}(x) \mid e^{M(h|x|)}; \alpha \in \mathbf{N}_0^n, x \in \mathbf{R}^n \right\}, \quad h > 0,$$

which is equivalent to $\{\sigma_{h,2}; h > 0\}$.

$\mathcal{S}^{(M_p)}$ and $\mathcal{S}^{\{M_p\}}$ are (FS) - and (LS) -spaces respectively. If (M.2) holds, they are (FN) - and (LN) -spaces respectively and $\mathcal{D}^* \hookrightarrow \mathcal{S}^* \hookrightarrow \mathcal{E}^*$, $\mathcal{S}^* \hookrightarrow \mathcal{S}$, where “ $A \hookrightarrow B$ ” means that A is dense in B and the inclusion mapping is continuous.

An $f \in \mathcal{D}'^*$ is in $\mathcal{S}'^*$ if and only if there exists a family $F_{\alpha,\beta}$, $\alpha, \beta \in \mathbf{N}_0^n$, in $L^2(\mathbf{R}^n)$ such that

$$f = \sum_{\alpha, \beta \in \mathbf{N}_0^n} ((1 + |x|^2)^{\beta/2} F_{\alpha,\beta})^{(\alpha)} \quad \text{in } \mathcal{S}'^*,$$

and in the case $\mathcal{S}'^{(M_p)}$, there exists $k > 0$ (resp., in the case $\mathcal{S}'^{\{M_p\}}$, for every $k > 0$) such that

$$\left(\sum_{\alpha, \beta \in \mathbf{N}_0^n} \int_{\mathbf{R}^n} \left| \frac{M_{|\alpha|} M_{|\beta|}}{k^{|\alpha+\beta|}} F_{\alpha,\beta}(x) \right|^2 dx \right)^{1/2} < \infty.$$

The Fourier transformation is an isomorphism of $\mathcal{S}^*$ onto itself.

We recall the next theorem for later use.

Theorem 1 ([8]). *Assume (M.1)*, (M.2) and (M.3)' hold for $\dagger = N_p$ ($N_0 = 1$). Let Γ be an open convex cone in $\mathbf{R}^n$ and F be an analytic function in*

$$Z = \{z \in \mathbf{C}^n; \text{Im } z \in \Gamma, |\text{Im } z| < \gamma\}$$

for some $\gamma > 0$. Moreover, assume

$$|F(x + \sqrt{-1}y)| \leq C_{a,b} e^{N(a|x|) + \tilde{N}(\frac{b}{|y|})}, \quad x + \sqrt{-1}y \in Z,$$

in the (N_p) case for some $a > 0$, $b > 0$, and $C_{a,b} > 0$, and in the $\{N_p\}$ case for every $a > 0$, $b > 0$ there exists $C_{a,b} > 0$. Then

$$F(x + \sqrt{-1}y) \xrightarrow{\mathcal{S}'^\dagger} F(x + \sqrt{-1}0), \quad y \rightarrow 0, \quad y \in \Gamma.$$

Note ([8]),

$$(2) \quad \langle F(x + \sqrt{-1}0), \varphi(x) \rangle = \int_{\mathbf{R}^n} F(x + \sqrt{-1}Y) \Phi(x + \sqrt{-1}Y) dx \\ + 2\sqrt{-1} \sum_{i=1}^n Y_i \int_0^1 \int_{\mathbf{R}^n} \frac{\partial}{\partial \bar{z}_i} \Phi(x + \sqrt{-1}tY) F(x + \sqrt{-1}Yt) dt dx.$$

Moreover, there exists $C > 0$ such that

$$(3) \quad |\langle F(x + \sqrt{-1}0), \varphi(x) \rangle| \leq C s_h(\varphi), \quad \varphi \in \mathcal{S}^\dagger,$$

where $\Phi(z)$ is the almost analytic extension of φ (cf. [6] or [8]).

4. MICROLOCAL ANALYSIS OF ULTRADISTRIBUTIONS

As in [2], put

$$I(\xi) = \int_{|\omega|=1} e^{-\langle \omega, \xi \rangle} d\omega, \quad \xi \in \mathbf{R}^n, \quad K(z) = (2\pi)^{-n} \int \frac{e^{\sqrt{-1}\langle z, \xi \rangle}}{I(\xi)} d\xi, \quad z \in D\mathbf{R}^n.$$

We call K Hörmander's kernel. Recall ([2], Lemma 8.4.10) that $K(z)$ is an analytic function in $\tilde{\Omega} = \{z \in \mathbf{C}^n; \langle z, z \rangle \notin (-\infty, -1]\}$. The properties of this function given in the quoted lemma imply

$$|K(x + \sqrt{-1}y)| \leq \frac{Cn!e^{-c|x|}}{(1-|y|)^n}, \quad z \in D\mathbf{R}^n,$$

for some $C > 0$ and $c > 0$. By using the Cauchy formula on the contour

$$\Gamma_{t+\sqrt{-1}y} = \Gamma_{t_1+\sqrt{-1}y_1} \times \cdots \times \Gamma_{t_n+\sqrt{-1}y_n},$$

where $\Gamma_{t_j+\sqrt{-1}y_j}$ is the boundary of

$$\{(\tau_j, u_j); |t_j - \tau_j| < 1, |y_j - u_j| < (1-|y|)/2\},$$

it follows that for some $C > 0$ and $c > 0$

$$|\frac{\partial^\alpha}{\partial t^\alpha} K(t + \sqrt{-1}y)| < C \frac{\alpha!n!}{(1-|y|)^{|\alpha|+n}} e^{-c|t|}, \quad t \in \mathbf{R}^n, |y| < 1, \quad \alpha \in \mathbf{N}_0^n.$$

This implies that $K(\cdot + \sqrt{-1}y) \in \mathcal{S}^*$ for every fixed y , $|y| < 1$.

The next theorem is proved in [2] for tempered distributions.

Theorem 2. Let $\dagger = N_p$ and $*$ = M_p satisfy (M.1)*, (M.2) and (M.3)' and let $N_p \leq M_p$. Let $u \in \mathcal{S}'^\dagger$ and $U(z) = (u * K)(z) = \langle u(t), K(x - t + \sqrt{-1}y) \rangle$, $z \in D\mathbf{R}^n$.

a) Then U is analytic in $D\mathbf{R}^n$ and

in the (N_p) case, for some $a > 0$, $b > 0$ there is $C_{a,b} > 0$,

in the $\{N_p\}$ case, for every $a > 0$, $b > 0$ there is $C_{a,b} > 0$, such that

$$(4) \quad |U(z)| \leq C_{a,b} e^{N(a|x|) + \tilde{N}(b/(1-|y|))}, \quad z = x + \sqrt{-1}y \in D\mathbf{R}^n.$$

For every $\omega \in S^{n-1}$ there exists the limit

$$\lim_{\substack{y \rightarrow \omega \\ z \in D\mathbf{R}^n}} \langle U(x + \sqrt{-1}y), \phi(x) \rangle = \langle U(x + \sqrt{-1}\omega), \phi(x) \rangle, \quad \phi \in \mathcal{S}^\dagger,$$

and the mapping $S^{n-1} \ni \omega \mapsto U(\cdot + \sqrt{-1}\omega) \in \mathcal{S}'^\dagger$ is continuous. Moreover,

$$(5) \quad \langle u, \phi \rangle = \int_{\mathbf{R}^n} \langle U(\cdot + \sqrt{-1}\omega), \phi \rangle d\omega, \quad \phi \in \mathcal{S}^\dagger.$$

b) Conversely, if U is given satisfying (4), then (5) defines an ultradistribution in $\mathcal{S}'^\dagger$.

c1) $q = (x_0, \omega_0) \notin WF^*u$ if and only if U is $\mathcal{O}^*$ in a neighborhood of $x_0 - \sqrt{-1}\omega_0$.

c2) $q \notin WF_*u$ if and only if U is $\mathcal{O}_*$ in a neighborhood of $x_0 - \sqrt{-1}\omega_0$.

c3) $q \notin WFAu$ if and only if U is analytic at $x_0 - \sqrt{-1}\omega_0$ (i.e. in a neighborhood of this point).

Proof. a) We will give only the proof in the (M_p) case. There exist $h > 0$, $C_h > 0$ and suitable constants which do not depend on h , such that

$$\begin{aligned} |\langle u(t), K(z-t) \rangle| &\leq C_h \sup_{\substack{\alpha \in \mathbf{N}_0^n \\ t \in \mathbf{R}^n}} \left\{ \frac{h^\alpha |\frac{\partial^\alpha}{\partial t^\alpha} K(z-t)| e^{N(h|t|)}}{N_{|\alpha|}} \right\} \\ &\leq CC_h \sup_{\substack{\alpha \in \mathbf{N}_0^n \\ t \in \mathbf{R}^n}} \left\{ \frac{h^\alpha \alpha! n! e^{-c|x-t|}}{N_{|\alpha|} (1-|y|)^{|\alpha|+n}} e^{N(h|t|)} \right\} \\ &\leq C_1 C_h e^{N(h|x|)} \sup_{\alpha \in \mathbf{N}_0^n} \frac{h^\alpha \alpha!}{(1-|y|)^{|\alpha|} N_{|\alpha|}} \\ &\leq C_1 C_h e^{N(h|x|) + \tilde{N}(\frac{h}{1-|y|})}, \quad z \in D\mathbf{R}^n. \end{aligned}$$

This proves (4). Since the Fourier transformation of $U(\cdot + \sqrt{-1}y)$, $|y| < 1$, is $e^{-\langle y, \xi \rangle} \hat{u}(\xi)/I(\xi)$, $\xi \in \mathbf{R}^n$, and

$$(6) \quad I(\xi) = (2\pi)^{(n-1)/2} \frac{e^{|\xi|}}{|\xi|^{(n-1)/2}} (1 + \mathcal{O}(1/|\xi|)), \quad |\xi| \rightarrow \infty$$

([2], p. 287), it follows that $y \mapsto U(\cdot + \sqrt{-1}y)$, $|y| \leq 1$, is a continuous function $\{y; |y| \leq 1\} \rightarrow \mathcal{S}'^{(N_p)}$.

Let us prove this. For any $\phi \in \mathcal{S}'$, $\xi \in \mathbf{R}^n$, $\alpha \in \mathbf{N}_0$ and $|y| \leq 1$ we have

$$\begin{aligned} &| (e^{-\langle y, \xi \rangle} (1/I(\xi)) \phi(\xi))^{(\alpha)} | \\ &= | \sum_{i \leq \alpha} \binom{\alpha}{i} (y)^{\alpha-i} e^{-\langle y, \xi \rangle} \sum_{p \leq i} \binom{i}{p} \phi^{(i-p)}(\xi) (1/I(\xi))^{(p)} |. \end{aligned}$$

Since $|(I(\xi))^{(\beta)}| \leq I(\xi)$, $\xi \in \mathbf{R}^n$, $\beta \in \mathbf{N}_0^n$, we conclude

$$|(1/I(\xi))^{(p)}| \leq 2^{|p|} p! / I(\xi), \quad \xi \in \mathbf{R}^n.$$

Now by (6),

$$|(e^{-\langle y, \xi \rangle} (1/I(\xi)) \phi(\xi))^{(\alpha)}| \leq C 3^{|\alpha|} (1 + |\xi|)^{(n-1)/2} \sup_{|i| \leq \alpha} |\phi^{(i)}(\xi)|$$

which implies:

$$\begin{aligned} \langle u, \phi \rangle &= \lim_{r \rightarrow 1} \left\langle \int_{|\omega|=1} \frac{e^{-\langle r\omega, \xi \rangle} \hat{u}(\xi)}{I(\xi)} d\omega, \hat{\phi}(\xi) \right\rangle \\ &= \lim_{r \rightarrow 1} \int_{|\omega|=1} \left\langle \frac{e^{-r\langle \omega, \xi \rangle} \hat{u}(\xi)}{I(\xi)}, \hat{\phi}(\xi) \right\rangle d\omega = \int_{|\omega|=1} \langle U(x + \sqrt{-1}\omega), \phi(x) \rangle d\omega, \end{aligned}$$

where $y = r\omega$, $\omega \in S^{n-1}$.

b) Conversely, if U is analytic in $D\mathbf{R}^n$ and satisfies (4), then Theorem 1, more precisely (2) and (3), imply that

$$\lim_{\substack{y \rightarrow \omega \\ z \in D\mathbf{R}^n}} U_y = U_\omega, \quad |(\omega_1, \dots, \omega_n)| = 1,$$

exists in $\mathcal{S}'^{(N_p)}$, where $U_y = U(\cdot + \sqrt{-1}y)$, $|y| < 1$ and

$$\begin{aligned} \langle U_\omega, \phi \rangle &= \int U(x) \Phi(x - \sqrt{-1}\omega) dx \\ &\quad - 2\sqrt{-1} \sum \omega_i \int_0^1 \int_{\mathbf{R}^n} U(x + \sqrt{-1}(1-t)\omega) \frac{\partial \Phi}{\partial \bar{z}_1}(x + \sqrt{-1}(1-t)\omega) dt dx. \end{aligned}$$

In fact, we have to put $U(z + \sqrt{-1}\omega) = F(z)$, $Y = -\omega$, and the conclusion follows as in Theorem 1.

Moreover, it follows that $S^{n-1} \ni \omega \mapsto U_\omega \in \mathcal{S}'^{(N_p)}$ is continuous, which implies that by

$$\phi \mapsto \langle u, \phi \rangle = \int \langle U_\omega(x), \phi(x) \rangle d\omega, \quad \phi \in \mathcal{S}'^{(M_p)},$$

an element from $\mathcal{S}'^{(N_p)}$ is defined. One can easily show that $U = u * K$.

c) We will prove the “if” parts.

Suppose $q \notin WF^*u$. Let $\varphi \in \mathcal{D}^*(U)$, where $U = \{x; |x - x_0| < 2r\}$ and $0 \leq \varphi \leq 1$, $\varphi \equiv 1$ on $\{x; |x - x_0| < r\}$.

Note that

$$(7) \quad |\widehat{\varphi u}(\xi)| < C_1 e^{M(L_1|\xi|)}, \quad \xi \in \mathbf{R}^n,$$

for some $L_1 > 0$ and some $C_1 > 0$ in the (M_p) case (resp., for every $L_1 > 0$ there is $C_1 > 0$ in the $\{M_p\}$ case). Put $U = \varphi u * K + (1 - \varphi)u * K = U_1 + U_2$. This decomposition is also used in parts b) and c). Let $x \in \{x; |x - x_0| < r/2\}$. Then,

$$(1 - \varphi(t))K(x + \sqrt{-1}y - t) = 0, \quad |t - x_0| < r, \quad x + \sqrt{-1}y \in \mathbf{C}^n.$$

If $|t - x_0| \geq r$, then $|x - t| \geq r/2$ and for $|y| < 1 + r/2$,

$$t \mapsto K(x + \sqrt{-1}y - t)$$

is a function which belongs to S^* .

This implies that

$$U_2(z) = \langle u(t), (1 - \varphi(t)), K(z - t) \rangle, \quad z \in D\mathbf{R}^n,$$

is analytic in

$$\{x + \sqrt{-1}y; |y| < 1 + r/2, |x - x_0| < r/2\},$$

which is a neighborhood of $x_0 - \sqrt{-1}\omega_0$. So in the proofs of c1), c2) and c3) below we have to consider U_1 .

c1) Let γ be an open convex cone which contains ω_0 , $\epsilon > 0$ and $|y + \omega_0| < \epsilon$, $|y| < 1$. We shall use the following inequalities:

$$(8) \quad M(L|\xi|) - \langle \xi, y \rangle - |\xi| \leq \tilde{M}(\tilde{L}/(1 - |y|)), \quad \xi \in \gamma,$$

which holds for some $\tilde{L} > 0$ (see (1)), and

$$(9) \quad -\langle y, \xi \rangle - |\xi| \leq -\epsilon|\xi|, \quad \xi \notin \gamma,$$

which holds if ϵ is small enough.

Let $|y + \omega_0| < \epsilon$, $|y| < 1$, $|x - x_0| < r$. Then $(z = x + \sqrt{-1}y)$,

$$\begin{aligned} |U_1(z)| &= |\mathcal{F}(\varphi u * K(z, \cdot))(\xi)| \\ &\leq \int_{\mathbf{R}^n} \frac{e^{-\langle y, \xi \rangle}}{I(\xi)} \widehat{\varphi u}(\xi) d\xi = \int_{\xi \in \gamma} + \int_{\xi \notin \gamma}. \end{aligned}$$

By using (7) and (8) for the first integral and (7) and (9) for the second integral, we obtain that U_1 is $\mathcal{O}^*$ in some neighborhood of $x_0 - \sqrt{-1}\omega_0$.

c2) Let $q \notin WF_* u$. By using

$$|P(\partial_z)(u\varphi * K)(z)| \leq \int_{\mathbf{R}^n} \frac{e^{-\langle y, \xi \rangle}}{I(\xi)} P(\sqrt{-1}\xi) \widehat{u\varphi}(\xi) d\xi,$$

in a similar way we obtain that for every ultradifferential operator $P(\partial_z)$ of the corresponding class, $P(\partial_z)U$ is bounded in a neighborhood of $x_0 - \sqrt{-1}\omega_0$.

c3) If $(x_0, \omega_0) \notin WF_A u$, then $(x_0, \omega_0) \notin WF_A u\phi$. By [2], Definition 9.3.2, it is equivalent to the analyticity of $u\varphi * K$ at $x_0 - \sqrt{-1}\omega_0$.

The converse parts in c1), c2) and c3) follow from the next lemma (see [2], Lemma 8.4, for tempered distributions).

Lemma 1. *Let $d\mu$ be a measure on S^{n-1} and Γ an open convex cone in $\mathbf{R}^n$ such that*

$$\langle y, \omega \rangle < 0 \quad \text{if } 0 \neq y \in \bar{\Gamma}, \quad \omega \in \text{supp } d\mu.$$

Let U be analytic in $D\mathbf{R}^n$, and satisfy (4) in $D\mathbf{R}^n$. Then

$$F(z) = \int_{S^{n-1}} U(z + \sqrt{-1}\omega) d\mu(\omega), \quad \text{Im } z \in \Gamma, |\text{Im } z| < \gamma,$$

is analytic and

$$(10) \quad |F(z)| \leq C_{a,b} e^{N(a|x|) + \tilde{N}(b/(1-|y|))}, \quad \text{Im } z \in \Gamma, |\text{Im } z| < \gamma,$$

for some a, b and some $C_{a,b} > 0$ in the (N_p) case, and for every a, b there is $C_{a,b} > 0$ in the $\{N_p\}$ case.

For every measure $d\mu$ on S^{n-1} and

$$U_\mu = \int_{S^{n-1}} U(\cdot + \sqrt{-1}\omega) d\mu(\omega),$$

there holds:

$$WF^* U_\mu \subset \{(x, \omega); -\omega \in \text{supp } d\mu \text{ and } U \text{ is not } \mathcal{O}^* \text{ at } x - \sqrt{-1}\omega\}.$$

$$WF_* U_\mu \subset \{(x, \omega); -\omega \in \text{supp } d\mu \text{ and } U \text{ is not } \mathcal{O}_* \text{ at } x - \sqrt{-1}\omega\}.$$

$$WF_A U_\mu \subset \{(x, \omega); -\omega \in \text{supp } d\mu \text{ and } U \text{ is not analytic at } x - \sqrt{-1}\omega\}.$$

Proof. Let $\omega \in \text{supp } d\mu$ and $\text{Im } z \in \Gamma$. Then for some $C > 0$,

$$1 - |\text{Im}(z + \sqrt{-1}\omega)| > \frac{|\text{Im } z|}{2} \quad \text{if } |\text{Im } z| < C \quad ([5], \text{Lemma 8.4.12}).$$

This implies that we may use Theorem 1 for F . Thus, $F(\dot{+}\sqrt{-1}\Gamma 0) \in \mathcal{S}'^\dagger$.

Note

$$WF^* U_\mu \subset WF_* U_\mu \subset WF_A U_\mu \subset \mathbf{R}^n \times \Gamma^0,$$

where $\Gamma^0 = \{\xi; \langle x, \xi \rangle \geq 0 \text{ for every } x \in \Gamma\}$ is the dual cone.

We shall prove only the estimate for $WF^* U_\mu$ since the other parts can be proved similarly. Denote $\text{sing supp}^* u = \pi_1 WF^* u$. There holds

$$\text{sing supp}^* U_\mu \subset \{x; U \text{ is not } \mathcal{O}^* \text{ at } x + \sqrt{-1}\omega \text{ for some } \omega \in \text{supp } d\mu\}.$$

Decompose

$$d\mu = \sum_{j=1}^r d\mu_j, \quad \text{supp } d\mu_j \subset \text{supp } d\mu \cap \Gamma_j,$$

where Γ_j are open convex cones such that $\text{int } \Gamma_j^0 \neq \emptyset$. By replacing $d\mu$ and Γ from the first part of the lemma by $d\mu_j$ and $-\text{int}(\Gamma_j^0)$ we obtain

$$WF^*U_\mu \subset \bigcup_{j=1}^r \{(x, \xi); -\xi \in \overline{\Gamma_j}, U \text{ is not } \mathcal{O}^* \text{ at } x - \sqrt{-1}\omega \text{ for some } \omega \in \Gamma_j \cap S^{n-1}\}.$$

If $-\frac{\xi}{|\xi|} \notin \text{supp } d\mu$ or U is not $\mathcal{O}^*$ at $x - \sqrt{-1}\frac{\xi}{|\xi|}$, there exists the decomposition of $d\mu$ such that $-\xi \notin \overline{\Gamma_j}$, for $j = 1, \dots, r$ or $-\xi \notin \overline{\Gamma_j}$ for $j = 2, \dots, r$ and U is $\mathcal{O}^*$ at $x + \sqrt{-1}\omega$ for every $\omega \in \Gamma_1 \cap S^{n-1}$. In both cases $(x, \xi) \notin WF^*U_\mu$.

Immediate consequences of this theorem are the next corollary and theorem (see [2], Corollary 8.4.13 and Theorem 8.4.15 for tempered distributions).

Corollary 1. *Let G_j be closed subsets of S^{n-1} such that*

$\bigcup_{j=1}^r G_j = S^{n-1}$. Any $u \in \mathcal{S}'^\dagger(\mathbf{R}^n)$ can be written $u = \sum_{j=1}^r u_j$, $u_j \in \mathcal{S}'^\dagger(\mathbf{R}^n)$ ($\dagger \leq *$) with

- a) $WF^*u_j \subset WF^*u \cap \mathbf{R}^n \times G_j$, $j = 1, \dots, r$.
- b) $WF_*u_j \subset WF_*u \cap \mathbf{R}^n \times G_j$, $j = 1, \dots, r$.
- c) $WF_Au_j \subset WF_Au \cap \mathbf{R}^n \times G_j$, $j = 1, \dots, r$.

If $u \in \mathcal{E}'^\dagger$, then u_j , $j = 1, \dots, r$, have compact supports as well.

If $u = \sum u'_j$ is another such decomposition, then

$$u'_j = u_j + \sum_k u_{jk}, \quad u_{jk} \in \mathcal{S}'^*,$$

$$u_{jk} = -u_{jk} \quad \text{and} \quad WF^*u_{jk} \subset (WF^*u) \cap \mathbf{R}^n \times (G_j \cap G_k).$$

The same holds for WF_*u_{jk} and WF_Au_{jk} .

Theorem 3. *Let Γ be an open convex cone in $\mathbf{R}^n$, $u \in \mathcal{D}'^*(\Omega)$, $\Omega \subset \mathbf{R}^n$, and $WF_*u \subset \Omega \times \Gamma^0$. If $\Omega_1 \subset \subset \Omega$ and Γ_1 is an open convex cone with closure $\subset \Gamma \cup \{0\}$, then there is an F analytic in $\{x + \sqrt{-1}y; x \in \Omega_1, y \in \Gamma_1, |y| < \gamma\}$ such that for some $k > 0$ and $C > 0$ in the (M_p) case (resp., for every $k > 0$ there is $C > 0$ in the $\{M_p\}$ case),*

$$|F(x + \sqrt{-1}y)| \leq Ce^{\tilde{M}(k/|y|)}, \quad x \in \Omega_1, y \in \Gamma_1, |y| < \gamma,$$

and $F(\cdot + \sqrt{-1}0) - u|_{\Omega_1} \in \mathcal{E}^(\Omega_1)$.*

REFERENCES

- [1] A. Eida, On the microlocal decomposition of ultradistributions and ultradifferentiable functions, Preprint.
- [2] L. Hörmander, The Analysis of Linear Partial Differential Operators I, Springer, Berlin - Heidelberg - New York - Tokyo, 1983. MR **91m**:35001b
- [3] H. Komatsu, Ultradistributions, I – III, J. Fac. Sci. Univ. Tokyo, Sect. IA 20(1973), 25-105; 24(1977) 607 - 628; 29(1982), 653 - 717. MR **47**:9277; MR **57**:17280; MR **84d**:46051
- [4] H. Komatsu, Microlocal Analytic in Gevrey Classes and in Convex Domains, Springer, Lec. Not. Math. 1726(1989), 426 - 493.
- [5] D. Kovačević, S. Pilipović, Structural properties of the spaces of tempered ultradistributions, Proc. Conf. "Complex Analysis and Generalized Functions", Varna 1991, 169 - 184. MR **95c**:46059

- [6] H. J. Petzsche, Generalized functions and the boundary values of holomorphic functions, J. Fac. Sci. Univ. Tokyo, Sect. IA Math. 31 (1984), 391-431. MR **85k**:46048
- [7] S. Pilipović, Characterization of bounded sets in spaces of ultradistributions, Proc. AMS, 120(1994), 1191-1206. MR **94f**:46048
- [8] S. Pilipović, Structural Theorems for Ultradistributions, Ann. Polon. Math. 340(1995), 223-235. MR **96f**:46076
- [9] J. W. de Roewer, Hyperfunctional singular support of distributions, J. Fasc. Sci. Univ. Tokyo, Sec IA Math. 31(1985), 585 - 631. MR **86d**:46039
- [10] M. Sato, T. Kawai, M. Kashiwara, Microfunctions and Pseudo - differential Equations, Springer, Lect. Not. Math. 287 (1973), 265-529. MR **54**:8747

INSTITUTE OF MATHEMATICS, UNIVERSITY OF NOVI SAD, TRG D. OBRADOVIĆA 4, 21000 NOVI SAD, YUGOSLAVIA