PRIMES OF THE FORM $p = 1 + m^2 + n^2$ IN SHORT INTERVALS

J. WU

(Communicated by Dennis A. Hejhal)

ABSTRACT. In this note, we prove that for every $\theta \geq \frac{115}{121}$ and $x \geq x_0(\theta)$, the short interval $(x, x + x^\theta]$ contains at least one prime number of the form $p = 1 + m^2 + n^2$ with $(m, n) = 1$. This improves a similar result due to Huxley and Iwaniec, which requires $\theta \geq \frac{99}{100}$.

§1. INTRODUCTION

The existence of prime numbers in short intervals is an important subject in analytic number theory. Huxley and Iwaniec [3] asked the following question: seek positive numbers θ , as small as possible, such that

(1.1). *the short interval $(x, x + x^\theta]$ contains at least one prime number of the form $p = 1 + m^2 + n^2$ with $(m, n) = 1$ for $x \geq x_0(\theta)$.*

Naturally this problem can be attacked by applying the half dimensional sieve to the sequence

$$\mathcal{A} := \{p - 1 : x < p \leq x + x^\theta, p \equiv 3 \pmod{8}\},$$

where, as in the sequel, the letter p , with or without subscript, denotes a prime number. In order to control error terms in formulas of sieves, a Bombieri–Vinogradov type mean-value theorem for short intervals is needed. Huxley and Iwaniec [3] have obtained a satisfactory generalization of Bombieri–Vinogradov’s theorem in the case of short intervals, using a zero-density theorem for Dirichlet L -functions. As an application, they have shown that (1.1) is true with $\theta = \frac{99}{100}$.

The aim of this note is to improve Huxley and Iwaniec’s exponent. More precisely we shall prove the following result.

Theorem. *For every $\theta \geq \frac{115}{121}$ and $x \geq x_0(\theta)$, we have*

$$(1.2) \quad \sum_{x < p \leq x + x^\theta} b^*(p - 1) \gg x^\theta / (\log x)^{3/2},$$

where

$$b^*(a) := \begin{cases} 1 & \text{if } a = m^2 + n^2 \text{ with } (m, n) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Received by the editors December 30, 1994.

1991 *Mathematics Subject Classification.* Primary 11N05, 11N36.

Key words and phrases. Distribution of primes, applications of sieves methods.

For comparison, we have $\frac{115}{121} \approx 0.9504$ and $\frac{99}{100} = 0.99$.

We begin in the same way as Huxley and Iwaniec. By (3.2) of [3], we have

$$(1.3) \quad \sum_{x < p \leq x+x^\theta} b^*(p-1) = S(\mathcal{A}; \mathcal{P}_3, x+x^\theta),$$

where $S(\mathcal{A}; \mathcal{P}_3, z) := |\{a \in \mathcal{A} : (a, \prod_{p < z, p \in \mathcal{P}_3} p) = 1\}|$ and $\mathcal{P}_3 := \{p : p \equiv 3 \pmod{4}\}$. Let $\alpha = \alpha(\theta) \in [2, 3)$ be a parameter to be chosen later. For $z = x^{1/\alpha}$, we write

$$(1.4) \quad S(\mathcal{A}; \mathcal{P}_3, x+x^\theta) = S(\mathcal{A}; \mathcal{P}_3, z) - T,$$

where $T := S(\mathcal{A}; \mathcal{P}_3, z) - S(\mathcal{A}; \mathcal{P}_3, x+x^\theta)$. A lower bound for $S(\mathcal{A}; \mathcal{P}_3, z)$ on the right-hand side of (1.4) will be obtained by the half dimensional sieve (as in [3]) in Section 3. We shall give a better upper bound for T than that of Huxley and Iwaniec. This improvement comes from our generalized Bombieri–Vinogradov type mean-value theorem for short intervals that we state in Section 2. One observes that each element $a \in \mathcal{A}$ is divisible by an even number of primes from $\mathcal{P}_3$ and $2 \parallel a$. Hence, for $z = x^{1/\alpha}$ with $2 \leq \alpha < 3$, we have obviously

$$(1.5) \quad T \leq \sum_{\substack{x < p \leq x+x^\theta \\ p=1+2np_1p_2}} 1,$$

where $p_1 \in \mathcal{P}_3$, $p_2 \in \mathcal{P}_3$, $p_1 \geq p_2 \geq x^{1/\alpha}$ and n is an integer divisible only by primes of the form $p \equiv 1 \pmod{4}$. We define

$$\mathcal{L} := \{l = 2np_2 : n \leq x^{1-2/\alpha}, p \mid n \Rightarrow p \equiv 1 \pmod{4}; \\ x^{1/\alpha} < p_2 \leq (x/n)^{1/2}, p_2 \in \mathcal{P}_3\}.$$

For every $l \in \mathcal{L}$, we put

$$\mathcal{M}(l) := \{m = lp_1 + 1 : x/l < p_1 \leq (x+x^\theta)/l, (l/2)p_1 \equiv 1 \pmod{4}\}.$$

It is clear that the sum on the right-hand side of (1.5) does not exceed the number of primes in the set $\bigcup_{l \in \mathcal{L}} \mathcal{M}(l)$; thus

$$(1.6) \quad T \leq \sum_{l \in \mathcal{L}} \{S(\mathcal{M}(l); \mathcal{P}(l), (x/l)^{\theta_1}) + O((x/l)^{\theta_1})\},$$

where $\mathcal{P}(l) := \{p : (p, l) = 1\}$ and $\theta_1 := (2\theta - 1)/4$. We shall apply an upper bound formula of the linear sieve to treat $S(\mathcal{M}(l); \mathcal{P}(l), (x/l)^{\theta_1})$ in Section 4.

It seems interesting to compare the theorem with the following results: for x sufficiently large, $(x, x+x^{0.973}]$ contains at least one prime of the form $p = -2 + P_2$ where P_2 denotes an integer having at most two prime factors [8]; for every $\varepsilon > 0$ and $x \geq x_0(\varepsilon)$, there exists at least one prime of the form $p = -2 + a$ in $(x, x+x^{3/4+\varepsilon}]$ where a is a $\mathcal{B}$ -free integer [7].

§2. TWO PRELIMINARY LEMMAS

We first recall some standard notations (cf. [2]). Let $\mathcal{F}$ be a finite sequence of integers and let $\mathcal{P}$ be a set of prime numbers. For $z \geq 2$, the sifting function is defined as follows:

$$S(\mathcal{F}; \mathcal{P}, z) := |\{a \in \mathcal{F} : (a, P(z)) = 1\}|,$$

where $P(z) := \prod_{p < z, p \in \mathcal{P}} p$. If d is a squarefree integer whose prime factors belong to $\mathcal{P}$, we let $\mathcal{F}_d := \{n \in \mathcal{F} : d|n\}$ and use $|\mathcal{F}_d|$ to denote the number of elements of $\mathcal{F}_d$. We write an approximate formula

$$|\mathcal{F}_d| = \frac{w(d)}{d} X + r(\mathcal{F}, d),$$

where $X > 1$ is independent of d , and $w(d)$ is a multiplicative function. We define

$$V(z) := \prod_{p < z, p \in \mathcal{P}} (1 - w(p)/p).$$

As usual, $\mu(n)$ is Möbius' function, $\varphi(n)$ Euler's function and $\omega(n)$ the number of distinct prime factors of n . Finally we write ε for an arbitrarily small positive number and γ for Euler's constant.

The first lemma is an upper bound formula of the linear sieve ([2], Theorem 8.3).

Lemma 1. *If there exist positive constants A_k ($k = 1, 2, 3$) such that*

$$\begin{aligned} 0 &\leq \frac{w(p)}{p} < 1 - \frac{1}{A_1}, \\ -A_2 &\leq \sum_{z_1 \leq p < z_2} w(p) \frac{\log p}{p} - \log \frac{z_2}{z_1} \leq A_3 \quad (z_2 \geq z_1 \geq 2), \end{aligned}$$

we then have, for $2 \leq z \leq Q$, that

$$S(\mathcal{F}; \mathcal{P}, z) \leq XV(z) \left\{ F\left(\frac{\log Q}{\log z}\right) + O\left(\frac{A_2}{(\log Q)^{1/14}}\right) \right\} + \sum_{d < Q, d|P(z)} 3^{\omega(d)} |r(\mathcal{F}, d)|,$$

where $F(t) = 2e^\gamma/t$ ($0 < t \leq 2$).

The following lemma is a direct consequence of Theorem 2 of [6].

Lemma 2. *Let $g(l)$ be an arithmetic function satisfying $g(l) \ll 1$ and let*

$$H(x', h, q, a, l) := \sum_{\substack{x' < lp \leq x' + h \\ lp \equiv a \pmod{q}}} -\frac{1}{\varphi(q)} \int_{x'/l}^{(x'+h)/l} \frac{dt}{\log t}.$$

Then for any $A > 0$, there exists a positive constant $B = B(A)$ such that

(2.1)

$$\sum_{q \leq Q} \max_{(a,q)=1} \max_{h \leq x^\theta} \max_{x/2 < x' \leq x} \left| \sum_{\substack{l \leq L \\ (l,q)=1}} g(l) H(x', h, q, a, l) \right| \ll \frac{x^\theta}{(\log x)^A},$$

$$(2.2) \quad \sum_{q \leq Q} \mu(q)^2 3^{\omega(q)} \max_{(a,q)=1} \max_{h \leq x^\theta} \max_{x/2 < x' \leq x} \left| \sum_{\substack{l \leq L \\ (l,q)=1}} g(l) H(x', h, q, a, l) \right| \ll \frac{x^\theta}{(\log x)^A}$$

for $x \geq 10$, $\frac{3}{5} + \varepsilon \leq \theta \leq 1$, $Q = x^{\theta-1/2}/(\log x)^B$ and $L = x^{(5\theta-3)/2-\varepsilon}$.

§3. LOWER BOUND FOR $S(\mathcal{A}; \mathcal{P}_3, x^{1/\alpha})$

The following proposition offers the required lower bound for $S(\mathcal{A}; \mathcal{P}_3, x^{1/\alpha})$.

Proposition 1. *Assume that $\frac{1}{2} \leq \theta \leq 1$ and $2/(2\theta - 1) \leq \alpha \leq 6/(2\theta - 1)$. We then have*

$$(3.1) \quad S(\mathcal{A}; \mathcal{P}_3, x^{1/\alpha}) \geq \{W_1(\theta, \alpha) + o(1)\} x^\theta / (\log x)^{3/2},$$

where

$$W_1(\theta, \alpha) := \frac{AC_3}{\sqrt{4\theta - 2}} \int_1^{\alpha(\theta-1/2)} \frac{dt}{\sqrt{t(t-1)}}$$

and $A := (1/2\sqrt{2}) \prod_{p \equiv 3 \pmod{4}} (1 - p^{-2})^{1/2}$, $C_3 := \prod_{p \equiv 3 \pmod{4}} (1 - (p-1)^{-2})$.

Proof. By (3.3) of [3] or Theorem 1 of [4], we have

$$(3.2) \quad S(\mathcal{A}; \mathcal{P}_3, x^{1/\alpha}) \geq \{1 + o(1)\} \frac{x^\theta V(x^{1/\alpha})}{4 \log x} \sqrt{\frac{e^\gamma}{\pi \xi}} \int_1^\xi \frac{dt}{\sqrt{t(t-1)}} - E(x, x^\theta, Q),$$

where

$$\xi = \alpha \frac{\log Q}{\log x} \in [1, 3], \quad V(x^{1/\alpha}) = \prod_{\substack{p < x^{1/\alpha} \\ p \equiv 3 \pmod{4}}} \left(1 - \frac{1}{p-1}\right),$$

$$E(x, x^\theta, Q) = \sum_{q \leq Q} \max_{(a,q)=1} \max_{h \leq x^\theta} \max_{x/2 < x' \leq x} \left| \sum_{\substack{x' < p \leq x'+h \\ p \equiv a \pmod{q}}} 1 - \frac{1}{\varphi(q)} \int_{x'}^{x'+h} \frac{dt}{\log t} \right|.$$

Let χ be the non-principal character modulo 4 and let $L(\chi, s)$ be the Dirichlet L -function associated with χ . Using the relation $L(\chi, 1; y) := \prod_{p < y} (1 - \chi(p)/p)^{-1} \rightarrow L(\chi, 1) = \frac{1}{4}\pi$ ($y \rightarrow \infty$) and Mertens' formula, we deduce that

$$(3.3) \quad \begin{aligned} V(x^{1/\alpha}) &= \sqrt{2L(\chi, 1; x^{1/\alpha})} \prod_{\substack{p < x^{1/\alpha} \\ p \equiv 3 \pmod{4}}} (1 - p^{-2})^{1/2} (1 - (p-1)^{-2}) \\ &\times \prod_{p < x^{1/\alpha}} (1 - p^{-1})^{1/2} \\ &= \{1 + o(1)\} 2AC_3 (\alpha \pi e^{-\gamma} / \log x)^{1/2} \quad (x \rightarrow \infty). \end{aligned}$$

Taking $Q = x^{\theta-1/2} / (\log x)^B$ and $g(1) = 1, g(l) = 0$ ($l \geq 2$) in (2.1) of Lemma 2, we obtain

$$(3.4) \quad E(x, x^\theta, Q) \ll x^\theta / (\log x)^2.$$

Now the required inequality (3.1) follows from (3.2)–(3.4). $\square$

§4. UPPER BOUND FOR T

The purpose of this section is to prove Proposition 2 below, which gives a better upper bound for T than that of [3]. To prepare for its proof, we first establish two auxiliary results.

Lemma 3. *Let $u(n)$ be the characteristic function of integers whose prime factors are of the form $4m + 1$ and let $f(n) := \prod_{p|n, p>2} (p-1)/(p-2)$. We then have*

$$(4.1) \quad \sum_{n \leq x} u(n)f(n) = (A/C_1)x/(\log x)^{1/2} + O(x/(\log x)^{3/2}),$$

where $C_1 := \prod_{p \equiv 1 \pmod{4}} (1 - (p-1)^{-2})$ and A is defined as in Proposition 1.

Proof. Let $\chi, L(s, \chi)$ be defined as above. It is clear that $u(n)f(n)$ is multiplicative and

$$u(p^\nu)f(p^\nu) = \begin{cases} (p-1)/(p-2) & \text{if } p \equiv 1 \pmod{4}, \\ 0 & \text{otherwise.} \end{cases}$$

A simple calculation shows that for $\Re s > 1$,

$$\sum_{n=1}^{\infty} u(n)f(n)n^{-s} = \prod_{p \equiv 1 \pmod{4}} (1 - p^{-s})^{-1} (1 + (p-2)^{-1}p^{-s}) = \zeta(s)^{1/2} G(s),$$

where $\zeta(s)$ is Riemann's zeta function and

$$G(s) := (L(s, \chi)(1 - 2^{-s}) \prod_{p \equiv 3 \pmod{4}} (1 - p^{-2s}))^{1/2} \prod_{p \equiv 1 \pmod{4}} (1 + (p-2)^{-1}p^{-s}).$$

Using the well known estimation (see [1], Theorem 8.1)

$$L(s, \chi) \ll \log(3 + |\Im s|) \quad \text{for } \Re s \geq 1 - 1/\log(3 + |\Im s|),$$

we see that $\sum_{n=1}^{\infty} u(n)f(n)n^{-s}$ is of type $\mathcal{T}(\frac{1}{2}, \frac{1}{2}; c_0, \delta, M)$, where c_0, δ, M are suitable positive constants (see page 185 of [5] for the definition of $\mathcal{T}(\frac{1}{2}, \frac{1}{2}; c_0, \delta, M)$). Thus Theorem II.5.3 of [5] is applicable. Now our required result follows immediately from this theorem with $N = 0$. $\square$

Lemma 4. *Let $\mathcal{L}, f(n), A, C_1$ be defined as above. Assume that $2 \leq \alpha < 3$. We have*

$$(4.2) \quad \sum_{l \in \mathcal{L}} \frac{f(l)}{l \log^2(x/l)} = \frac{1 + o(1)}{(\log x)^{3/2}} \cdot \frac{A\alpha}{4C_1} \int_2^\alpha \frac{t-2 + (t-1) \log(t-1)}{t^2(t-1)(1-t/\alpha)^{1/2}} dt.$$

Proof. Let Y be the sum on the left-hand side of (4.2) and let $u(n)$ be the function defined as in Lemma 3. We have

$$Y = \frac{1 + o(1)}{2} \sum_{n \leq x^{1-2/\alpha}} \frac{u(n)f(n)}{n} \sum_{\substack{x^{1/\alpha} < p_2 \leq (x/n)^{1/2} \\ p_2 \equiv 3 \pmod{4}}} \frac{1}{p_2 \log^2(x/np_2)}.$$

As usual we put $\pi(t; 4, 3) := \sum_{p \leq t, p \equiv 3 \pmod{4}} 1$. By the Siegel–Walfisz theorem

$$\pi(t; 4, 3) = \frac{1}{2} \int_2^t \frac{dv}{\log v} + O(t e^{-\sqrt{\log t}})$$

and by partial integration, we easily deduce that

$$\begin{aligned}
 (4.3) \quad Y &= \frac{1+o(1)}{2} \sum_{n \leq x^{1-2/\alpha}} \frac{u(n)f(n)}{n} \int_{x^{1/\alpha}}^{(x/n)^{1/2}} \frac{d\pi(t; 4, 3)}{t \log^2(x/nt)} \\
 &= \frac{1+o(1)}{4} \sum_{n \leq x^{1-2/\alpha}} \frac{u(n)f(n)}{n} \int_{x^{1/\alpha}}^{(x/n)^{1/2}} \frac{dt}{\log^2(x/nt) t \log t} \\
 &= \frac{1+o(1)}{4 \log^2 x} \sum_{n \leq x^{1-2/\alpha}} \frac{u(n)f(n)}{nh(n)^2} \left(\frac{\alpha h(n) - 2}{\alpha h(n) - 1} + \log(\alpha h(n) - 1) \right)
 \end{aligned}$$

with $h(n) := 1 - \log n / \log x$. We define

$$U(t) := \sum_{n \leq t} u(n)f(n), \quad K(t) := \frac{1}{th(t)^2} \left(\frac{\alpha h(t) - 2}{\alpha h(t) - 1} + \log(\alpha h(t) - 1) \right).$$

It is not difficult to show that we have, uniformly for $x \geq 10$ and $1 \leq t \leq x^{1-2/\alpha}$,

$$(4.4) \quad K'(t) = -\frac{1}{t^2 h(t)^2} \left(\frac{\alpha h(t) - 2}{\alpha h(t) - 1} + \log(\alpha h(t) - 1) \right) + O\left(\frac{1}{t^2 \log x}\right).$$

Let Z be the last sum on the right-hand side of (4.3). Noticing that $U(1-) = K(x^{1-2/\alpha}) = 0$, using (4.1) and (4.4), we deduce, by partial integration, that

$$\begin{aligned}
 Z &= \int_{1-}^{x^{1-2/\alpha}} K(v) dU(v) = - \int_1^{x^{1-2/\alpha}} U(v) K'(v) dv \\
 &= \frac{A}{C_1} \int_1^{x^{1-2/\alpha}} \left(\frac{\alpha h(v) - 2}{\alpha h(v) - 1} + \log(\alpha h(v) - 1) \right) \frac{dv}{vh(v)^2 (\log v)^{1/2}} + O(1).
 \end{aligned}$$

By the change of variables $t = \alpha h(v)$, we obtain

$$Z = \sqrt{\log x} \frac{A\alpha}{C_1} \int_2^\alpha \frac{t - 2 + (t - 1) \log(t - 1)}{t^2(t - 1)(1 - t/\alpha)^{1/2}} dt + O(1).$$

Inserting this in (4.3) yields (4.2). This completes the proof. $\square$

An upper bound for T is obtained in the following proposition.

Proposition 2. *Assume that $\frac{4}{5} < \theta < 1$ and $2 \leq \alpha < \min\{3, 2/(5 - 5\theta)\}$. We then have*

$$T \leq \{W_2(\theta, \alpha) + o(1)\} x^\theta / (\log x)^{3/2},$$

where

$$W_2(\theta, \alpha) := \frac{AC_3\alpha}{2\theta - 1} \int_2^\alpha \frac{t - 2 + (t - 1) \log(t - 1)}{t^2(t - 1)(1 - t/\alpha)^{1/2}} dt,$$

and A, C_3 are defined as in Proposition 1.

Proof. For every $l \in \mathcal{L}$, it is natural to choose, in Lemma 1,

$$\begin{aligned}
 \mathcal{F} &= \mathcal{M}(l), \quad \mathcal{P} = \mathcal{P}(l), \quad X = \frac{1}{2} \int_{x/l}^{(x+x^\theta)/l} \frac{dt}{\log t}, \\
 w(p) &= \begin{cases} p/(p-1) & \text{if } p \in \mathcal{P}(l), \\ 0 & \text{otherwise.} \end{cases}
 \end{aligned}$$

Let d be a squarefree integer whose prime factors belong to $\mathcal{P}(l)$. By the Chinese Remainder Theorem, the system of congruences $(l/2)p_1 \equiv 1 \pmod{4}$, $lp_1 \equiv -1 \pmod{d}$ has exactly one solution $a^* \pmod{4d}$. Thus we have

$$|\mathcal{M}(l)_d| = \sum_{\substack{x/l < p_1 \leq (x+x^\theta)/l \\ p_1 \equiv a^* \pmod{4d}}} 1, \quad r(\mathcal{M}(l), d) = H(x/l, x^\theta/l, 4d, a^*, 1).$$

According to Lemma 1, one has

(4.5)

$$S(\mathcal{M}(l); \mathcal{P}(l), (x/l)^{\theta_1}) \leq XV((x/l)^{\theta_1}) \left\{ F\left(\frac{\log Q}{\theta_1 \log(x/l)}\right) + o(1) \right\} + R(x, x^\theta, Q, l),$$

where $\theta_1 = (2\theta - 1)/4$ and

$$R(x, x^\theta, Q, l) := \sum_{d < Q, d|P((x/l)^{\theta_1})} 3^{\omega(d)} |r(\mathcal{M}(l), d)|.$$

Using Mertens' formula, we have

$$(4.6) \quad V((x/l)^{\theta_1}) = \prod_{p < (x/l)^{\theta_1}, (p, l)=1} \left(1 - \frac{1}{p-1}\right) = \{1 + o(1)\} \frac{2C_1 C_3 e^{-\gamma} f(l)}{\theta_1 \log(x/l)},$$

where $f(l)$, C_1 are defined as in Lemma 3. Since $\frac{4}{5} < \theta < 1$ and $2 \leq \alpha < 2/(5 - 5\theta)$, we have $\max_{l \in \mathcal{L}} l = 2x^{1-1/\alpha} \leq x^{(5\theta-3)/2-\varepsilon}$ and $x^\theta/l \geq (x/l)^{3/5+\varepsilon/5}$. Taking $Q = (x/l)^{\theta-1/2}/\log^B(x/l)$ and $g(1) = 1$, $g(m) = 0$ ($m \geq 2$), it follows from (2.2) of Lemma 2 that

$$(4.7) \quad R(x, x^\theta, Q, l) \ll x^\theta/l \log^3(x/l).$$

Combining (4.5)–(4.7), one has the following estimate

$$S(\mathcal{M}(l); \mathcal{P}(l), (x/l)^{\theta_1}) \leq \{1 + o(1)\} \frac{C_1 C_3 f(l) x^\theta}{\theta_1 l \log^2(x/l)}.$$

Then summing over l and using Lemma 4, we obtain the desired inequality. $\square$

§5. THE END OF THE PROOF OF THE THEOREM

Assuming that

$$(5.1) \quad \frac{6}{7} < \theta < 1 \quad \text{and} \quad 2/(2\theta - 1) \leq \alpha < \min\{3, 2/(5 - 5\theta)\},$$

we have, by (1.3)–(1.6) and Propositions 1–2, that

$$\sum_{x < p \leq x+x^\theta} b^*(p-1) \geq \{ (AC_3/(2\theta-1))W(\theta, \alpha) + o(1) \} x^\theta / (\log x)^{3/2},$$

where

(5.2)

$$W(\theta, \alpha) := \sqrt{\theta-1/2} \int_1^{\alpha(\theta-1/2)} \frac{dt}{\sqrt{t(t-1)}} - \alpha \int_2^\alpha \frac{t-2+(t-1)\log(t-1)}{t^2(t-1)(1-t/\alpha)^{1/2}} dt.$$

In order to facilitate the calculation on a computer, we eliminate, by integration by parts, the singularity of two integrands on the right-hand side of (5.2). We have

$$\begin{aligned} W(\theta, \alpha) = & \sqrt{4\theta - 2 - 4/\alpha} + \sqrt{\theta - 1/2} \int_1^{\alpha^{(\theta-1/2)}} (t-1)^{1/2} t^{-3/2} dt \\ & - 2\alpha^2 \int_2^\alpha \frac{-t^2 + 6t - 4 - 2(t-1)^2 \log(t-1)}{(t-1)^2 t^3} (1-t/\alpha)^{1/2} dt. \end{aligned}$$

We choose $\theta = \frac{115}{121}$ and $\alpha = 2.349$, which satisfy (5.1). A numerical computation gives us

$$\begin{aligned} W\left(\frac{115}{121}, 2.349\right) \\ = 0.314325 \cdots + 0.671128 \cdots \times 0.008853 \cdots - 2 \times 2.349^2 \times 0.028982 \cdots \\ \geq 4 \times 10^{-4}. \end{aligned}$$

This completes the proof.

REFERENCES

- [1] W.J. Ellison and M. Mendès-France, *Les nombres premiers*, Hermann, 1974. MR **54**:5138
- [2] H. Halberstam and H.-E. Richert, *Sieve Methods*, Academic Press, London 1974. MR **54**:12689
- [3] M.N. Huxley and H. Iwaniec, “Bombieri’s theorem in short intervals”, *Mathematika* **22** (1975), 188–194. MR **52**:10620
- [4] H. Iwaniec, “The half dimensional sieve”, *Acta Arith.* **29** (1976), 69–95. MR **54**:261
- [5] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge University Press, Cambridge, 1995. MR **97e**:11005b
- [6] J. Wu, “Théorèmes généralisés de Bombieri–Vinogradov dans les petits intervalles”, *Quart. J. Math. Oxford* (2), **44** (1993), 109–128. MR **93m**:10090
- [7] J. Wu, “Distribution des nombres $\mathcal{B}$ -libres dans les petits intervalles”, *J. Théorie des Nombres de Bordeaux* **5** (1993), 151–163. MR **94m**:11125
- [8] J. Wu, “Sur l’équation $p + 2 = P_2$ dans les petits intervalles”, *J. London Math. Soc.* (2) **49** (1994), 61–72. MR **94k**:11118

LABORATOIRE DE MATHÉMATIQUES, INSTITUT ELIE CARTAN – CNRS UMR 9973, UNIVERSITÉ
HENRI POINCARÉ (NANCY 1), 54506 VANDŒUVRE-LÈS-NANCY, FRANCE
E-mail address: `wujie@iecn.u-nancy.fr`