UNIQUENESS OF POSITIVE SOLUTIONS FOR STURM-LIOUVILLE BOUNDARY VALUE PROBLEMS

FU-HSIANG WONG

(Communicated by Hal L. Smith)

ABSTRACT. Sufficient conditions for the uniqueness of positive solutions of singular Sturm-Liouville boundary value problems

$$(BVP) \quad \begin{cases} (E) \quad (|u'|^{m-2}u')' + f(t, u, u') = 0, & \text{in } (\theta_1, \theta_2), m \geq 2, \\ (BC) \quad \begin{cases} \alpha_1 u(\theta_1) - \beta_1 u'(\theta_1) = 0, \\ \alpha_2 u(\theta_2) + \beta_2 u'(\theta_2) = 0, \end{cases} \end{cases}$$

where $\alpha_i, \beta_i \geq 0$ and $\alpha_i^2 + \beta_i^2 \neq 0$ ($i = 1, 2$), are established.

1. INTRODUCTION

In this paper, we are concerned with the uniqueness of positive solutions of boundary value problems for the nonlinear differential equation

$$(E) \quad (|u'|^{m-2}u')' + f(t, u, u') = 0, \quad \theta_1 < t < \theta_2, \quad m \geq 2,$$

subject to one of the following sets of boundary conditions:

$$(BC.1) \quad u(\theta_1) = \xi_1 \geq 0, \quad u'(\theta_2) = \xi_2 \geq 0,$$

$$(BC.2) \quad u'(\theta_1) = \xi_1 \leq 0, \quad u(\theta_2) = \xi_2 \geq 0,$$

$$(BC.3) \quad u(\theta_1) = \xi_1 \geq 0, \quad u(\theta_2) = \xi_2 \geq 0,$$

where $m \geq 2$, $(\theta_1, \theta_2) \subseteq (-\infty, \infty)$ and $f: (\theta_1, \theta_2) \times (0, \infty) \times (-\infty, \infty) \rightarrow (0, \infty)$ satisfies

$$(H) \quad \begin{aligned} & f(t, x, y) \text{ is locally Lipschitz continuous for } (x, y) \text{ in} \\ & (0, \infty) \times \{(-\infty, 0) \cup (0, \infty)\}; \quad f(t, x, y)/x^{m-1} \\ & \text{is strictly decreasing with respect to } x \in (0, \infty) \text{ for} \\ & \text{each fixed } (t, y) \in (\theta_1, \theta_2) \times (-\infty, \infty); \text{ and} \\ & \operatorname{sgn}(y)f(t, x, y) \text{ is decreasing with respect to } y \in (-\infty, \infty) \\ & \text{for each fixed } (t, x) \in (\theta_1, \theta_2) \times (0, \infty). \end{aligned}$$

Received by the editors April 17, 1996.

1991 *Mathematics Subject Classification*. Primary 34B15; Secondary 35J25, 35J65, 47H15.

Key words and phrases. Sturm-Liouville BVPs, positive solution, singular and uniqueness.

Furthermore, we use the uniqueness theorems of (E) with respect to the boundary conditions (BC.i) ($i = 1, 2, 3$) to show that

$$(BVP) \quad \begin{cases} (E) \quad (|u'|^{m-2}u')' + f(t, u, u') = 0, & \text{in } (\theta_1, \theta_2), \quad m \geq 2, \\ (BC) \quad \begin{cases} \alpha_1 u(\theta_1) - \beta_1 u'(\theta_1) = 0, \\ \alpha_2 u(\theta_2) + \beta_2 u'(\theta_2) = 0, \end{cases} \end{cases}$$

where $\alpha_i, \beta_i \geq 0$ and $\alpha_i^2 + \beta_i^2 \neq 0$ ($i = 1, 2$) has at most one positive solution in $C^1([\theta_1, \theta_2])$.

Equations of the type (E) arise in studies of radially symmetric solutions (i.e., solutions u that depend only on the variable $r = |x|$) of the m -Laplace equation,

$$(E_1) \quad \nabla \cdot (|\nabla u|^{m-2} \nabla u) + g(|x|, u, \nabla u) = 0, \quad R_0 < |x| < R_1, \quad x \in \mathbb{R}^N, \quad N \geq 2.$$

A radially symmetric solution of (E₁) satisfies the ordinary differential equation

$$(E_2) \quad (|u'|^{m-2}u')' + \frac{N-1}{r}|u'|^{m-2}u' + g(r, u, u') = 0, \quad R_0 < r < R_1.$$

With the change of variables $t = r^{\frac{m-N}{m-1}}$ (for $m \neq N$) or $t = \log r$ (for $m = N$), equation (E₂) can be reduced to an equation of the type (E) or

$$(E^*) \quad (m-1)|u'|^{m-2}u'' + f(t, u, u') = 0, \quad \theta_1 < t < \theta_2, \quad m \geq 2.$$

Conditions for the existence of solutions of equation (E) with respect to (BC.1)–(BC.3) were studied by many authors; see for instance, De Figueiredo, Lions and Nussbaum [6], Granas, Guenther and Lee [9], Kaper, Knaap and Kwong [12], Lions [16], del Pino, Elgueta and Manasevich [19], Rabinowitz [21], Wong [24], and the references therein. The uniqueness problem concerning (E), for the case $m = 2$, has been studied by many authors. For example, Gatica, Olikar and Waltman [7], Kwong [14], Dalmasso [4, 5], Brezis and Oswald [3], Krasnoselskii [13], and the excellent book by Agarwal and Lakshmikantham [1]. However, it seems that very little is known for the case $m \neq 2$. Recently, Naito [17] considered the case $f(t, u, u') = p(t)f(u)$ and established some excellent conditions for uniqueness by using the generalized Prüfer transformation and comparison theorems. In this article, the author attempts to afford a concise approach to study the uniqueness of positive solutions of (E) with boundary conditions (BC.1)–(BC.3) and (BC).

For other related results, we refer the reader to Bobisud [2], Dalmasso [4, 5], Guedda and Veron [10], Naito [17], del Pino and Manasevich [20], O'Regan [22], and Wong and Yu [23].

2. MAIN RESULT

Let u and v be two distinct positive solutions of (E). We define

$$(1) \quad w(t) := \{u(t)\}^{m-1}(|v'(t)|^{m-2}v'(t)) - (|u'(t)|^{m-2}u'(t))\{v(t)\}^{m-1}$$

for $t \in [a, b] \subseteq [\theta_1, \theta_2]$.

It is clear that $w(t)$ satisfies

$$\begin{aligned}
 w'(t) &= u^{m-1}(|v'|^{m-2}v')' - (|u'|^{m-2}u')'v^{m-1} \\
 &\quad + (m-1)u'v'\{|v'|^{m-2}u^{m-2} - |u'|^{m-2}v^{m-2}\} \\
 &= u^{m-1}\{-f(t, v, v')\} - \{-f(t, u, u')\}v^{m-1} \\
 &\quad + (m-1)u'v'\{|v'u|^{m-2} - |u'v|^{m-2}\} \\
 (2) \quad &= (uv)^{m-1} \left\{ \frac{f(t, u, u')}{u^{m-1}} - \frac{f(t, v, v')}{v^{m-1}} \right\} \\
 &\quad + (m-1)u'v'\{|v'u|^{m-2} - |u'v|^{m-2}\} \\
 &= (uv)^{m-1} \left\{ \frac{f(t, u, u')}{u^{m-1}} - \frac{f(t, v, u')}{v^{m-1}} + \frac{f(t, v, u')}{v^{m-1}} - \frac{f(t, v, v')}{v^{m-1}} \right\} \\
 &\quad + (m-1)u'v'\{|v'u|^{m-2} - |u'v|^{m-2}\}
 \end{aligned}$$

for $t \in (a, b) \subseteq (\theta_1, \theta_2)$.

In order to treat our main results, we need the following:

Lemma 2.1. *Let u and v be distinct positive solutions of (E)–(BC.i) in $C^1([\theta_1, \theta_2])$ and $u > v$ on (θ_1, θ_2) for $i = 1, 2, 3$. Then $w'(t) < 0$ in (θ_1, θ_2) , that is, $w(t)$ is strictly decreasing in $[\theta_1, \theta_2]$.*

Proof. We separate the proof into the following cases:

Case (1). Suppose that u and v are two distinct positive solutions of (E)–(BC.1). First, we claim that $u'(t)v(t) \geq v'(t)u(t)$ (≥ 0) in $[\theta_1, \theta_2]$.

(1°) Assume that there exists $t_1 \in (\theta_1, \theta_2)$ such that

$$u'(t)v(t) \geq v'(t)u(t) \quad (\geq 0) \quad \text{on } [\theta_1, t_1] \quad \text{and} \quad u'(t_1)v(t_1) = v'(t_1)u(t_1) \quad (\geq 0),$$

which imply $u'(t) \geq v'(t)$ (≥ 0) on $[\theta_1, t_1]$. It follows from (2), $f(t, x, y)/x^{m-1}$ is strictly decreasing with respect to $x \in (0, \infty)$ and $f(\cdot, \cdot, y)$ is decreasing in $(0, \infty)$ that $w'(t) < 0$ on (θ_1, t_1) . Thus $w(t)$ is a strictly decreasing function on $[\theta_1, t_1]$. Therefore

$$\begin{aligned}
 0 &= (v'(t_1)u(t_1))^{m-1} - (u'(t_1)v(t_1))^{m-1} = w(t_1) \\
 &< w(\theta_1) = \{u(\theta_1)\}^{m-1} [|v'(\theta_1)|^{m-2}v'(\theta_1) - |u'(\theta_1)|^{m-2}u'(\theta_1)] \leq 0,
 \end{aligned}$$

which gives a contradiction.

(2°) Assume that there exists a strictly decreasing sequence $\{t_n\}_{n=1}^\infty$ satisfying $\lim_{n \rightarrow \infty} t_n = \theta_1$, $(u'v - v'u)(t_n) = 0$, $(u'v - v'u)'(t_{2n}) = (u''v - v''u)(t_{2n}) \leq 0$ and $(u'v - v'u)'(t_{2n-1}) = (u''v - v''u)(t_{2n-1}) \geq 0$ for all $n \in \mathbb{N}$. It follows from (E*), (H) and $u'(t_{2n}) \geq v'(t_{2n}) \geq 0$ that

$$\begin{aligned}
 0 &\geq (m-1)|u'|^{m-2}v^{m-2}(u''v - v''u)(t_{2n}) \\
 &= (uv)^{m-1}(t_{2n}) \left\{ \frac{f(t_{2n}, v(t_{2n}), v'(t_{2n}))}{v^{m-1}(t_{2n})} - \frac{f(t_{2n}, u(t_{2n}), u'(t_{2n}))}{u^{m-1}(t_{2n})} \right\} \\
 &= (uv)^{m-1}(t_{2n}) \left\{ \frac{f(t_{2n}, v(t_{2n}), v'(t_{2n}))}{v^{m-1}(t_{2n})} - \frac{f(t_{2n}, u(t_{2n}), v'(t_{2n}))}{u^{m-1}(t_{2n})} \right\} \\
 &\quad + (uv)^{m-1}(t_{2n}) \left\{ \frac{f(t_{2n}, u(t_{2n}), v'(t_{2n}))}{u^{m-1}(t_{2n})} - \frac{f(t_{2n}, u(t_{2n}), u'(t_{2n}))}{u^{m-1}(t_{2n})} \right\} > 0.
 \end{aligned}$$

This gives a contradiction. Hence, we have $u'(t)v(t) \geq v'(t)u(t)$ (≥ 0) in $[\theta_1, \theta_2]$, which implies $w'(t) < 0$ in (θ_1, θ_2) .

Case (2). Suppose that u and v are two distinct positive solutions of (E)–(BC.2). The proof is quite similar to Case (1), thus we omit the details.

Case (3). Suppose that u and v are two distinct positive solutions of (E)–(BC.3). By virtue of Case (1) and Case (2), we need only consider the case $u'(\theta_1) \neq v'(\theta_1)$ and $u'(\theta_2) \neq v'(\theta_2)$. Without loss of generality, we may assume that $\xi_1 \leq \xi_2$ (resp. $\xi_2 \leq \xi_1$), which implies $u'(\theta_1) > v'(\theta_1) \geq 0$ (resp. $u'(\theta_2) < v'(\theta_2) \leq 0$).

(3°) Assume that there exist $t_1, t_2 \in (\theta_1, \theta_2)$ such that $u'(t_1) = v'(t_2) = 0$. Since $|u'|^{m-2}u'$ and $|v'|^{m-2}v'$ are strictly decreasing in (θ_1, θ_2) , t_1 and t_2 are determined uniquely. If $t_1 < t_2$, it follows from $u'(\theta_1) > v'(\theta_1) \geq 0$ and $u'(t_1) = 0 < v'(t_1)$ that there exists $t_3 \in (\theta_1, t_1)$ satisfying

$$u'(t) > v'(t) > 0 \text{ on } [\theta_1, t_3] \quad \text{and} \quad u'(t_3) = v'(t_3) > 0.$$

It follows from $u(\theta_1) = v(\theta_1) = \xi_1 \geq 0$, $u'(t_3) = v'(t_3) > 0$ and Case (1) that $w'(t) < 0$ on (θ_1, t_3) . Thus $w(t)$ is a strictly decreasing function on $[\theta_1, t_3]$. Therefore

$$\begin{aligned} 0 &< \{u'(t_3)\}^{m-1}[\{u(t_3)\}^{m-1} - \{v(t_3)\}^{m-1}] = w(t_3) \\ &< w(\theta_1) = \{u(\theta_1)\}^{m-1}[|v'(\theta_1)|^{m-2}v'(\theta_1) - |u'(\theta_1)|^{m-2}u'(\theta_1)] \leq 0, \end{aligned}$$

which gives a contradiction. If $t_1 > t_2$, it follows from $u'(\theta_2) < v'(\theta_2) \leq 0$ and $v'(t_1) < 0 = u'(t_1)$ that there exists $t_4 \in (t_1, \theta_2)$ such that

$$u'(t_4) = v'(t_4) < 0 \quad \text{and} \quad u'(t) < v'(t) \leq 0 \text{ on } (t_4, \theta_2].$$

It follows from $u(\theta_2) = v(\theta_2) = \xi_2 \geq 0$, $u'(t_4) = v'(t_4) < 0$ and Case (2) that $w'(t) < 0$ on (t_4, θ_2) . Thus $w(t)$ is a strictly decreasing function on $[t_4, \theta_2]$. Therefore

$$\begin{aligned} 0 &\leq \{u(\theta_2)\}^{m-1}[|v'(\theta_2)|^{m-2}v'(\theta_2) - |u'(\theta_2)|^{m-2}u'(\theta_2)] = w(\theta_2) \\ &< w(t_4) = |u'(t_4)|^{m-2}u'(t_4)[\{u(t_4)\}^{m-1} - \{v(t_4)\}^{m-1}] \leq 0, \end{aligned}$$

which gives a contradiction, too. Thus, $t_1 = t_2$. By Cases (1)–(2), we see that $w'(t) < 0$ in (θ_1, θ_2) .

(4°) Assume that there exists $t_1 \in (\theta_1, \theta_2)$ such that $u'(t_1) = 0$ and $v'(t) \neq 0$ in (θ_1, θ_2) . It follows from $u'(\theta_1) > v'(\theta_1) (\geq 0)$ and $u'(t_1) = 0 < v'(t_1)$ that there exists $t_5 \in (\theta_1, t_1)$ satisfying

$$u'(t) > v'(t) > 0 \text{ on } [\theta_1, t_5] \quad \text{and} \quad u'(t_5) = v'(t_5) > 0.$$

Just as in the proof in (3°), we get a contradiction.

(5°) Assume that there exists $t_2 \in (\theta_1, \theta_2)$ such that $v'(t_2) = 0$ and $u'(t) \neq 0$ in (θ_1, θ_2) . Therefore, we obtain $0 \leq u'(\theta_2) < v'(\theta_2) \leq 0$, which gives a contradiction.

(6°) Assume that $u'(t) \neq 0$ and $v'(t) \neq 0$ in (θ_1, θ_2) . It follows from $u'(\theta_1) > v'(\theta_1) (\geq 0)$ and $u'(\theta_2) < v'(\theta_2) (\geq 0)$ that there exists $t_6 \in (\theta_1, \theta_2)$ satisfying

$$u'(t) > v'(t) > 0 \text{ on } [\theta_1, t_6] \quad \text{and} \quad u'(t_6) = v'(t_6) > 0.$$

Just as in the proof in (3°), we get a contradiction.

Theorem 2.2. *The boundary value problem (E)–(BC.1) has at most one positive solution in $C^1([\theta_1, \theta_2])$.*

Proof. Assume to the contrary that u and v are two distinct positive solutions of (E)–(BC.1). We claim that u and v intersect in (θ_1, θ_2) . Suppose, on the contrary,

that $u(t) > v(t)$ in (θ_1, θ_2) . It follows from Lemma 2.1 and $u'(\theta_1) \geq v'(\theta_1) (\geq 0)$ that

$$\begin{aligned} 0 &\leq [\{u(\theta_2)\}^{m-1} - \{v(\theta_2)\}^{m-1}]|v'(\theta_2)|^{m-2}v'(\theta_2) = w(\theta_2) \\ &< w(\theta_1) = \{u(\theta_1)\}^{m-1}[|v'(\theta_1)|^{m-2}v'(\theta_1) - |u'(\theta_1)|^{m-2}u'(\theta_1)] \leq 0, \end{aligned}$$

which gives a contradiction. Hence, there exists $t_1 \in (\theta_1, \theta_2)$ such that $u(t_1) = v(t_1) > 0$. Following from $u(t_1) = v(t_1) > 0$, $u'(\theta_2) = v'(\theta_2) = \xi_2 \geq 0$, and repeating the same process as above, we obtain a $t_2 \in (t_1, \theta_2)$ such that $u(t_2) = v(t_2) > 0$.

Now, we claim that u and v intersect in (t_1, t_2) . Assume, on the contrary, that $u(t) > v(t)$ in (t_1, t_2) ; then $u'(t_1) \geq v'(t_1) \geq 0$ and $0 \leq u'(t_2) \leq v'(t_2)$. From Lemma 2.1 we see that

$$\begin{aligned} 0 &\leq \{u(t_2)\}^{m-1}[|v'(t_2)|^{m-2}v'(t_2) - |u'(t_2)|^{m-2}u'(t_2)] = w(t_2) \\ &< w(t_1) = \{u(t_1)\}^{m-1}[|v'(t_1)|^{m-2}v'(t_1) - |u'(t_1)|^{m-2}u'(t_1)] \leq 0, \end{aligned}$$

which gives a contradiction, too. Hence, there exists $t_3 \in (t_1, t_2)$ such that $u(t_3) = v(t_3) > 0$. Repeating the same argument, we obtain a strictly decreasing sequence $\{t_n\}_{n=3}^\infty \subset (t_1, t_2) \subset (\theta_1, \theta_2)$ such that $t_n \in (t_1, t_{n-1})$ and $u(t_n) = v(t_n)$ for all $n = 3, 4, \dots$. By the Bolzano-Weierstrass theorem, we see that $\{t_n\}_{n=3}^\infty$ has an accumulation point, say η , in $[t_1, t_2]$. It is clear that $u(\eta) = v(\eta) > 0$ and $u'(\eta) = v'(\eta) > 0$. Since $f(t, x, y)$ satisfies (H), it follows from the uniqueness of the non-zero initial value problem that $u(t) = v(t)$ in $[\theta_1, \theta_2]$ (see, for example, Hartman [11]).

Theorem 2.3. *The boundary value problem (E)–(BC.2) has at most one positive solution in $C^1([\theta_1, \theta_2])$.*

Proof. Assume to the contrary that u and v are two distinct positive solutions of (E)–(BC.2). Just as in the proof of Theorem 2.2, we claim that u and v intersect in (θ_1, θ_2) . Suppose, on the contrary, that $u(t) > v(t)$ in (θ_1, θ_2) . It follows from Lemma 2.1 and $u'(\theta_2) \leq v'(\theta_2) (\leq 0)$ that

$$\begin{aligned} 0 &\leq \{u(\theta_2)\}^{m-1}[|v'(\theta_2)|^{m-2}v'(\theta_2) - |u'(\theta_2)|^{m-2}u'(\theta_2)] = w(\theta_2) \\ &< w(\theta_1) = [\{u(\theta_1)\}^{m-1} - \{v(\theta_1)\}^{m-1}]|v'(\theta_1)|^{m-2}v'(\theta_1) \leq 0, \end{aligned}$$

which gives a contradiction. Hence, there exists $t_1 \in (\theta_1, \theta_2)$ such that $u(t_1) = v(t_1) > 0$. Since $u(t_1) = v(t_1) > 0$ and $u'(\theta_2) = v'(\theta_2) = \xi_2 \geq 0$, it follows from Theorem 2.2 that $u(t) = v(t)$ on $[t_1, \theta_2]$. Therefore, $u(t) = v(t)$ in $[\theta_1, \theta_2]$.

Theorem 2.4. *The boundary value problem (E)–(BC.3) has at most one positive solution in $C^1([\theta_1, \theta_2])$.*

Proof. Assume to the contrary that u and v are two distinct positive solutions of (E)–(BC.3). By virtue of Theorems 2.2 and 2.3, we see that $u'(\theta_1) \neq v'(\theta_1)$ and $u'(\theta_2) \neq v'(\theta_2)$. Without loss of generality, we may assume that $u(t) > v(t)$ in (θ_1, θ_2) . Thus $u'(\theta_1) > v'(\theta_1)$. Define t_1 and t_2 so that $u'(t_1) = v'(t_2) = 0$. Similar to the proof of Lemma 2.1, we have that $t_1 = t_2$. Applying Theorems 2.2 and 2.3, we obtain $u \equiv v$ on $[\theta_1, t_1]$ and $u \equiv v$ on $[t_1, \theta_2]$. Therefore, we obtain the desired results.

Theorem 2.5. *The boundary value problem (BVP) has at most one positive solution in $C^1([\theta_1, \theta_2])$.*

Proof. Assume to the contrary that u and v are two distinct positive solutions of (BVP). We split the proof into the following cases.

Case (1). Assume that $\alpha_1 = 0$, that is, $u'(\theta_1) = v'(\theta_1) = 0$. Since $|u'|^{m-2}u'$ and $|v'|^{m-2}v'$ are strictly decreasing in (θ_1, θ_2) , $u'(t) < 0$ and $v'(t) < 0$ in $(\theta_1, \theta_2]$. Now, we claim that u and v intersect in (θ_1, θ_2) . Suppose to the contrary that $u(t) > v(t) > 0$ in (θ_1, θ_2) .

(1°) If $\alpha_2 = 0$, then $u'(\theta_2) = v'(\theta_2) = 0$. This contradicts the fact that $u'(t) < 0$ and $v'(t) < 0$ in $(\theta_1, \theta_2]$.

(2°) If $\beta_2 = 0$, then $u(\theta_2) = v(\theta_2) = 0$. It follows from $u'(\theta_1) = v'(\theta_1) = 0$, $u(\theta_2) = v(\theta_2) = 0$ and Theorem 2.3 that $u(t) = v(t)$ on $[\theta_1, \theta_2]$, which gives a contradiction.

(3°) If $\alpha_2\beta_2 \neq 0$, then $u'(\theta_2)v(\theta_2) = v'(\theta_2)u(\theta_2)$. It is clear that $u(\theta_2) > v(\theta_2) > 0$, and thus $u'(\theta_2) < v'(\theta_2) < 0$. In fact, if $v(\theta_2) = 0$ (resp. $u(\theta_2) = v(\theta_2)$), it follows from (BC) and $u'(\theta_2)v(\theta_2) = v'(\theta_2)u(\theta_2)$ that $u(\theta_2) = v(\theta_2) \geq 0$, with Theorem 2.3 gives a contradiction.

Repeating the similar argument in Case (1) of Lemma 2.1, we can see that $w(t)$ is strictly decreasing on $[\theta_1, \theta_2]$. Therefore, we obtain

$$\begin{aligned} 0 &= (-1)^{m-2}[(v'(\theta_1)u(\theta_1))^{m-1} - (u'(\theta_1)v(\theta_1))^{m-1}] = w(\theta_1) \\ &> w(\theta_2) = (-1)^{m-2}[(v'(\theta_2)u(\theta_2))^{m-1} - (u'(\theta_2)v(\theta_2))^{m-1}] = 0, \end{aligned}$$

which gives a contradiction.

Hence, there is $t_2 \in (\theta_1, \theta_2)$ such that $u(t_2) = v(t_2) > 0$. Since $u'(\theta_1) = v'(\theta_1) = 0$ and $u(t_2) = v(t_2) > 0$, it follows from Theorem 2.3 that $u(t) = v(t)$ on $[\theta_1, t_2]$. Therefore, $u(t) = v(t)$ in $[\theta_1, \theta_2]$.

Case (2). Assume that $\beta_1 = 0$, that is, $u(\theta_1) = v(\theta_1) = 0$. It follows from $|u'|^{m-2}u'$, $|v'|^{m-2}v'$ strictly decreasing in (θ_1, θ_2) and $u(\theta_1) = v(\theta_1) = 0$ that $u'(\theta_1) > 0$ and $v'(\theta_1) > 0$. Now, we claim that u and v intersect in (θ_1, θ_2) . Suppose to the contrary that $u(t) > v(t) > 0$ in (θ_1, θ_2) , and this implies $u'(\theta_1) \geq v'(\theta_1) > 0$.

(4°) If $\alpha_2 = 0$, then $u'(\theta_2) = v'(\theta_2) = 0$. It follows from Theorem 2.2 that $u(t) = v(t)$ in $[\theta_1, \theta_2]$, which gives a contradiction.

(5°) If $\beta_2 = 0$, then $u(\theta_2) = v(\theta_2) = 0$. It follows from Theorem 2.4 that $u(t) = v(t)$ in $[\theta_1, \theta_2]$, which gives a contradiction.

(6°) If $\alpha_2\beta_2 \neq 0$, then $u'(\theta_2)v(\theta_2) = v'(\theta_2)u(\theta_2)$. Repeating the same argument in Case (1) of Lemma 2.1 (or cf. Case (1)–(3°)), we see that $w(t)$ is strictly decreasing on $[\theta_1, \theta_2]$. Therefore, we obtain

$$\begin{aligned} 0 &= \{u(\theta_1)\}^{m-1}[(v'(\theta_1)|v(\theta_1)|^{m-2}v'(\theta_1) - |u'(\theta_1)|^{m-2}u'(\theta_1))] = w(\theta_1) \\ &> w(\theta_2) = \{u(\theta_2)\}^{m-1}(|v'(\theta_2)|^{m-2}v'(\theta_2) - (|u'(\theta_2)|^{m-2}u'(\theta_2))\{v(\theta_2)\}^{m-1}) = 0, \end{aligned}$$

which gives a contradiction.

Hence, there is $t_2 \in (\theta_1, \theta_2)$ such that $u(t_2) = v(t_2) > 0$. Since $u(\theta_1) = v(\theta_1) = 0$ and $u(t_2) = v(t_2) > 0$, it follows from Theorem 2.4 that $u(t) = v(t)$ on $[\theta_1, t_2]$. Therefore, $u(t) = v(t)$ in $[\theta_1, \theta_2]$.

Just as in the proof of Cases (1)–(2), we can exclude the possibility of $\alpha_2 = 0$ or $\beta_2 = 0$.

Case (3). Assume that $\alpha_1\alpha_2\beta_1\beta_2 \neq 0$, that is, $u'(\theta_i)v(\theta_i) = v'(\theta_i)u(\theta_i)$ for $i = 1, 2$. The rest of the proof is quite similar to the proofs in Cases (1)–(3°) and Cases (2)–(6°), so we omit the details.

By Cases (1)–(3), we obtain the desired results.

3. REMARKS AND EXAMPLES

Recently, Gatica, Olikar and Waltman [7], Kwong [14], Naito [17], Brezis and Oswald [3], and Dalmasso [5] showed the following important results:

Theorem 3.A ([3, Theorem 1]). *Consider the problem*

$$(BVP.1) \quad \begin{cases} \Delta u + f(x, u) = 0 & \text{in } \Omega, \\ u \geq 0, \ u \neq 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

and make the following assumptions:

(A₁) For a.e. $x \in \Omega$ the function $u \rightarrow f(x, u)$ is continuous on $[0, \infty)$ and the function $u \rightarrow f(x, u)/u$ is strictly decreasing in $(0, \infty)$.

(A₂) For each $u \geq 0$ the function $u \rightarrow f(x, u)$ belongs to $L^\infty(\Omega)$. Then, there exists at most one solution of (BVP.1) in $H_0^1 \cap L^\infty(\Omega)$.

Theorem 3.B ([5, Theorem 1]). *Let $f \in C^1([0, R] \times [0, \infty))$ satisfying*

(C₁) $uf_u(t, u) > f(t, u)$ for $(t, u) \in [0, R] \times [0, \infty)$, that is, $\frac{f(t, u)}{u}$ is strictly increasing in $u \in [0, \infty)$ for each fixed $t \in [0, R]$;

(C₂) $f_t(u, u) \leq 0$ for $(t, u) \in [0, R] \times [0, \infty)$;

(C₃) there exists $u > 0$ such that $f(R, u) \geq 0$.

Then

$$(BVP.2) \quad \begin{cases} u''(t) + f(|t|, u(t)) = 0, & -R < t < R, \\ u(-R) = u(R) = 0 \end{cases}$$

has at most one positive solution in $C^2[-R, R]$.

Theorem 3.C ([7, Theorem 4.1]). *Let $k \in \{1, 2, 3, \dots\}$, $p \in (0, 1)$ and $h \in C([0, 1]; [0, \infty))$ such that*

$$0 < \int_0^1 (1-t)^{-p} h(t) dt < \infty.$$

Then

$$(E_3) \quad u''(t) + \frac{k}{t} u'(t) + h(t) u^{-p}(t) = 0 \quad \text{in } (0, 1)$$

has at least one positive solution satisfying (BC.2) in $C^1[0, 1] \cap C^2(0, 1)$.

Theorem 3.D ([14, Theorem 2]). *Assume that $q(t) > 0$ in $(0, 1)$ and $f(u)/u$ is decreasing in $(0, \infty)$ and not constant in any neighborhood of $u = 0$. Then*

$$(E_4) \quad u''(t) + q(t)f(u(t)) = 0 \quad \text{in } (0, 1)$$

has a unique positive solution satisfying (BC.2) in $C^1[0, 1] \cap C^2(0, 1)$.

Theorem 3.E ([17, Theorems 1–3]). *Let*

(B₁) $p \in C[\theta_1, \theta_2]$ and $p(t) > 0$ in (θ_1, θ_2) ;

(B₂) $f \in C[0, \infty)$, $f(u) > 0$ in $(0, \infty)$ and $\frac{f(u)}{u^{m-1}}$ is decreasing in $u \in (0, \infty)$;

(B₃) for any $\lambda > 0$ (resp. $\lambda < 0$), a solution u of

$$(E_5) \quad (|u'|^{m-2} u')' + p(t)f(u) = 0, \quad \theta_1 < t < \theta_2, \ m \geq 2,$$

satisfying $u(\theta_1) = 0$ and $u'(\theta_1) = \lambda$ (resp. $u(\theta_2) = 0$ and $u'(\theta_2) = \lambda$) is determined uniquely as long as $u'(t) > 0$ (resp. $u'(t) < 0$).

Then, (E_5) has at most one positive solution in $C^1[\theta_1, \theta_2]$ satisfying (BC.1)–(BC.3).

Remark 3.F. Comparing our uniqueness theorems with the above-mentioned results, we have the following remarks:

(I) The assumption “ $u \rightarrow f(x, u)$ is continuous on $[0, \infty)$ ” in Theorem 3.A implies $f(x, u) \neq u^p$ for $p < 0$, and “ $f(u)/u$ is decreasing in $(0, \infty)$ and is not a constant in any neighborhood of $u = 0$ ” in Theorem 3.D is imposed to exclude the situation in which $f(u)$ behaves like a linear function in a neighborhood of $u = 0$, that is, $f(u)/u$ behaves like a strictly decreasing function near $u = 0$.

(II) It is clear that if $f(t, u)$ is (strictly) decreasing in $u \in (0, \infty)$ and

$$f(t, u) \equiv h(t)u^{-p}, h(t)u^q, u^\alpha + u^{-\alpha}, \sin(t)u^{-p} + \cos(t)u^q$$

for any given $p \in [0, \infty)$, $q \in [0, m-1]$, $\alpha \in [0, m-1]$ and $h \in C((0, 1); [0, \infty))$, then $f(t, u)$ satisfies “ $f(t, u)/u^{m-1}$ is strictly decreasing in u ”. But Theorems 3.A, 3.C, 3.D and 3.E cannot be applied to most of these functions, for example,

$$f(t, u) \equiv u^{1/2}, u^{-1} + u \text{ and } tu^{\frac{1}{3}} + e^t u^{\frac{1}{2}}.$$

Furthermore, Theorem 3.C does not tell us “the uniqueness of positive solution of (E_3) with (BC.2)”.

(III) Our main results generalize Theorems 3.A, 3.C, 3.D, 3.E and also confirm the uniqueness of Theorem 3.C.

(IV) The excellent uniqueness Theorem 3.B of Dalmasso [5] combines with the main results in this article. They can criticize almost all the uniqueness of positive solutions of (E) –(BC. i) ($i = 1, 2, 3$) and (BVP).

Example 3.G. (I) It follows from Theorem 2.2 (resp. Theorems 2.3, 2.4, 2.5) that the boundary value problem

(BVP.3)

$$\begin{cases} (|u'|^{m-2}u')' + 2[t(1-t)]^2u^{-p} + \frac{\sin(t)|u'|^{-\frac{1}{3}}}{1+t^2} = 0 & \text{in } (0, 1), \\ u(0) = u'(1) = 0 \\ \text{(resp. } u'(0) = u(1) = 0, u(0) = u(1) = 0, u(0) = u(1) + 2u'(1) = 0) \end{cases}$$

has at most one positive solution in $C^1[0, 1]$, where $p \in (0, \infty)$.

(II) It follows from Theorem 2.3 (resp. Theorems 2.2, 2.4, 2.5) that the boundary value problem

(BVP.4)

$$\begin{cases} (|u'|^{m-2}u')' + \frac{1}{1+t^2}u^p = 0 & \text{in } (0, 1), p \in (-\infty, m-1), \\ u'(0) = u(1) = 0 \\ \text{(resp. } u(0) = u'(1) = 0, u(0) = u(1) = 0, u(0) = u(1) + 2u'(1) = 0) \end{cases}$$

has at most one positive solution in $C^1[0, 1]$.

(III) It follows from Theorem 2.4 (resp. Theorems 2.2, 2.3, 2.5) that the boundary value problem

$$(BVP.5) \quad \begin{cases} (|u'|^{m-2}u')' + \frac{1}{t^{\alpha+1}}(u^\alpha + u^{-\alpha}) + \cos(t)|u'|^{-\beta} = 0 & \text{in } (0, 1), \\ u(0) = u(1) = 0 \\ \text{(resp. } u(0) = u'(1) = 0, u'(0) = u(1) = 0, u(0) = u(1) + 2u'(1) = 0) \end{cases}$$

has at most one positive solution in $C^1[0, 1]$, where $\alpha \in [0, m - 1]$, $\beta > 0$.

REFERENCES

- [1] R. P. Agarwal and V. Lakshmikantham, *Uniqueness and nonuniqueness criteria for ordinary differential equations*, World Scientific, Singapore, 1993. MR **96e**:34002
- [2] L. E. Bobisud, *Steady state turbulent flow with reaction*, Rocky Mount. J. Math. **21** (1991), 993–1007. MR **93a**:34026
- [3] H. Brezis and L. Oswald, *Remarks on sublinear elliptic equations*, Nonlinear Analysis T. M. & A. **19** (1986), 55–64. MR **87c**:35057
- [4] R. Dalmasso, *Uniqueness of positive and nonnegative solutions of nonlinear equations*, Funkcial. Ekvac **37** (1994), 461–482. MR **97c**:34037
- [5] ———, *Uniqueness of positive solutions of nonlinear second-order equations*, Proc. Amer. Math. Soc. **123** (1995), 3417–3424. MR **96a**:34035
- [6] D. De Figueiredo, P. L. Lions and R. Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures. App. **61** (1982), 41–63. MR **83h**:35039
- [7] J. A. Gatica, V. Oliker and P. Waltman, *Singular nonlinear boundary value problems for second-order ordinary differential equations*, **79** (1989), 62–78. MR **90f**:34030
- [8] B. Gidas, W. M. Ni and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209–243. MR **80h**:35043
- [9] A. Granas, R. B. Guenther and J. W. Lee, *Nonlinear boundary value problems for ordinary differential equations*, Dissertationes (Rozprawy Mat) **244** (1985). MR **87a**:34025
- [10] M. Guedda and L. Veron, *Bifurcation phenomena associated to the p -Laplace operator*, Trans. Amer. Math. Soc. **310** (1988), 419–431. MR **89j**:35024
- [11] P. Hartman, *Ordinary differential equations*, Birkhauser, Boston, 1982. MR **83e**:34002
- [12] H. G. Kaper, M. Knapp and M. K. Kwong, *Existence theorems for second order boundary value problems*, Differential Integral Equations **4** (1991), 543–554. MR **92m**:34059
- [13] M. Krasnoselskii, *Positive solutions of operator equations*, Noordhoff, Groningen, 1964. MR **31**:6107
- [14] M. K. Kwong, *On the Kolodner-Coffman method for the uniqueness problem of Emden-Fowler BVP*, Z. Angew. Math. Phys. **41** (1990), 79–104. MR **91b**:34037
- [15] ———, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in $\mathbb{R}^n$* , Arch. Rational. Mech. Anal. **105** (1989), 243–266. MR **90d**:35015
- [16] P. L. Lions, *On the existence of positive solutions of semilinear elliptic equations*, SIAM Rev. **24** (1982), 441–467. MR **84a**:35093
- [17] Y. Naito, *Uniqueness of positive solutions of quasilinear differential equations*, Differential Integral Equations **8** (1995), 1813–1822. MR **96g**:34039
- [18] W. M. Ni and R. Nussbaum, *Uniqueness and nonuniqueness for positive radial solutions of $\Delta u + f(r, u) = 0$* , Comm. Pure. Appl. Math. **38** (1985), 69–108. MR **86d**:35055
- [19] M. del Pino, M. Elgueta and R. Manasevich, *A homotopic deformation along p of a Leray-Schauder degree result and existence for $(|u'|^{p-2}u')' + f(t, u) = 0$, $u(0) = u(T) = 0$, $p > 1$* , J. Differential Equations **80** (1989), 1–13. MR **91i**:34018
- [20] M. del Pino and R. Manasevich, *Multiple solutions for the p -Laplacian under global nonresonance*, Proc. Amer. Math. Soc. **112** (1991), 131–138. MR **91h**:34026
- [21] P. H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, CBMS Regional Conf. Ser. in Math., Amer. Math. Soc., Providence, RI, 1986. MR **87j**:58024
- [22] Donal O'Regan, *Theory of singular boundary value problems*, World Scientific, Singapore, 1994. MR **95g**:34003

- [23] F. H. Wong and S. L. Yu, *Uniqueness of positive solutions for singular nonlinear boundary value problems*, Hiroshima Mathematical Journal **26** (1996), 233–243. MR **97d**:35044
- [24] F. H. Wong, *Existence of positive solutions of singular boundary value problems*, Nonlinear analysis T.M. & A. **21** (1993), 397–406. MR **94i**:34056

DEPARTMENT OF MATHEMATICS AND SCIENCE, NATIONAL TAIPEI TEACHER'S COLLEGE, 134,
HO-PING E. RD. SEC. 2, TAIPEI 10659, TAIWAN, REPUBLIC OF CHINA
E-mail address: `wong@tea.ntptc.edu.tw`