FINITISTIC DIMENSION AND ZIEGLER SPECTRUM

HENNING KRAUSE

(Communicated by Ken Goodearl)

ABSTRACT. Given a two-sided artinian ring Λ , it is shown that the Ziegler spectrum of Λ forms a test class for certain homological properties of Λ . We discuss the finitistic dimension of Λ , Nunke's condition, and also the relation between the big and the little finitistic dimension.

Let Λ be a two-sided artinian ring. Denote by $\text{Mod } \Lambda$ the category of (right) Λ -modules and by $\text{mod } \Lambda$ the full subcategory of all finitely presented Λ -modules. Given a Λ -module M , we denote by $\text{pd } M$ its projective dimension, and the *finitistic dimension* $\text{Fin. dim } \Lambda$ of Λ is the supremum of the projective dimensions of the Λ -modules with finite projective dimension. The *Ziegler spectrum* $\text{Zsp } \Lambda$ of Λ is by definition the set of isomorphism classes of indecomposable pure-injective Λ -modules [14].

The aim of this note is to show that the Ziegler spectrum forms a test class for certain homological properties of Λ . Furthermore, the Ziegler spectrum carries a topology, and we shall use its compactness to obtain an equivalent formulation of Nunke's condition. In the final part of this note we study the relation between $\text{Fin. dim } \Lambda$ and the little finitistic dimension $\text{fin. dim } \Lambda$ of Λ .

Our first result is easily stated as follows.

Theorem 1. $\text{Fin. dim } \Lambda = \sup\{\text{pd } M \mid M \in \text{Zsp } \Lambda \text{ and } \text{pd } M < \infty\}$.

We postpone the proof and first discuss some related homological properties of Λ . Suppose that Λ has a self-duality D between $\text{mod } \Lambda$ and $\text{mod } \Lambda^{\text{op}}$. We are interested in *Nunke's condition* [9]:

$$\text{Nd } M < \infty \text{ for every } \Lambda\text{-module } M \neq 0,$$

where $\text{Nd } M$ denotes the smallest integer n (or ∞ if such an integer does not exist) such that $\text{Ext}_{\Lambda}^n(D\Lambda, M) \neq 0$. It is convenient to define the *Nunke dimension* of Λ as follows:

$$\text{Nun. dim } \Lambda = \sup\{\text{Nd } M \mid M \in \text{Mod } \Lambda \text{ and } M \neq 0\}.$$

The following relation between Nunke dimension and finitistic dimension is essentially due to Jans [10] (see also [5]).

Proposition 2. $\text{Nun. dim } \Lambda \leq \text{Fin. dim } \Lambda$.

Proof. We use the fact that the functor

$$\text{Inj } \Lambda \longrightarrow \text{Proj } \Lambda, \quad X \mapsto \text{Hom}_{\Lambda}(D\Lambda, X)$$

Received by the editors June 7, 1996 and, in revised form, September 25, 1996.
 1991 *Mathematics Subject Classification.* Primary 16E10, 16G10.

is an equivalence between the full subcategories of injective and projective Λ -modules. Suppose now that $\text{Nun. dim } \Lambda \geq n$ and choose a non-zero Λ -module M with $\coprod_{i=0}^{n-1} \text{Ext}_{\Lambda}^i(D\Lambda, M) = 0$. Take an injective coresolution

$$0 \longrightarrow M \longrightarrow I_0 \xrightarrow{\psi_0} I_1 \xrightarrow{\psi_1} I_2 \xrightarrow{\psi_2} \dots$$

and put $N = \text{Coker Hom}_{\Lambda}(D\Lambda, \psi_{n-1})$. It is easily checked that $\text{pd } N = n$. Thus $\text{Fin. dim } \Lambda \geq n$. $\square$

Remark. Given any $n \in \mathbb{N}_0$, there is a finite dimensional algebra Λ over a field (with $\text{rad}^2 \Lambda = 0$ and $\text{Nun. dim } \Lambda = 2$) such that $\text{Fin. dim } \Lambda - \text{Nun. dim } \Lambda = n$ [7].

Theorem 3. $\text{Nun. dim } \Lambda = \sup\{\text{Nd } M \mid M \in \text{Zsp } \Lambda\}$, and Nunke's condition holds if and only if $\text{Nun. dim } \Lambda < \infty$.

Some further definitions are needed in order to give the proof of both theorems. An additive functor $F: \text{Mod } \Lambda \rightarrow \text{Ab}$ into the category of abelian groups is *finitely presented* if there is a presentation

$$\text{Hom}_{\Lambda}(Y, _) \longrightarrow \text{Hom}_{\Lambda}(X, _) \longrightarrow F \longrightarrow 0$$

with X and Y in $\text{mod } \Lambda$. We denote by $\text{Ker } F$ the full subcategory of Λ -modules M with $F(M) = 0$. A full subcategory $\mathcal{X}$ of $\text{Mod } \Lambda$ is *definable* if there is a family F_i , $i \in I$, of finitely presented functors such that $\mathcal{X} = \bigcap_{i \in I} \text{Ker } F_i$. A definable subcategory is automatically closed under direct limits, products and pure submodules, and it can be shown that this property characterizes a definable subcategory [12]. The notion of a definable subcategory allows us to describe the topology on $\text{Zsp } \Lambda$ which was first introduced by Ziegler in model theoretic terms [14].

Proposition 4. (1) *The assignment $\mathcal{X} \mapsto \mathcal{X} \cap \text{Zsp } \Lambda$ defines a bijection between the definable subcategories of $\text{Mod } \Lambda$ and the closed subsets of $\text{Zsp } \Lambda$.*

(2) *$\text{Zsp } \Lambda$ is a quasi-compact space.*

Proof. (1) This result is due to Crawley-Boevey [3], and he relies on work of Herzog [6] and Ziegler [14]. We refer to [12] for a proof which uses the localization theory for locally coherent categories developed in [11]. However, the inverse of the assignment $\mathcal{X} \mapsto \mathcal{X} \cap \text{Zsp } \Lambda$ is easily constructed; it sends a closed subset $\mathcal{U}$ of $\text{Zsp } \Lambda$ to $\bigcap_{F(\mathcal{U})=0} \text{Ker } F$.

(2) See [14] or [11]. $\square$

The next lemma shows that some relevant subcategories of $\text{Mod } \Lambda$ are definable.

Lemma 5. *Let $n \in \mathbb{N}_0$.*

(1) *The full subcategory $\mathcal{P}_n$ of Λ -modules M with $\text{pd } M \leq n$ is definable.*

(2) *The full subcategory $\mathcal{Q}_n$ of Λ -modules M with $\text{Nd } M \geq n$ is definable.*

Proof. Projective dimension and flat dimension coincide for any module over an artinian ring, and therefore $\mathcal{P}_n = \bigcap_{X \in \text{mod } \Lambda^{\text{op}}} \text{Ker Tor}_{n+1}^{\Lambda}(_, X)$ since $\text{Tor}_{n+1}^{\Lambda}(M, _)$ commutes with direct limits and every module can be written as a direct limit of finitely presented modules. Therefore $\mathcal{P}_n$ is definable, and $\mathcal{Q}_n$ is definable since $\mathcal{Q}_n = \text{Ker } \coprod_{i=0}^{n-1} \text{Ext}_{\Lambda}^i(D\Lambda, _)$. $\square$

An easy consequence of the preceding lemma is the following.

Proposition 6. *Let $\mathcal{U}$ be a subset of $\text{Zsp } \Lambda$. Then $\text{pd } M \leq \sup\{\text{pd } N \mid N \in \mathcal{U}\}$ for every module M in the closure of $\mathcal{U}$.*

Proof of Theorem 1. Suppose that $\mathcal{P}_n$ is properly contained in $\mathcal{P}_{n+1}$. The first part of Proposition 4 then implies there is $M \in \text{Zsp } \Lambda$ with $M \in \mathcal{P}_{n+1} \setminus \mathcal{P}_n$, since $\mathcal{P}_n$ and $\mathcal{P}_{n+1}$ are definable. Thus the assertion follows. $\square$

Proof of Theorem 3. The proof of the first part of the assertion is analogous to that of Theorem 1. Suppose now that Nunke's condition holds. Thus $\bigcap_{n \geq 0} \mathcal{Q}_n = 0$, and the compactness of $\text{Zsp } \Lambda$ then implies the existence of $d \in \mathbb{N}_0$ with $\bigcap_{n=0}^d \mathcal{Q}_n = 0$. We conclude that $\text{Nun. dim } \Lambda < d$. $\square$

Remark. The finitistic dimension of Λ is finite if and only if the full subcategory of all Λ -modules with finite projective dimension is definable. Therefore one might conjecture that $\text{Fin. dim } \Lambda < \infty$ if and only if the modules in the Ziegler spectrum having finite projective dimension form a closed subset of $\text{Zsp } \Lambda$.

The rest of this paper is devoted to studying the relation between the finitistic dimension $\text{Fin. dim } \Lambda$ and the *little finitistic dimension* $\text{fin. dim } \Lambda$, which is the supremum of the projective dimensions of the finitely presented Λ -modules with finite projective dimension. Note that there are examples of finite dimensional algebras where $\text{fin. dim } \Lambda$ is different from $\text{Fin. dim } \Lambda$ [15]. Our results are motivated by recent work of Huisgen-Zimmermann and Smalø, who have shown that both dimensions coincide for an interesting class of Artin algebras [8].

In order to state the next result we recall from [2] that for any subcategory $\mathcal{C}$ of $\text{Mod } \Lambda$ a morphism $M \rightarrow M'$ is a *left $\mathcal{C}$ -approximation* of M if M' lies in $\mathcal{C}$ and the induced map $\text{Hom}_\Lambda(M', X) \rightarrow \text{Hom}_\Lambda(M, X)$ is surjective for every X in $\mathcal{C}$. We denote by $\mathcal{P}_\infty$ the full subcategory of all Λ -modules with finite projective dimension.

Theorem 7. *The following conditions are equivalent.*

- (1) *A Λ -module has finite projective dimension if and only if it is a direct limit of finitely presented modules having finite projective dimension.*
- (2) *The little finitistic dimension $\text{fin. dim } \Lambda$ is finite, and every finitely presented Λ -module has a left $\mathcal{P}_\infty$ -approximation which is finitely presented.*

Moreover, if (1)–(2) hold, then $\text{fin. dim } \Lambda = \text{Fin. dim } \Lambda < \infty$.

Examples of rings which satisfy the conditions in this theorem are Artin algebras Λ where every finitely presented Λ -module has a right $\mathcal{P}_\infty \cap \text{mod } \Lambda$ -approximation [1, 8]. There are also examples of Artin algebras Λ which show that in condition (2) the term “ $\mathcal{P}_\infty$ -approximation” cannot be replaced by “ $\mathcal{P}_\infty \cap \text{mod } \Lambda$ -approximation” [13]. The proof of Theorem 7 is based on the following lemma.

Lemma 8. *Let $\mathcal{C}$ be any additive subcategory of $\text{mod } \Lambda$. Then a Λ -module M is a direct limit of objects in $\mathcal{C}$ iff every morphism $X \rightarrow M$ with $X \in \text{mod } \Lambda$ factors through an object in $\mathcal{C}$.*

Proof. See [4, Lemma 4.1]. $\square$

Proof of Theorem 7. (1) $\Rightarrow$ (2) In order to show that $\text{fin. dim } \Lambda < \infty$ assume there is a family M_i , $i \in \mathbb{N}$, in $\text{mod } \Lambda$ with $\text{pd } M_i < \text{pd } M_{i+1} < \infty$ for all i . Then $\coprod_{i \in \mathbb{N}} M_i$ is a direct limit of finitely presented objects in $\mathcal{P}_\infty$, but $\text{pd } \coprod_{i \in \mathbb{N}} M_i = \infty$. Thus (1)

implies $\text{fin. dim } \Lambda = n < \infty$ and therefore $\text{Fin. dim } \Lambda = n$, since $\mathcal{P}_n$ is closed under direct limits by Lemma 5. To show the second part of the assertion we use an idea from [4]. Let M be in $\text{mod } \Lambda$ and choose a representative set $\varphi_i: M \rightarrow N_i$, $i \in I$, of morphisms in $\text{mod } \Lambda$ such that the domain has finite projective dimension. Then $N = \prod_{i \in I} N_i$ lies in $\mathcal{P}_\infty$, since $\mathcal{P}_\infty = \mathcal{P}_n$ is closed under products by Lemma 5. Thus the induced morphism $\varphi: M \rightarrow N$ factors through a finitely presented object M' in $\mathcal{P}_\infty$ by the preceding lemma. It is easily checked that the corresponding morphism $M \rightarrow M'$ is a left $\mathcal{P}_\infty$ -approximation of M .

(2) $\Rightarrow$ (1) It follows immediately from the preceding lemma that every module in $\mathcal{P}_\infty$ is a direct limit of finitely presented modules in $\mathcal{P}_\infty$. Conversely, the projective dimension of a direct limit of finitely presented Λ -modules in $\mathcal{P}_\infty$ is bounded by $\text{fin. dim } \Lambda = n$, since $\mathcal{P}_n$ is closed under direct limits by Lemma 5. $\square$

If the ring Λ has the property that every finitely presented Λ -module is pure-injective, for instance if Λ is an Artin algebra, then condition (1) in the preceding theorem implies that the finitely presented modules form a dense subset of $\{M \in \text{Zsp } \Lambda \mid \text{pd } M < \infty\}$. We point out that this condition is already sufficient for the equality $\text{fin. dim } \Lambda = \text{Fin. dim } \Lambda$.

Corollary 9. *Let $n \in \mathbb{N}_0$ and suppose that the finitely presented modules $M \in \text{Zsp } \Lambda$ with $\text{pd } M \leq n$ form a dense subset of $\{M \in \text{Zsp } \Lambda \mid \text{pd } M < \infty\}$. Then $\text{fin. dim } \Lambda = \text{Fin. dim } \Lambda \leq n$.*

Proof. Let $\mathcal{U}_n = \{M \in \text{Zsp } \Lambda \mid \text{pd } M \leq n \text{ and } M \in \text{mod } \Lambda\}$, and we may assume that $n \in \mathbb{N}_0$ is the minimal choice with $\{M \in \text{Zsp } \Lambda \mid \text{pd } M < \infty\} \subseteq \overline{\mathcal{U}_n}$. It follows from Proposition 6 and Theorem 1 that $\text{Fin. dim } \Lambda \leq n$, and we therefore obtain $n \leq \text{fin. dim } \Lambda \leq \text{Fin. dim } \Lambda \leq n$. Thus $\text{fin. dim } \Lambda = \text{Fin. dim } \Lambda = n$. $\square$

REFERENCES

- [1] M. Auslander and I. Reiten, Applications of contravariantly finite subcategories, *Adv. Math.* **86** (1991) 111–152. MR **92e**:16009
- [2] M. Auslander and S.O. Smalø, Preprojective modules over Artin algebras, *J. Algebra* **66** (1980) 61–122. MR **83a**:16039
- [3] W. Crawley-Boevey, Seminar talk at Universität Bielefeld (1993).
- [4] W. Crawley-Boevey, Locally finitely presented additive categories, *Comm. Algebra* **22** (1994) 1641–1674. MR **95h**:18009
- [5] D. Happel, On Gorenstein algebras, in: *Representation Theory of Finite Groups and Finite-dimensional Algebras*, Birkhäuser Verlag, Basel (1991) 389–404. MR **92k**:16022
- [6] I. Herzog, The Ziegler spectrum of a locally coherent Grothendieck category, *Proc. London Math. Soc.* (3) **74** (1997), 503–558. CMP 97:08
- [7] B. Huisgen-Zimmermann, Private communication (1996).
- [8] B. Huisgen-Zimmermann and S.O. Smalø, A homological bridge between finite and infinite dimensional representations of algebras, preprint (1996).
- [9] J.P. Jans, Duality in Noetherian rings, *Proc. Amer. Math. Soc.* **12** (1961) 829–835. MR **25**:1192
- [10] J.P. Jans, Some generalizations of finite projective dimension, *Illinois J. Math.* **5** (1961) 334–344. MR **32**:1226
- [11] H. Krause, The spectrum of a locally coherent category, *J. Pure Appl. Algebra*, **114** (1997), 259–271. CMP 97:06
- [12] H. Krause, Exactly definable categories, *J. Algebra*, to appear.
- [13] S.O. Smalø, The supremum of the difference between the big and little finitistic dimensions is infinite, preprint (1996).

- [14] M. Ziegler, Model theory of modules, Ann. of Pure and Appl. Logic **26** (1984) 149–213. MR **86c**:03034
- [15] B. Zimmermann-Huisgen, Homological domino effects and the first finitistic dimension conjecture, Invent. Math. **108** (1992) 369–383. MR **93i**:16016

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, 33501 BIELEFELD, GERMANY
E-mail address: `henning@mathematik.uni-bielefeld.de`