SOME NEARLY BOOLEAN ORTHOMODULAR POSETS

PAVEL PTÁK

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let L be an orthomodular partially ordered set (“a quantum logic”). Let us say that L is nearly Boolean if L is set-representable and if every state on L is subadditive. We first discuss conditions under which a nearly Boolean OMP must be Boolean. Then we show that in general a nearly Boolean OMP does not have to be Boolean. Moreover, we prove that an arbitrary Boolean algebra may serve as the centre of a (non-Boolean) nearly Boolean OMP.

1. INTRODUCTION AND BASIC NOTIONS

Orthomodular posets (OMPs) are certain “nondistributive” generalizations of Boolean algebras. They are sometimes called “quantum logics” when viewed as event structures of quantum experiments. The states of the experiments are then associated with probability measures on the corresponding OMPs, called “states” for this reason, which do not in general enjoy standard Boolean features. For instance, they do not have to be subadditive (a state s on an orthomodular poset L is said to be subadditive if for any $a, b \in L$ there exists $c \in L$ such that $c \geq a, c \geq b$ and $s(c) \leq s(a) + s(b)$). In fact, the subadditivity of all states is quite a strong condition imposed on L . If, moreover, we require L to be set-representable, which is a conceptually important “Boolean-like” condition, we find L to be intrinsically very near a Boolean algebra. A natural question arises whether these “nearly Boolean” OMPs do not have to be Boolean. This turned out to be a fairly nontrivial problem. Moreover, since subadditive states can be viewed as “strongly Jauch-Piron states” (see [1] and [13] for comments on the Jauch-Piron property in quantum theories), the problem has a bearing on quantum axiomatics. We show in this paper that non-Boolean nearly Boolean OMPs exist abundantly. (Observe that the σ -complete version of this problem, which is quite different in its mathematical nature, has been solved in [3]. Notice also that for lattices the abundance of subadditive states implies distributivity [14], and therefore in this case one finds oneself in the realm of Boolean algebras.)

It should be noted that the construction we present here is not entirely new – the technique of [6] and [7] designed for Jauch-Piron states is used as an essential

Received by the editors December 16, 1996.

1991 *Mathematics Subject Classification.* Primary 28A60, 06C15, 81P10.

Key words and phrases. Orthomodular partially ordered set, Boolean algebra, state (= finitely additive probability measure), subadditivity.

The author acknowledges the support by the grant GA 201/96/0117 of the Czech Grant Agency.

tool. We have adapted it to our purpose in some places. Explicit novelties seem to be Th. 2.1 (v), Prop. 2.2 and Th. 2.5.

Let us now introduce the notions we shall need in the sequel.

1.1. Definition. Let Ω be a set and let Δ be a collection of subsets of Ω . The pair (Ω, Δ) is called a set-representable orthomodular poset (a SROMP) if the following three conditions are satisfied:

- (i) $\Omega \in \Delta$,
- (ii) if $A \in \Delta$, then $\Omega - A \in \Delta$,
- (iii) if $A, B \in \Delta$ and $A \cap B = \emptyset$, then $A \cup B \in \Delta$.

The SROMPs have proved to be of a considerable conceptual value for both noncommutative measure theory and quantum axiomatics (see [11] and [4]). It can be easily shown ([13]) that the SROMPs are exactly those (general) orthomodular posets which admit a set-theoretic representation with set-theoretic union and set-theoretic orthocomplementation for the OMP operations. Obviously, a SROMP is a Boolean algebra exactly when it is closed under the formation of intersections.

1.2. Definition. Let (Ω, Δ) be a SROMP. By a state on (Ω, Δ) we mean a mapping $s: \Delta \rightarrow \langle 0, 1 \rangle$ such that

- (i) $s(\Omega) = 1$,
- (ii) $s(A \cup B) = s(A) + s(B)$ provided $A, B \in \Delta$ and $A \cap B = \emptyset$.

Let us denote by $\mathcal{S}(\Omega, \Delta)$ the set of all states on (Ω, Δ) . We will often write simply $\mathcal{S}(\Delta)$ instead of $\mathcal{S}(\Omega, \Delta)$ if Ω is clear from the context or not needed in the reasoning. By a standard argument, $\mathcal{S}(\Delta)$ is easily seen to be a convex set compact in the pointwise topology. We will use this fact a few times in the sequel.

1.3. Definition. Let (Ω, Δ) be a SROMP and let $s \in \mathcal{S}(\Delta)$. We say that the state s is subadditive if the following condition is satisfied: If $A, B \in \Delta$, then there is a set $C \in \Delta$ such that $C \supset A \cup B$ and, moreover, $s(A) + s(B) \geq s(C)$.

If (Ω, Δ) is a Boolean algebra, then all states on Δ are subadditive (we simply take $A \cup B$ for C). In SROMPs, we typically have both subadditive and non-subadditive states. The following example well illustrates the situation.

1.4. Example. Let r be a positive rational number. Let $\Omega = \langle 0, r \rangle \times \langle 0, r \rangle$ and, further, let $\Delta = \{A \subset \Omega \mid A \text{ is a Lebesgue measurable set in } \Omega \text{ such that } \mu(A) \text{ is a rational number, where } \mu \text{ is the Lebesgue measure}\}$. Then a routine verification gives that (Ω, Δ) is a SROMP. Fix a set $B \in \Delta$ with $\mu(B) > 0$, and define a mapping $s_B: \Delta \rightarrow \langle 0, 1 \rangle$ by putting $s_B(A) = \frac{\mu(B \cap A)}{\mu(B)}$. Then s_B is a state on (Ω, Δ) . It can be shown that s_B is subadditive if, and only if, $\mu(B)$ is a rational number.

Let us now introduce the central notion of this paper.

1.5. Definition. Let (Ω, Δ) be a SROMP. If every state s on Ω, Δ is subadditive, then (Ω, Δ) is called *nearly Boolean*.

In what follows, we will analyse how “near” the nearly Boolean OMPs are to Boolean algebras.

2. RESULTS

The following theorem lists the hitherto known cases in which nearly Boolean implies Boolean (the statement (vi) is auxiliary). We provide sketches of all proofs for the sake of completeness, though some of the proofs can be extracted from [8] and [12]. (Let us recall that a maximal Boolean subalgebra of an OMP is called a block. Obviously, every Boolean subalgebra of an OMP can be enlarged to a block.)

2.1. Theorem. *Let (Ω, Δ) be a SROMP. Then the following statements hold true:*

- (i) *If Δ is a lattice, then Δ is Boolean.*
- (ii) *If Ω is countable, then Δ is Boolean.*
- (iii) *If Δ possesses only finite blocks, then Δ is Boolean.*
- (iv) *If Δ possesses only finitely many blocks, then Δ is Boolean.*
- (v) *If Δ is 2^{χ_0} -complete, then Δ is Boolean.*
- (vi) *If $A_1, A_2, \dots, A_n$ are sets from Δ , then there is a finite collection of sets from Δ , call them $B_1, B_2, \dots, B_m$, such that $\bigcap_{i \leq n} A_i = \bigcup_{j \leq m} B_j$. If, moreover, a positive ε is given, then the sets $B_1, B_2, \dots, B_m$ can be chosen so that for any state $s \in \mathcal{S}(\Delta)$ such that $s(A_i) = 1$ for each $i \leq n$ there is a B_j ($j \leq m$) such that $s(B_j) > 1 - \varepsilon$.*

Proof. (i) Take two sets $A, B \in \Delta$ and consider $A \wedge B$. We want to show that $A \wedge B = A \cap B$. Suppose that it is not the case. Then there is a point $p \in \Omega$ such that $p \in (A \cap B) - (A \wedge B)$. Consider the two-valued state s_p concentrated at the point p . Thus, $s_p(c) = 1$ if, and only if, $p \in c$. Then $s_p(\Omega - A) = s_p(\Omega - B) = 0$. Since s_p is supposed to be subadditive, there is a set $\tilde{C}$ such that $\tilde{C} \supset (\Omega - A) \cup (\Omega - B)$ and $s_p(\tilde{C}) \leq s_p(\Omega - A) + s_p(\Omega - B) = 0$. It follows that $s_p(\tilde{C}) = 0$. If we put $C = \Omega - \tilde{C}$, then $s_p(C) = 1$ and therefore $C \neq \emptyset$. Moreover, $C \subset A \cap B$ and $p \in C$. By the definition of infima, $C \subset A \wedge B$, and this is a contradiction. We have shown that $A \cap B \in \Delta$ for any $A, B \in \Delta$, and this means that (Ω, Δ) is Boolean.

(ii) Suppose that $A, B \in \Delta$ and $A \cap B \neq \emptyset$. Put $A \cap B = \{p_n | n \in N\}$ and

$$s = \sum_{n \in N} \frac{1}{2^n} s_{p_n},$$

where s_{p_n} are states concentrated at p_n . Obviously, $s(A) = s(B) = 1$, and the subadditivity of s guarantees a set C such that $C \in \Delta$, $C \subset A \cap B$ and $s(C) = 1$. But then $C = A \cap B$, and this is what we wanted to show.

For the proof of the remaining properties, let us first verify the condition (vi) of Th. 2.1.

(vi) Let us first observe a simple fact which we formulate as a lemma.

Lemma. *Suppose that (Ω, Δ) is a SROMP. Suppose that $\{A_1, A_2, \dots, A_n\} \subset \Delta$ and suppose further that $s(A_1) = s(A_2) = \dots = s(A_n) = 1$ for a subadditive state $s \in \mathcal{S}(\Omega, \Delta)$. Then there is a set B , $B \in \Delta$, such that $B \subset \bigcap_{i \leq n} A_i$ and $s(B) = 1$.*

Proof of the Lemma. We will first prove the lemma for two sets, A_1 and A_2 . Suppose that $s(A_1) = s(A_2) = 1$. Then $s(\Omega - A_1) = s(\Omega - A_2) = 0$. Since s is subadditive, there is a set $\tilde{B}$ such that $\tilde{B} \supset (\Omega - A_1) \cup (\Omega - A_2)$ and $s(\tilde{B}) \leq s(\Omega - A_1) + s(\Omega - A_2) = 0$. If we put $B_1 = \Omega - \tilde{B}$, then $B_1 \subset A_1 \cap A_2$ and $s(B_1) = 1$.

Let us now proceed inductively. Suppose that for A_1, A_2 we have the required set B_1 . Let us consider the sets B_1, A_3 , and find the set B_2 such that $B_2 \subset B_1 \cap A_3 \subset$

$A_1 \cap A_2 \cap A_3$ and $s(B_2) = 1$. Going on in this way, we finally take B_{n-1} for B , which establishes the lemma. $\square$

Let us return to the proof of Th. 2.1(vi). Suppose that $\{A_1, A_2, \dots, A_n\} \subset \Delta$. If $\bigcap_{i \leq n} A_i = \emptyset$, then there is nothing to be proved. Suppose therefore that $\bigcap_{i \leq n} A_i \neq \emptyset$. Put $\mathcal{S} = \{s \mid s \in \mathcal{S}(\Omega, \Delta) \text{ and } s(A_1) = s(A_2) = \dots = s(A_n) = 1\}$. Obviously, $\mathcal{S} \neq \emptyset$. Moreover, since $\mathcal{S}$ is a closed subset of the (compact) set $\mathcal{S}(\Omega, \Delta)$, the set $\mathcal{S}$ is also compact. Put $\mathcal{B} = \{B \mid B \in \Delta \text{ and } B \subset \bigcap_{i \leq n} A_i\}$ and set, for any $B \in \mathcal{B}$, $S_B = \{s \in \mathcal{S}(\Omega, \Delta) \mid s(B) > 0\}$. Every set S_B is obviously an open set in $\mathcal{S}$ and, by our lemma, $\mathcal{S} = \bigcup_{B \in \mathcal{B}} S_B$. This means that $\{S_B \mid B \in \mathcal{B}\}$ is an open covering of $\mathcal{S}$, and therefore there is a finite collection $B_1, B_2, \dots, B_m$ such that $\mathcal{S} = \bigcup_{i \leq m} S_{B_i}$. It remains to be shown that $\bigcup_{i \leq m} B_i = \bigcap_{i \leq n} A_i$. If this is not the case, then there is a point $p \in \bigcap_{i \leq n} A_i - \bigcup_{i \leq m} B_i$. Let us consider the state s_p concentrated at p . It follows that $s_p(A_i) = 1$ for any i , $i \leq n$, and therefore $s_p \in \mathcal{S}$. On the other hand, $s_p(B_i) = 0$ for any B_i ($i \leq m$), and this is absurd. (For the additional statement of Th. 2.1(vi), we simply take the sets $S_B^\varepsilon = \{s \in \mathcal{S} \mid s(B) > 1 - \varepsilon\}$ in the above proof instead of S_B .)

(iii) Let us suppose that (Ω, Δ) possesses only finite blocks. If $A, B \in \Delta$ and $A \cap B \neq \emptyset$, then there is a nonempty set $C_1 \in \Delta$ such that $C_1 \subset A \cap B$ (Th. 2.1(vi)). If $(A - C_1) \cap (B - C_1) = \emptyset$, we are done, for $A \cap B = \Delta$. If not, then there is a nonempty set C_2 such that

$$C_2 \subset (A - C_1) \cap (B - C_1).$$

If $(A - (C_1 \cup C_2)) \cap (B - (C_1 \cup C_2)) = \emptyset$, we are done. If not, then there is a nonempty set C_3 such that $C_3 \subset (A - (C_1 \cup C_2)) \cap (B - (C_1 \cup C_2))$, etc. If $\bigcup_{i \leq n} C_i = A \cap B$ for some $n \in \mathbb{N}$, we are done. If not, then there is a countable collection $\{C_i \mid i \in \mathbb{N}\}$ of mutually disjoint sets which belong to Δ . But then the Boolean algebra generated by $\{C_i \mid i \in \mathbb{N}\}$ in Δ is infinite, which is a contradiction. Thus, (Ω, Δ) must be Boolean.

(iv) Suppose that (Ω, Δ) possesses exactly n maximal Boolean subalgebras. Take any two sets $A_1, A_2 \subset \Delta$, and consider Th. 2.1 (vi) for $\varepsilon = \frac{1}{n}$. We see that there is a collection $\{B_1, B_2, \dots, B_m\} \subset \Delta$ such that $\bigcup_{j \leq m} B_j = A_1 \cap A_2$ and, moreover, for any state $s \in \mathcal{S}(\Delta)$ with $s(A_1) = s(A_2) = 1$ there is B_j such that $s(B_j) > 1 - \frac{1}{n}$. Since every B_j ($j \leq m$) must belong to a maximal Boolean subalgebra of Δ , we may (and shall) assume that $m \leq n$ (otherwise we simply take the unions of those B_j which belong to the same maximal Boolean subalgebra of (Ω, Δ)). We may further assume that $B_j \neq A_1 \cap A_2$ for any j ($j \leq m$). This means that there are points $p_j \in \Omega$ ($j \leq m$) such that, for any $j \leq m$, $p_j \in (A_1 \cap A_2) - B_j$. Put $s = \sum_{j \leq m} \frac{1}{m} s_{p_j}$ (s_{p_j} is the state concentrated at p_j). Then $s(B_j) \leq 1 - \frac{1}{m} \leq 1 - \frac{1}{n}$ for any $j \leq m$, and this is a contradiction. It follows that (Ω, Δ) is a Boolean algebra.

(v) This part strengthens a result of the paper [6], where the author proves the same fact for complete SROMPs. We have not been able to prove (or disprove) the same statement for σ -complete SROMPs (this problem has already been raised in [6]).

Let (Ω, Δ) be a $2^{\aleph_0}$ -complete SROMP. In other words, let (Ω, Δ) be closed under the formation of the unions of the pairwise disjoint collections of elements from Δ consisting of at most $2^{\aleph_0}$ elements. Take arbitrary sets $A, B \in \Delta$ and find the collection $B_1, B_2, \dots, B_m \in \Delta$ such that $\bigcup_{j \leq m} B_j = A \cap B$ (Th. 2.1 (vi)). Put

$\mathcal{B}_1 = \{B_j | j \leq m\}$. We will show that $\bigcup_{j \leq m} B_j \in \Delta$, proving that $A \cap B \in \Delta$. This will mean that (Ω, Δ) is Boolean.

Form the coarsest partition, $\mathcal{P}_1$, of $A \cap B$ that refines the covering $\mathcal{B}_1$ of the set $A \cap B$. Thus, $\mathcal{P}_1$ consists of the sets $\bigcap_{j \leq m} \tilde{B}_j$, where $\tilde{B}_j$ is either B_j or $\Omega - B_j$. Write $\mathcal{P}_1 = \{C_k | k \leq r\}$. By Th. 2.1 (vi), every C_k can be expressed as a union of the type $\bigcup_{p \leq s} B_p^k$ for some $s \in N$, where $B_p^k \in \Delta$ for any $p, p \leq s$. Obviously,

$$\bigcup_{\substack{k \leq r \\ p \leq s}} B_p^k = A \cap B.$$

Put $\mathcal{B}_2 = \{B_p^k | k \leq r, p \leq s\}$ and denote by $\mathcal{P}_2$ the coarsest partition of $A \cap B$ that refines $\mathcal{B}_2$. Going on in this way, we obtain a countable collection $\mathcal{B}_1, \mathcal{B}_2, \dots$ of coverings of $A \cap B$ with the following property. If $P \in \mathcal{B}_u$ ($u \in N$) and $Q \in \mathcal{B}_v$ ($v \in N$) for $u < v$, then the following implication holds true: If $P \cap Q \neq \emptyset$, then $P \subset Q$. It can be easily seen that the sets of the type $P_1 \cap P_2 \cap P_3 \cap \dots$, where $P_i \in \mathcal{B}_i$, belong to Δ (here even σ -completeness would suffice) and, moreover, these sets constitute a partition of $A \cap B$ into not more than $2^{\aleph_0}$ sets. This means that $A \cap B \in \Delta$, and the proof is complete. $\square$

The previous results indicate that if “properly” nearly Boolean OMP exist, they must be rather complicated. It therefore seems probable that the following procedure for constructing non-Boolean nearly Boolean OMP can hardly be simplified. The following result, which generalizes step VI in the proof of Theorem 1 in [6], provides an essential argument.

2.2. Proposition. *Let (Ω, Δ) be a SROMP, and suppose that for any pair $A, B \in \Delta$ there exist an uncountable set I and a collection $C_\alpha, D_\alpha \in \Delta$ ($\alpha \in I$) such that the following two conditions are satisfied:*

- (i) *For any $\alpha \in I$, $C_\alpha \cup D_\alpha = A \cap B$.*
- (ii) *If $\alpha \neq \beta$, then $C_\alpha \cap C_\beta = \emptyset$.*

Then (Ω, Δ) is nearly Boolean.

Proof. Let $s \in \mathcal{S}(\Delta)$ be a state on Δ . We must show that s is subadditive. Suppose that $A, B \in \Delta$. We have to exhibit a set $D, D \in \Delta$, such that $D \supset A \cup B$ and $s(D) \leq s(A) + s(B)$. Let $\tilde{A} = \Omega - A$ and $\tilde{B} = \Omega - B$. Let C_α, D_α ($\alpha \in I$) be the collection ensured by the assumption of Prop. 2.2 taken for $\tilde{A}$ and $\tilde{B}$. It can be seen easily that there is an index $\alpha_0 \in I$ such that $s(C_{\alpha_0}) = 0$. Indeed, if $s(C_\alpha) > 0$ for any $\alpha \in I$, then there is an infinite set $J, J \subset I$, such that $s(C_j) > \frac{1}{n}$ for some $n \in N$. Since the collection C_α ($\alpha \in I$) consists of mutually disjoint sets, we reach a contradiction with the additivity of s . Consider now the set C_α . Since $C_{\alpha_0} \cup D_{\alpha_0} = \tilde{A} \cap \tilde{B}$, we see that $(\tilde{A} - C_{\alpha_0}) \cap (\tilde{B} - D_{\alpha_0}) = \emptyset$. It follows that $s(\tilde{A} - C_{\alpha_0}) + s(\tilde{B} - D_{\alpha_0}) \leq 1$ and therefore $s(\tilde{A}) - s(C_{\alpha_0}) + s(\tilde{B}) - s(D_{\alpha_0}) \leq 1$. Since $s(C_{\alpha_0}) = 0$, we obtain $s(\tilde{A}) + s(\tilde{B}) \leq 1 + s(D_{\alpha_0})$. Put $D = \Omega - D_{\alpha_0}$. Then $D \supset A \cup B$, because $D_\alpha \subset \tilde{A} \cap \tilde{B}$. Also, $D_\alpha = \Omega - D$ and $\tilde{A} = \Omega - A, \tilde{B} = \Omega - B$. Substituting into the inequality $s(\tilde{A}) + s(\tilde{B}) \leq 1 + s(D_\alpha)$, we have $s(\Omega - A) + s(\Omega - B) \leq 1 + s(\Omega - D)$. This yields $1 - s(A) + 1 - s(B) \leq 1 + 1 - s(D)$, which gives the inequality $s(D) \geq s(A) + s(B)$. The proof of Prop 2.2 is complete. $\square$

Before we take up the construction of the desired example, let us recall the following simple but useful fact (see [7] for the definition and other considerations).

2.3. Proposition. *Let (Ω, Σ) be a Boolean algebra of subsets of Ω . Let G be a group. Let $m: \Sigma \rightarrow G$ be a G -valued measure (i.e., $m(\emptyset) = 0$, and $m(A \cup B) = m(A) + m(B)$ whenever $A \cap B = \emptyset$). Then if H is a subgroup of G such that $m(\Omega) \in H$, then the set $\Delta = \{A \in \Sigma \mid m(S) \in H\}$ is a SROMP.*

Proof. Obviously, $\emptyset \in \Delta$. If $A \in \Delta$, then $m(\Omega - A) = m(\Omega) - m(A) \in H$. Thus, $\Omega - A \in \Delta$. Finally, if $A, B \in \Delta$ and $A \cap B = \emptyset$, then $m(A \cup B) = m(A) + m(B) \in H$. This means that $A \cup B \in \Delta$. The proof is complete. $\square$

We are ready to present the example of a non-Boolean nearly Boolean OMP. We omit some technical details, assuming the reader to be familiar with the standard construction of Boolean field product and basic extension technique for measures (see e.g. [5] and [15]).

2.4. Theorem. *There is a non-Boolean nearly Boolean OMP.*

Proof. Let P, Q be two disjoint uncountable sets of equal cardinality. Put $S = P \cup Q$ and consider the Boolean algebra $\tilde{B} = (S, \tilde{\Sigma})$ where $\tilde{\Sigma}$ consists of finite-cofinite subsets of S . Let Z denote the group of whole numbers, and let $\tilde{m}_1$ (resp. $\tilde{m}_2$) be the Z -valued measure $\tilde{m}_1: \tilde{\Sigma} \rightarrow Z$ (resp., $\tilde{m}_2: \tilde{\Sigma} \rightarrow Z$) which is determined as follows: $\tilde{m}_1(\{x\}) = 1$ if $x \in P$, $\tilde{m}_1(\{x\}) = 0$ if $x \in Q$ and $\tilde{m}_1(S) = 1$ (resp., $\tilde{m}_2(\{x\}) = 0$ if $x \in P$, $\tilde{m}_2(\{x\}) = 1$ if $x \in Q$ and $\tilde{m}_2(S) = 1$). Now, take countably many copies of the Boolean algebra $\tilde{B}$ and form the corresponding countable Boolean field product B . Thus, $B = (T, \Sigma)$, where T denotes the cartesian product of countably many copies of the set S and Σ consists of finite disjoint unions of “generalized rectangles” in T (by a generalized rectangle in T we mean a set of the type

$$A_1 \times A_2 \times \cdots \times A_n \times \prod_{j \geq n+1} S_j,$$

where $A_j \in \tilde{\Sigma}$ for $j \leq n$ and $S_j = S$ for $j \geq n+1$). Since $\tilde{m}_1(S) = 1$ (resp., $\tilde{m}_2(S) = 1$), we can extend $\tilde{m}_1$ (resp., $\tilde{m}_2$) over the entire Σ , obtaining measures m_1 and m_2 on Σ . (Indeed, if $V = A_1 \times A_2 \times \cdots \times A_n \times \prod_{j \geq n+1} S_j$ is a generalized rectangle, we put

$$m_1(V) = \prod_{i \leq n} \tilde{m}_1(A_i), \quad m_2(V) = \prod_{i \leq n} \tilde{m}_2(A_i)$$

and continue in the standard way - see e. g. [15].) Let $m = m_1 + m_2$. Obviously, m is a measure, $m: \Sigma \rightarrow Z$, and $m(T) = 2$. Let us denote by Z_2 the group of even numbers considered as a subgroup of Z . Let $\Delta = \{A \in \Sigma \mid m(A) \in Z_2\}$. Then (T, Δ) is a SROMP (Prop. 2.3). We claim that (T, Δ) is a non-Boolean nearly Boolean OMP.

First, let us show that (T, Δ) is not Boolean. Pick a point $x \in P$ and two distinct points $y_1, y_2 \in Q$. Let $V_1 = \{x, y_1\} \times \prod_{j \geq 2} S_j$, $V_2 = \{x, y_2\} \times \prod_{j \geq 2} S_j$ ($S_j = S$ for $j \geq 2$). Then $m(V_1) = m(V_2) = 2$, and therefore $A, B \in \Delta$. On the other hand, $m(V_1 \cap V_2) = m(\{x\} \times \prod_{j \geq 2} S_j) = 1$, and therefore $V_1 \cap V_2 \notin \Delta$. Thus, (T, Δ) is not Boolean.

Let us prove that (T, Δ) is nearly Boolean. We will verify that (T, Δ) satisfies the assumption of Prop. 2.2. Suppose that $A, B \in \Delta$. If $A \cap B \in \Delta$, then we put $C_\alpha = \emptyset$, $D_\alpha = A \cap B$, the index α formally varying over an uncountable set. Suppose that $A \cap B \notin \Delta$. Then $m(A \cap B)$ is an odd number. We have defined $m = m_1 + m_2$, and we also know that $A \cap B$ can be expressed as a disjoint union

of rectangles, $A \cap B = \bigcup_{j \leq n} R_j$. Since there is a rectangle, R_{j_0} , for which either $m_1(R_{j_0}) \neq 0$ or $m_2(R_{j_0}) \neq 0$, we see that we can find points $p_1, p_2, \dots, p_m$ such that the set $V = \{p_1\} \times \{p_2\} \times \dots \times \{p_m\} \times \prod_{j \geq m+1} S_j$ ($S_j = S$ for $j \geq m+1$) is a subset of $A \cap B$ and, moreover, either $m_1(V) = 1$ or $m_2(V) = 1$. Observe that if $m_1(V) = 1$ then $m_2(V) = 0$, and if $m_2(V) = 1$ then $m_1(V) = 0$.

Suppose that $m_1(V) = 1$ and $m_2(V) = 0$ (the other case can be argued similarly). Consider the set S_{m+1} ($S_{m+1} = S$) in the above expression of the set V . Take the partition $\mathcal{P}$, $\mathcal{P} = \{E_\alpha | \alpha \in I\}$, of S_{m+1} such that the index set I is the set of quadruples of elements of S_{m+1} and the sets E_α are defined as follows: Every E_α consists of four points, $E_\alpha = \{x_1, x_2, x_3, x_4\}$, where $x_1, x_2 \in P$ and $x_3, x_4 \in Q$. For any $E_\alpha = \{x_1, x_2, x_3, x_4\}$, let us set

$$C_\alpha = \{p_1\} \times \{p_2\} \times \dots \times \{p_m\} \times E_\alpha \times \prod_{j \geq n+2} S_j$$

and

$$D_\alpha = A \cap B - (\{p_1\} \times \{p_2\} \times \dots \times \{p_m\} \times \{x_1\} \times \prod_{j \geq n+2} S_j).$$

Obviously, $C_\alpha \cup D_\alpha = A \cap B$ for any $\alpha \in I$, and all sets C_α belong to Δ . Since the number $m(D_\alpha)$ can be expressed as a difference of two odd numbers, we see that $m(D_\alpha)$ is an even number and therefore $D_\alpha \in \Delta$ for any $\alpha \in I$. The proof is complete. $\square$

The above result can be improved by ensuring an arbitrary degree of “absolute compatibility” in nearly Boolean OMPs. In other words, it can be proved that there are nearly Boolean OMPs with arbitrary centres (for the notion of compatibility and centre in OMP, see e.g. [13]). This might be relevant to the logico-algebraic foundations of quantum theories. Also, it establishes a proper class of nearly Boolean OMPs. To keep the exposition short, we assume that the reader is acquainted with the construction of a Boolean power of OMPs (see e.g. [9] and [2]).

2.5. Proposition. *Let $L = (\Omega, \Delta)$ be a nearly Boolean OMP. Let B be a Boolean algebra. Then the Boolean power, P , of B and L is a nearly Boolean OMP. Moreover, if $C(\Delta)$ denotes the centre of Δ and if $C(\Delta) = \{0, 1\}$, then $C(P) = B$.*

Proof. This immediately follows from the description of the states on P (see e.g. [10]) and from compatibility properties of a Boolean power (see e.g. [9] and [2]). $\square$

2.6. Theorem. *Suppose that B is a Boolean algebra. Then there is a non-Boolean nearly Boolean OMP, L , such that $C(L) = B$.*

Proof. Observe that the nearly Boolean OMP constructed in Th. 2.3 has a centre consisting only of $\{0, 1\}$. The rest follows from Prop. 2.4. $\square$

Let us conclude this paper by formulating an open question that arises naturally from Th. 2.3. Can every SROMP be faithfully embedded in a nearly Boolean OMP? (An OMP embedding $e: L \rightarrow K$ is called faithful if a is compatible with b in L exactly when $e(a)$ is compatible with $e(b)$ in K .) A positive answer to this question could have a bearing on quantum axiomatics. It should be noted that the σ -additive version of this question has been positively answered in [3].

REFERENCES

- [1] Beltrametti, E., Cassinelli, G., *The logic of quantum mechanics*, Addison-Wesley, Reading, Massachusetts, 1981. MR **83d**:81008
- [2] Bruns, G., Greechie, R.J., Harding, J., Roddy, M., *Completions of orthomodular lattices*, Order **7** (1990), 67–76. MR **92b**:06029
- [3] DeLucia, P., Pták, P., *Quantum probability spaces that are nearly classical*, Bull. Polish Acad. Sciences – Math. **40** (2) (1992), 163–173. MR **97i**:81015
- [4] Gudder, S., *Stochastic Methods in Quantum Mechanics*, North-Holland, Amsterdam, 1979. MR **84j**:81003
- [5] Halmos, P., *Measure Theory*, Van Nostrand, New York, 1950. MR **11**:504d
- [6] Müller, V., *Jauch-Piron states on concrete quantum logics*, Int. Journ. Theor. Phys. **32** (3) (1993), 433–442. MR **94g**:81011
- [7] Navara, M., *Kernel logics*, Czechoslovak Math. J., (to appear).
- [8] Navara, M., Pták, P., *Almost Boolean orthomodular posets*, Jour. Pure Applied Algebra **60** (1989), 105–111. MR **90h**:06003
- [9] Pták, P., *Logics with given centers and state spaces*, Proc. Amer. Math. Soc. **88** (1983), 106–109. MR **84f**:06016
- [10] Pták, P., *Summing of Boolean algebras and logics*, Demonstratio Math. **19** (1986), 349–357. MR **89b**:03104
- [11] Pták, P., *Jauch-Piron property (everywhere!) in the logico-algebraic foundation of quantum theories*, Int. Journ. Theor. Phys. **32** (10) (1993), 1985–1991. MR **94j**:81024
- [12] Müller, V., Pták, P., Tkadlec, J., *Concrete quantum logics with covering properties*, Int. Journ. Theor. Phys. **31** (5) (1992), 843–854. MR **93a**:81021
- [13] Pták, P., Pulmannová, S., *Orthomodular Structures as Quantum Logics*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1991. MR **94d**:81018b
- [14] Pták, P., Pulmannová, S., *A measure theoretic characterization of Boolean algebras among orthomodular lattices*, Comment. Math. Univ. Carolinae **35** (1) (1994), 205–208. MR **95i**:06014
- [15] Sikorski, R., *Boolean Algebras*, Springer Verlag, Berlin-Heidelberg-New York, 1969. MR **39**:4053
- [16] Varadarajan, V., *Geometry of Quantum Theory I*, Van Nostrand, Princeton, 1968. MR **57**:11399

CZECH TECHNICAL UNIVERSITY, FACULTY OF ELECTRICAL ENGINEERING, DEPARTMENT OF MATHEMATICS, 16627 PRAGUE 6, CZECH REPUBLIC

E-mail address: ptak@math.feld.cvut.cz