A NEW PROOF OF THE SOLOMON-TITS THEOREM

CHAO KU

(Communicated by Ronald M. Solomon)

ABSTRACT. We give a new proof of the Solomon-Tits Theorem which asserts that the Tits building of a finite group of Lie type has the homotopy type of a bouquet of spheres.

1. INTRODUCTION

Let G be a finite group of Lie type of rank $n \geq 3$, (G, B, N, S) a Tits system for G , $H = B \cap N$, $B = UH$ where $U = O_p(B)$ and p is the defining prime for G . $\mathbf{F} = \{G_i; 1 \leq i \leq n\}$ is the set of maximal parabolics of G over B and $\mathcal{B} = \mathcal{C}(G, \mathbf{F})$ the simplicial complex whose vertex set is the union of coset spaces G/G_i , $i \in I$, and whose simplices are collection of vertices with nonempty intersection. $\mathcal{B}$ is called the Tits building of G . The following theorem is well known:

Theorem (Solomon-Tits). *The Tits building $\mathcal{B}$ has the homotopy type of a wedge of $|U|$ $(n-1)$ -spheres.*

In his seminar paper [S] Solomon first mentioned the theorem and sketched a proof. Later Garland ([G]) gave a proof which is essentially the same as Solomon's, and Curtis and Lehrer ([CL]) gave a proof of a related result based on the computation of the $\mathbb{Q}G$ -endomorphism algebra of $H_*(\mathcal{B})$. We provide a new proof here which should help to better understand the structure of the Tits building. Namely, $\mathcal{B}$ is the union of the $|U|$ apartments containing a fixed chamber, each having the homotopy type of an $(n-1)$ -sphere as it is a Coxeter complex. We show that the intersection of any k ($k \geq 2$) of these subcomplexes is contractible. Then it follows from an easy lemma in topology that $\mathcal{B}$ has the homotopy type of a wedge of these spheres.

2. PRELIMINARY LEMMAS

Let us begin by recording a well known lemma from topology.

Lemma 1. *Suppose X is a simplicial complex and $X_1, X_2, \dots, X_m$, $m > 1$, are subcomplexes of X with $X = \bigcup_{1 \leq i \leq m} X_i$ such that for any $1 \leq i_1 < i_2 < \dots < i_k \leq m$, $k > 1$, the set $\bigcap_{1 \leq j \leq k} X_{i_j}$ is contractible. Then X has the homotopy type of $X_1 \vee X_2 \vee \dots \vee X_m$.*

Received by the editors December 19, 1996.

1991 *Mathematics Subject Classification.* Primary 20E42.

©1998 American Mathematical Society

Proof: By induction. Note that if B is a contractible subcomplex of a simplicial complex A , then the quotient map $q : A \rightarrow A/B$ is a homotopy equivalence. If $m = 2$, as $Y = X_1 \cap X_2$ is contractible, $X/Y = X_1/Y \cup X_2/Y$ and $X_1/Y \cap X_2/Y = Y/Y$ is a point, the lemma follows easily in this case.

Assume the lemma holds for $< m$. Then $X_1 \cap (\bigcup_{i>1} X_i) = \bigcup_{i>1} (X_1 \cap X_i)$ is homotopy equivalent to $\bigvee_{i>1} (X_1 \cap X_i)$ by induction and hence contractible. Therefore X is homotopy equivalent to $X_1 \vee (\bigcup_{i>1} X_i)$, which is in turn homotopy equivalent to $X_1 \vee X_2 \vee \cdots \vee X_m$. $\square$

We refer readers who are unfamiliar with the geometry of groups of Lie type and the theory of Coxeter groups to [A] or [B]. From time to time by abuse of notation we identify an element of N with its image in the Weyl group $W = N/H$. We may also use the same notation for a simplex of a complex and the subcomplex consisting of its faces.

Lemma 2. *Let $W_J = G_J \cap N/H$ be a parabolic subgroup of W , where $G_J = \bigcap_{j \in J \subseteq I} G_i$. Then each coset $W_J d$ has a unique element d_J of minimal length such that $l(wd_J) = l(w) + l(d_J)$ for all $w \in W_J$. Here l is the length function on W with respect to S .*

Proof. This is exercise 3, page 37 in [B]. $\square$

Define a partial ordering $\leq$ on W by $v \leq w$ if and only if $w = uv$ and $l(w) = l(u) + l(v)$ for some $u \in W$. This is a well defined ordering, and w_0 , the longest element of W , is the unique maximal element with respect to this order. See for instance exercise 6, page 155 in [A].

Lemma 3. *Let Φ be the Coxeter complex (see page 210 in [A] for the definition) for W , $\mathbf{C} \in \Phi$ the fundamental chamber and $\sigma \in \Phi$ a simplex. Then there is a unique $w_\sigma \in W$ with $\sigma \subseteq \mathbf{C}w_\sigma$ such that for any chamber $\mathbf{C}w$ containing σ , $w_\sigma \leq w$. Moreover, there is a sequence of chambers $\mathbf{C}w_\sigma = \mathbf{C}w_1, \mathbf{C}w_2, \dots, \mathbf{C}w_n = \mathbf{C}w$, each containing σ , such that $w_i \leq w_{i+1}$ with $l(w_{i+1}) = l(w_i) + 1$, $1 \leq i < n$.*

Proof. This is a direct consequence of Lemma 2. $\square$

3. PROOF OF THE THEOREM

Let $\mathbf{C} = \{G_i; 1 \leq i \leq n\}$ be the fundamental chamber of $\mathcal{B}$ and $\Sigma = \bigcup_{w \in W} \mathbf{C}w$. Then Σ is an apartment stabilized by N isomorphic to the Coxeter complex Φ . Hence, as Φ has the homotopy type of a $(n-1)$ -sphere,

Proposition 1. Σ has the homotopy type of a $(n-1)$ -sphere.

As $G = BNB = BNU$, it is obvious that

Proposition 2. $\mathcal{B} = \bigcup_{u \in U} \Sigma^u$, where $\Sigma^u = \bigcup_{w \in W} \mathbf{C}wu$.

Notice that for $u_i \in U$, $1 \leq i \leq k$, $\bigcap_{1 \leq i \leq k} \Sigma^{u_i}$ is isomorphic to

$$\left(\bigcap_{1 \leq i \leq k} \Sigma^{u_i} \right)^{u_1^{-1}} = \bigcap_{1 \leq i \leq k} \Sigma^{u_i u_1^{-1}}.$$

So by the argument in the introduction, a proof of the following proposition will complete the proof of the Solomon-Tits Theorem.

Proposition 3. For distinct $u_1, u_2, \dots, u_k \in U - \{1\}$, $k \geq 1$, $\Sigma \cap \bigcap \Sigma^{u_k}$ is contractible.

The rest of paper is dedicated to the proof of Proposition 3. For simplicity, we deal only with the case $k = 1$. The same proof works for $k > 1$.

Lemma 4. (1) For $1 \neq u \in U$, $\mathcal{T} = \Sigma \cap \Sigma^u$ is the subcomplex of Σ consisting of simplices fixed by u . Moreover, $\mathcal{T}$ is a union of chambers.
 (2) If $Cw \in \mathcal{T}$ and $v \leq w$, then $Cv \in \mathcal{T}$.

Proof. Let $\mathbf{C}_J = \{G_j; j \in J \subseteq I\} \subseteq \mathbf{C}$ and assume $\mathbf{C}_J v_1 u = \mathbf{C}_J v_2$ for some $v_1, v_2 \in W$. Then $G_J v_1 u = G_J v_2$. As $N \cap G_J = N_J$ is the preimage of W_J in N , we may choose v_1 and v_2 as in Lemma 2. Let $l(v_k) = r_k$, $k = 1, 2$. Without loss assume $r_1 \leq r_2$. Otherwise we consider u^{-1} instead. Set $v_1 = x_1 x_2 \dots x_{r_1}$ with x_k 's from S . As $sBw \subseteq BwB \cup BswB$ for $s \in S$ and $w \in W$, we have

$$v_1 u v_2^{-1} \in B v_2^{-1} B \cup \bigcup_{1 \leq k_1 < k_2 < \dots < k_t \leq r_1} B x_{k_1} x_{k_2} \dots x_{k_t} v_2^{-1} B.$$

Then as $v_1 u v_2^{-1} \in G_J = B N_J B$, either $v_1 u v_2^{-1} \in B v_2^{-1} B \subseteq G_J$ or there are $1 \leq k_1 < k_2 < \dots < k_t \leq r_1$ such that $v_1 u v_2^{-1} \in B x_{k_1} \dots x_{k_t} v_2^{-1} B \subseteq G_J$. In the first case, $v_2 \in G_J$ and by Lemma 2 $v_2 = 1$. Consequently $v_1 \in G_J v_2 u^{-1} = G_J$, and again by Lemma 2 $v_1 = 1$. In the second case, set $v_0 = x_{k_1} \dots x_{k_t}$. Then $v_0 \in G_J v_2$ with $l(v_0) \leq t \leq r_1 \leq r_2 = l(v_2)$; hence by Lemma 2, all equalities must hold and $t = r_1$, $r_1 = r_2$, $v_0 = v_1 = v_2$. Therefore $v_1 u v_2^{-1} \in B v_0 v_2^{-1} B = B$, i.e. $u \in B \cap B^{v_1}$. In either case, $v_1 = v_2$, $u \in B \cap B^{v_1}$ and $\mathbf{C}_J v_1 \subseteq \mathbf{C}_{v_1} \subseteq \mathcal{T}$. Part (1) is proved.

For $v \leq w \in W$ and $Cw \in \mathcal{T}$, assume $w = xv$ with $l(w) = l(x) + l(v)$. Let $x = x_1 x_2 \dots x_r$ with $r = l(x)$ and $x_i \in S$. For $t \leq r$, set $w_t = x_t \dots x_r v$. Now for any $s \in S$ and any $w \in W$ with $sw \leq w$, we have $B \cap B^w \leq B^{sw}$. Applying the observation to x_1 and w , we have $u \in B \cap B^w \in B^{w_2}$, so $Cw_2 \in \mathcal{T}$. Now part (2) follows inductively. $\square$

Lemma 5. $\mathcal{T}$ is contractible.

Proof. As $\mathbf{C} \in \mathcal{T}$, $\mathcal{T}$ is nonempty. $\mathcal{T}$ is connected by Lemma 3 and Lemma 4.2. Recall that w_0 is the longest element in W . Notice that $Cw_0 \notin \mathcal{T}$, as $u \notin B \cap B^{w_0} = H$.

Pick a $\bar{w} \in W$ with $\mathbf{C}\bar{w} \subseteq \mathcal{T}$ such that $l(\bar{w})$ is maximal. Set

$$\mathcal{T}_1 = \bigcup_{w \neq \bar{w}, u \in B \cap B^w} \mathbf{C}w \subseteq \mathcal{T}.$$

For a simplex $\sigma \in \mathcal{T}_1 \cap \mathbf{C}\bar{w}$, choose $v \in W$ as in Lemma 3 such that $\sigma \in \mathbf{C}v$. Then by Lemma 3, $v \leq \bar{w}$. So $\mathbf{C}v \in \mathcal{T}$ by Lemma 4.2. But $\sigma \in \mathcal{T}_1$, and consequently there is a $w \in W$ with $\sigma \in \mathbf{C}w \in \mathcal{T}_1$. This forces $v \neq \bar{w}$ and $\mathbf{C}v \in \mathcal{T}_1$. Therefore by Lemma 3 there is an $s \in S$ with $v \leq s\bar{w} \leq \bar{w}$ and $\sigma \in \mathbf{C}s\bar{w} \subseteq \mathcal{T}_1$. So $\sigma \subseteq \mathbf{C}\bar{w} \cap \mathbf{C}s\bar{w}$, an $(n-2)$ -simplex of $\mathbf{C}\bar{w}$.

On the other hand, for any $s \in S$ with $s\bar{w} \leq \bar{w}$, it is obvious that $\mathbf{C}s\bar{w} \in \mathcal{T}_1$ and $\mathbf{C}\bar{w} \cap \mathbf{C}s\bar{w} \subseteq \mathcal{T}_1 \cap \mathbf{C}\bar{w}$. We have shown that $\mathcal{T}_1 \cap \mathbf{C}\bar{w}$ is the union of $(n-2)$ -simplices $\mathbf{C}\bar{w} \cap \mathbf{C}s\bar{w}$ of $\mathbf{C}\bar{w}$ with $s\bar{w} \leq \bar{w}$. As $\bar{w} \neq w_0$, there is an $s \in S$ with $s\bar{w} > \bar{w}$. Therefore there are at most $n-1$ $(n-2)$ -simplices of $\mathbf{C}\bar{w}$ in $\mathcal{T}_1$, and their intersection contains a vertex; hence $\mathcal{T}_1 \cap \mathbf{C}\bar{w}$ is contractible. Applying Lemma 1, we conclude that $\mathcal{T}$ is homotopy equivalent to $\mathcal{T}_1$.

Notice that $\mathcal{T}_1$ is also a union of chambers and satisfies Lemma 3, part (2). Therefore, repeating the above procedure gives that $\mathcal{T}$ is homotopy equivalent to $\mathbf{C}$, the fundamental chamber, hence contractible. $\square$

REFERENCES

- [A] M. Aschbacher, *Finite Group Theory*, Cambridge University Press, Cambridge, 1986. MR **95b**:20002
- [B] Bourbaki, N., *Groupes et algebras de Lie*, 4, 5, 6, Masson, Paris, 1981. MR **83g**:17001
- [CL] C.W. Curtis and G. I. Lehrer, A new proof of a theorem of Solomon-Tits, *Proc. of AMS* v. 82, No. 2 (1982) 154-156. MR **83j**:20048
- [G] H. Garland, p -adic curvature and the cohomology of discrete subgroups of p -adic groups, *Ann. Math.(2)* 97 (1973) 375-393. MR **47**:8719
- [S] L. Solomon, The Steinberg character of a finite group with a BN -pair, *Theory of Finite Groups*, R. Brauer and C.H. Sah, Eds., Benjamin, New York, 1969, pp. 213-221. MR **40**:220

DEPARTMENT OF MATHEMATICS, CALTECH, PASADENA, CALIFORNIA 91125
E-mail address: `chaoku@cco.caltech.edu`