C^* -ALGEBRAS THAT ARE ONLY WEAKLY SEMIPROJECTIVE

TERRY A. LORING

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We show that the C^* -algebra of continuous functions on the Cantor set is a weakly semiprojective C^* -algebra that is not semiprojective.

The two most basic forms of semiprojectivity, both equivalent to useful notions of stability for C^* -algebra relations, are called semiprojectivity and weak semiprojectivity. A C^* -algebra A is *weakly semiprojective* if we can always solve the $*$ -homomorphism lifting problem

$$\begin{array}{ccc} & \prod_N B_n & (b_N, b_{N+1}, \dots) \\ & \downarrow \rho_N & \downarrow \\ A \xrightarrow{\varphi} \prod_1^\infty B_n / \bigoplus_1^\infty B_n & & [(0, \dots, 0, b_N, b_{N+1}, \dots)] \end{array}$$

and it is *semiprojective* if we can always solve the lifting problem

$$\begin{array}{ccc} & B / J_N & \\ & \downarrow & \\ A \xrightarrow{\varphi} B / \bigcup_1^\infty J_n & & (J_1 \triangleleft J_2 \triangleleft \dots \triangleleft B). \end{array}$$

If A is finitely generated, we have an advantage proving weak semiprojectivity, as it suffices to prove that the lifting can be done only to produce a $*$ -homomorphism $\bar{\varphi}$ so that

$$\|\rho_N \circ \bar{\varphi}(g) - \varphi(g)\| \leq \epsilon \quad (\forall g \in G)$$

for preordained positive ϵ and finite subset G . (See [9] or [3].)

What has been lacking is an example of a C^* -algebra that is weakly semiprojective but not semiprojective. With the Cantor set

$$X = \prod_1^\infty \{0, 1\}, \quad X = \varprojlim X_j, \quad X_j = \prod_1^j \{0, 1\},$$

we get such an example, $C(X)$. It is not semiprojective because its spectrum X is not an ANR, as is necessary for a commutative C^* -algebra to be semiprojective. It

Received by the editors November 19, 1996 and, in revised form, February 4, 1997.

1991 *Mathematics Subject Classification*. Primary 46L05.

Key words and phrases. Stable relations, semiprojectivity.

The research summarized here was supported, in part, by the National Science Foundation, DMS-9531841.

is weakly semiprojective by the following theorem, the well-known semiprojectivity of finite-dimensional C^* -algebras, and the existence of the $*$ -homomorphisms

$$\theta_j : C(X) \longrightarrow C(X_j) \quad \gamma_j : C(X_j) \longrightarrow C(X)$$

of truncation and zero-padding, whose composition $\gamma_j \circ \theta_j$ converges pointwise to the identity. Although $C(X)$ is the direct limit of the $C(X_j)$, this is not very relevant. The point is that $C(X)$ is an “approximate retract” of $C(X_j)$. A retract of a semiprojective is semiprojective ([1]), so it is not surprising that we can conclude that $C(X)$ is weakly semiprojective.

The proof of Theorem 0.1 was inspired by similar arguments by Lin. See any of the references for interesting results regarding semiprojectivity and stable relations.

Theorem 0.1. *Suppose that $A_1, A_2, \dots$ are semiprojective C^* -algebras and that A is a finitely generated C^* -algebra. If there are $*$ -homomorphisms $\theta_j : A \rightarrow A_j$ and $\gamma_j : A_j \rightarrow A$ such that*

$$\|\gamma_j \circ \theta_j(a) - a\| \rightarrow 0$$

for all a in A , then A is weakly semiprojective.

Proof. Let $\epsilon > 0$ and a finite set $G \subset A$ be given, as well as a $*$ -homomorphism of the form

$$\varphi : A \longrightarrow \prod_1^\infty B_n \bigg/ \bigoplus_1^\infty B_n,$$

which we now must lift, approximately. Fix j such that $\|\gamma_j \circ \theta_j(g) - g\| \leq \epsilon$ for all g in G . Since A_j is semiprojective, there exist N and a $*$ -homomorphism

$$\psi : A_j \longrightarrow \prod_N^\infty B_n$$

such that $\rho_N \circ \psi = \varphi \circ \gamma_j$. Let $\bar{\varphi} = \psi \circ \theta_j$, so that, for any g in G ,

$$\begin{aligned} \|\rho_K \circ \bar{\varphi}(g) - \varphi(g)\| &= \|\varphi \circ \gamma_j \circ \theta_j(g) - \varphi(g)\| \\ (0.1) \qquad \qquad \qquad &\leq \|\gamma_j \circ \theta_j(g) - g\| \\ &\leq \epsilon. \end{aligned}$$

□

REFERENCES

1. B. Blackadar, *Shape theory for C^* -algebras*, Math. Scand. **56** (1985), 249–275. MR **87b**:46074
2. S. Eilers, T. A. Loring and G. K. Pedersen *Morphisms of extensions of C^* -algebras: pushing forward the Busby invariant*, Adv. Math., to appear.
3. S. Eilers, T. A. Loring and G. K. Pedersen *Stability of anticommutation relations. An application of noncommutative CW complexes*, J. reine angew. Math., to appear.
4. E. G. Effros and J. Kaminker, *Homotopy continuity and shape theory for C^* -algebras*, in Geometric Methods in Operator Algebras, Pitman, 1985. MR **88a**:46082
5. P. Friis and M. Rørdam *Almost commuting self-adjoint matrices—a short proof of Huaxin Lin’s theorem*, J. reine angew. Math., **479** (1996), 121–131. MR **97i**:46097
6. H. Lin *Homomorphisms from $C(X)$ into C^* -algebras*, preprint.
7. H. Lin *Almost commuting unitary elements in purely infinite simple C^* -algebras*, Math. Ann. **303** (1995), 599–616. MR **96k**:46101

8. H. Lin *Almost commuting self-adjoint matrices and applications*, Operator Algebras and Their Applications, Fields Institute Communications, vol. 13, Amer. Math. Soc., Providence, RI, 1997, pp. 193–233. CMP 97:05
9. T. A. Loring, *Lifting Solutions to Perturbing Problems*, Fields Institute Monograph Series, vol. 8, Amer. Math. Soc., Providence, RI, 1997. MR 97:04

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE,
NEW MEXICO 87131

E-mail address: loring@math.unm.edu