CAUCHY-SCHWARZ AND MEANS INEQUALITIES FOR ELEMENTARY OPERATORS INTO NORM IDEALS

DANKO R. JOCIĆ

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. The Cauchy-Schwarz norm inequality for normal elementary operators

$$\left\| \sum_{n=1}^{\infty} A_n X B_n \right\| \leq \left\| \left(\sum_{n=1}^{\infty} A_n^* A_n \right)^{1/2} X \left(\sum_{n=1}^{\infty} B_n^* B_n \right)^{1/2} \right\|,$$

implies a means inequality for generalized normal derivations

$$\left\| \frac{AX + XB}{2} \right\| \leq \|X\|^{1-\frac{1}{r}} \left\| \frac{|A|^r X + X|B|^r}{2} \right\|^{\frac{1}{r}},$$

for all $r \geq 2$, as well as an inequality for normal contractions A and B

$$\left\| (I - A^* A)^{\frac{1}{2}} X (I - B^* B)^{\frac{1}{2}} \right\| \leq \|X - AXB\|,$$

for all X in $B(H)$ and for all unitarily invariant norms $\|\cdot\|$.

1. INTRODUCTION

Let $B(H)$ and $\mathcal{C}_\infty$ stand respectively for spaces of all bounded and all compact linear operators acting on a separable, infinite-dimensional, complex Hilbert space H . For an $X \in B(H)$ let $\|X\|$ denote its norm, and for an arbitrary $X \in \mathcal{C}_\infty$ let $s_1(X) \geq s_2(X) \geq \cdots$ denote the singular values of X , i.e., the eigenvalues of $|X| = (X^* X)^{1/2}$, arranged in a non-increasing order, with their multiplicities counted. Each “symmetric gauge function” Φ on sequences gives rise to a unitarily invariant (u.i.) norm on operators defined by $\|A\|_\Phi = \Phi(\{s_n(A)\})$. We will denote by the symbol $\|\cdot\|$ any such norm. Any such norm is defined on a natural subclass $\mathcal{C}_{\|\cdot\|}$ of $\mathcal{C}_\infty$ called the norm ideal associated with the norm $\|\cdot\|$, and satisfies the invariance property $\|UAV\| = \|A\|$ for all A in this ideal and for all unitary operators U, V . Each norm ideal $\mathcal{C}_{\|\cdot\|}$ is closed in the topology generated by the norm $\|\cdot\|$. Particularly well known among unitarily invariant norms are the Schatten p -norms, defined as $\|X\|_p = (\sum_{n=1}^{\infty} s_n^p(A))^{1/p}$ for $1 \leq p < \infty$ and $\|X\|_\infty = \|X\| = s_1(X)$, which represent the norms on the Schatten p -ideals $\mathcal{C}_p$. The Ky Fan norms, defined as $\|A\|_k = \Phi_k(s_i(A)) = \sum_{i=1}^k s_i(A)$ for $k = 1, 2, \dots$, represent another interesting family of unitarily invariant norms. The associated ideals $\mathcal{C}_\infty^{(k)}$ consist of all compact operators as every Ky Fan k -norm is equivalent to the norm in $\mathcal{C}_\infty$. The property saying that for all $X \in \mathcal{C}_\infty$ and $Y \in \mathcal{C}_{\|\cdot\|}$ with $\|X\|_k \leq \|Y\|_k$ for all $k \geq 1$ we have

Received by the editors March 12, 1996 and, in revised form, February 4, 1997.

1991 *Mathematics Subject Classification*. Primary 47A30; Secondary 47B05, 47B10, 47B15.

Key words and phrases. Unitarily invariant norms, Ky Fan dominance property.

$X \in \mathcal{C}_{\|\cdot\|}$ with $\|X\| \leq \|Y\|$ is known as the Ky Fan dominance property ([GK], ch. 3, §4). For a complete account of the theory of norm ideals, the reader is referred to [GK], [Sch] and [Si].

If $\mathcal{A} = (A_1, \dots, A_N)$ and $\mathcal{B} = (B_1, \dots, B_N)$ are N -tuples of bounded Hilbert space operators, then the elementary operator $R = R_{\mathcal{A}, \mathcal{B}}$ on $B(H)$ is defined by $R(X) = \sum_{n=1}^N A_n X B_n$. Elementary operators were introduced by Lumer and Rosenblum in [LR], who studied their spectral properties. In this setting many authors subsequently studied spectral, algebraic, metric and structural properties of elementary operators (see [F83], [F85], [F87], [FL], [McI], and the references therein).

If both $\{A_n\}_{n=1}^N$ and $\{B_n\}_{n=1}^N$ are families of commuting normal operators, one can easily show that the associated elementary operator $R_{\mathcal{A}, \mathcal{B}}$ is normal when restricted to the Hilbert space $\mathcal{C}_2$. So, in the sequel, by a normal elementary operator we will always mean such an operator, including those cases of $N = \infty$ with the convergence guaranteed. As was shown by G. Weiss in his paper [W83], a famous Fuglede-Putnam type theorem extends to such operators. Very important examples of normal elementary operators are so called “pinching” operators $R_{\mathcal{P}}(X) = \sum_{n=1}^{\infty} P_n X P_n$, generated by a family of mutually orthogonal self-adjoint projections $\{P_n\}_{n=1}^{\infty}$.

2. MAIN RESULTS

We start with the basic Cauchy-Schwarz norm inequality for normal elementary operators. The following theorem extends “pinching” theorems 2.5.1 of [GK] and 1.19 of [Si].

Theorem 2.1. *If $\sum_{n=1}^{\infty} C_n^* C_n \leq 1$, $\sum_{n=1}^{\infty} C_n C_n^* \leq 1$, $\sum_{n=1}^{\infty} D_n^* D_n \leq 1$ and $\sum_{n=1}^{\infty} D_n D_n^* \leq 1$ for some operator families $\{C_n\}_{n=1}^{\infty}$ and $\{D_n\}_{n=1}^{\infty}$, then also $\sum_{n=1}^{\infty} C_n Y D_n \in \mathcal{C}_{\|\cdot\|}$ whenever $Y \in \mathcal{C}_{\|\cdot\|}$ for some unitarily invariant norm $\|\cdot\|$, and moreover*

$$(2.1) \quad \left\| \sum_{n=1}^{\infty} C_n Y D_n \right\| \leq \|Y\|.$$

Proof. For arbitrary f and g in H a straightforward calculation gives

$$\begin{aligned} | \langle (\sum_{n=1}^{\infty} C_n Y D_n) f, g \rangle | &\leq \sum_{n=1}^{\infty} \|Y\| \|D_n f\| \|C_n^* g\| \\ &\leq \|Y\| \left(\sum_{n=1}^{\infty} \|D_n f\|^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \|C_n^* g\|^2 \right)^{1/2} \\ &= \|Y\| \left\langle \sum_{n=1}^{\infty} D_n^* D_n f, f \right\rangle^{1/2} \left\langle \sum_{n=1}^{\infty} C_n C_n^* g, g \right\rangle^{1/2} \\ &= \|Y\| \left\| \left(\sum_{n=1}^{\infty} C_n C_n^* \right)^{1/2} g \right\| \left\| \left(\sum_{n=1}^{\infty} D_n^* D_n \right)^{1/2} f \right\| \leq \|Y\| \|f\| \|g\|, \end{aligned}$$

from which we conclude that

$$(2.2) \quad \left\| \sum_{n=1}^{\infty} C_n Y D_n \right\| \leq \|Y\|.$$

Therefore, for all $N = 1, 2, \dots$, for $Y \in \mathcal{C}_1$ and for all $W \in B(H)$ we have

$$\begin{aligned} |tr(\sum_{n=1}^N C_n Y D_n W^*)| &= |tr(Y(\sum_{n=1}^N C_n^* W D_n^*)^*)| \\ &\leq \|Y\|_1 \|\sum_{n=1}^N C_n^* W D_n^*\| \leq \|Y\|_1 \|W\|, \end{aligned}$$

according to (2.2), from which we deduce that

$$(2.3) \quad \|\sum_{n=1}^N C_n Y D_n\|_1 \leq \|Y\|_1.$$

If $Y \in \mathcal{C}_\infty$, let $Y = \sum_{n=1}^\infty s_n(Y) \langle \cdot, e_n \rangle f_n$ be a singular value decomposition for some orthonormal systems $\{e_n\}$ and $\{f_n\}$. For all $k \geq 2$ we introduce operators $Z = \sum_{n=1}^{k-1} (s_n(Y) - s_{n+1}(Y)) \sum_{j=1}^n \langle \cdot, e_j \rangle f_j$ and $V = s_k(Y) \sum_{n=1}^k \langle \cdot, e_n \rangle f_n + \sum_{n=k+1}^\infty s_n(Y) \langle \cdot, e_n \rangle f_n$. We see that

$$\begin{aligned} Z &= \sum_{n=1}^{k-1} \sum_{j=1}^n (s_n(Y) - s_{n+1}(Y)) \langle \cdot, e_j \rangle f_j \\ &= \sum_{j=1}^k (s_j(Y) - s_k(Y)) \langle \cdot, e_j \rangle f_j \\ &= \sum_{n=1}^k s_n(Y) \langle \cdot, e_n \rangle f_n + s_k(Y) \sum_{n=1}^k \langle \cdot, e_n \rangle f_n = Y - V. \end{aligned}$$

We can also note that $s_1(V) = \dots = s_k(V) = s_k(Y)$, due to orthogonality of the systems $\{e_n\}$ and $\{f_n\}$. That will allow us to conclude that for all Ky Fan k -norms we have

$$\begin{aligned} \left\| \sum_{n=1}^N C_n Y D_n \right\|_k &\leq \left\| \sum_{n=1}^N C_n Z D_n \right\|_k + \left\| \sum_{n=1}^N C_n V D_n \right\|_k \\ (2.4) \quad &\leq \|Z\|_1 + k \left\| \sum_{n=1}^N C_n Z D_n \right\|_\infty \end{aligned}$$

$$(2.5) \quad \leq \sum_{n=1}^{k-1} (s_n(Y) - s_{n+1}(Y)) \sum_{j=1}^n \|\langle \cdot, e_j \rangle f_j\|_\infty + k \|V\|_\infty$$

$$(2.6) \quad \leq \sum_{n=1}^{k-1} n(s_n(Y) - s_{n+1}(Y)) + k s_k(Y) = \sum_{n=1}^k s_n(Y) = \|Y\|_k,$$

with (2.4) following from (2.3) and (2.5) from (2.2).

Moreover, if Y is in $\mathcal{C}_\infty$ then also $\sum_{n=1}^\infty C_n Y D_n \in \mathcal{C}_\infty$. Indeed, elementary operators $R_N(Y) = \sum_{n=1}^N C_n Y D_n$ acting on $\mathcal{C}_\infty^{(k)}$ represent a bounded family, because $\|R_N(Y)\|_k \leq \|Y\|_k$ for all $Y \in \mathcal{C}_\infty$ by (2.6). Also, for one dimensional operators

$f \otimes g$ and $M > N$ we have

$$\begin{aligned} \|R_M(f \otimes g) - R_N(f \otimes g)\|_k &\leq \left\| \sum_{n=N+1}^M D_n^* f \otimes C_n g \right\|_1 \\ &\leq \sum_{n=N+1}^M \|D_n^* f\| \|C_n g\| \leq \left\| \left(\sum_{n=N+1}^M C_n C_n^* \right)^{\frac{1}{2}} g \right\| \left\| \left(\sum_{n=N+1}^M D_n^* D_n \right)^{\frac{1}{2}} f \right\|, \end{aligned}$$

which $\rightarrow 0$ as $M, N \rightarrow \infty$. Therefore $R_N(Y)$ converge in $\mathcal{C}_\infty^{(k)}$ for all finite dimensional Y to a **compact** operator. By the uniform boundedness principle the same is true for all $Y \in \mathcal{C}_\infty^{(k)}$, due to its separability. So (2.1) holds for all Ky Fan k -norms, and we therefore invoke the Ky Fan dominance property to conclude that (2.1) holds for all unitarily invariant norms, as required. $\square$

In the sequel we will refer to a family $\{A_n\}_{n=1}^\infty$ in $B(H)$ as square summable if $\sum_{n=1}^\infty \|A_n f\|^2 < \infty$ for all $f \in H$. Though this means just the weak convergence of $\sum_{n=1}^\infty A_n^* A_n$, an appeal to the resonance principle shows that $\sum_{n=1}^\infty A_n^* A_n$ actually defines a **bounded** Hilbert space operator, and due to the monotonicity of its partial sums, the convergence is moreover strong. For such families the following Cauchy-Schwarz inequality holds:

Theorem 2.2. *For a square summable families $\{A_n\}_{n=1}^\infty$ and $\{B_n\}_{n=1}^\infty$ of commuting normal operators*

$$(2.7) \quad \left\| \sum_{n=1}^\infty A_n X B_n \right\| \leq \left\| \left(\sum_{n=1}^\infty A_n^* A_n \right)^{1/2} X \left(\sum_{n=1}^\infty B_n^* B_n \right)^{1/2} \right\|,$$

for all $X \in B(H)$ and for all u.i. norms $\|\cdot\|$. If $\mathcal{C}_{\|\cdot\|}$ is separable and $X \in \mathcal{C}_{\|\cdot\|}$, then the left-hand side sum converges in the norm of this ideal.

Proof. First, we need a suitable factorization for Hilbert space operators A_n and B_n . Let $A = (\sum_{n=1}^\infty A_n^* A_n)^{\frac{1}{2}}$ and $B = (\sum_{n=1}^\infty B_n^* B_n)^{\frac{1}{2}}$, and let P and Q denote respectively the orthogonal projections on $R(A)$ and $R(B)$. If for a given $f \in H$ we have that $Pf = \lim_{k \rightarrow \infty} Ag_k$ for some sequence $\{g_k\}$ in H , then $\lim_{k \rightarrow \infty} A_n g_k$ exists for all $n \geq 1$ and does not depend on the chosen sequence. Indeed,

$$\|A_n g_k - A_n g_l\| \leq \|A(g_k - g_l)\| \rightarrow \|Pf - Pf\| = 0$$

as $k, l \rightarrow \infty$, and also $\|A_n g_k - A_n h_k\| \leq \|A(g_k - h_k)\| \rightarrow 0$ as $k \rightarrow \infty$ whenever $\lim_{k \rightarrow \infty} Ah_k = Pf$ for some other sequence $\{h_k\}$. Thus we can correctly introduce operators $C_n, n = 1, 2, \dots$, by $C_n f = \lim_{k \rightarrow \infty} A_n g_k$, where $\{g_k\}$ is any sequence in H such that $\lim_{k \rightarrow \infty} Ag_k = Pf$. Let us note that due to our definition every C_n vanishes on $N(A)$, i.e., $C_n = C_n P$, and also $C_n A = AC_n = A_n$. Moreover, $\sum_{n=1}^\infty C_n^* C_n = P$. Indeed, $\sum_{n=1}^\infty C_n^* C_n A^2 = \sum_{n=1}^\infty A_n^* A_n = A^2$ implies $\sum_{n=1}^\infty C_n^* C_n P = P$, which together with the fact that $C_n(I - P) = 0$ gives the desired conclusion. For all $m, n = 1, 2, \dots$, C_m^* and C_n commute on $R(A^2)$ and $N(A^2)$, and so also on all of H . Thus $\{C_n\}_{n=1}^\infty$ is a commuting family of normal contractions which realize the factorizations $C_n A = AC_n = A_n$, with $\sum_{n=1}^\infty C_n^* C_n = P$, and which commute with the family $\{A_n\}_{n=1}^\infty$. Similarly we get a commuting family $\{D_n\}_{n=1}^\infty$ of normal contractions which also commute with $\{B_n\}_{n=1}^\infty$ and satisfy $D_n B = BD_n = B_n$ and $\sum_{n=1}^\infty D_n^* D_n = Q$. One could easily derive the next explicit

formula: $C_n = \overline{A_n A^\dagger} = \overline{A^\dagger A_n}$, where $A^\dagger$ denotes a (densely defined) Moore-Penrose (generalized) inverse for A .

For $Y = AXB \in \mathcal{C}_{\|\cdot\|}$ (there is nothing to prove in the opposite case), an application of Theorem 2.1 gives

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} A_n X B_n \right\| &= \left\| \sum_{n=1}^{\infty} C_n Y D_n \right\| \\ (2.8) \quad &\leq \|Y\| = \left\| \left(\sum_{n=1}^{\infty} A_n^* A_n \right)^{1/2} X \left(\sum_{n=1}^{\infty} B_n^* B_n \right)^{1/2} \right\|, \end{aligned}$$

which proves the first part of theorem.

Finally, if $\mathcal{C}_{\|\cdot\|}$ is separable, then for all $N = 1, 2, \dots$, an application of the just proven part of theorem combined with the arithmetic-geometric means inequality in [BhD] gives

$$\begin{aligned} \left\| \sum_{n=N}^{\infty} A_n X B_n \right\| &\leq \left\| \left(\sum_{n=N}^{\infty} A_n^* A_n \right)^{\frac{1}{2}} X \left(\sum_{n=N}^{\infty} B_n^* B_n \right)^{\frac{1}{2}} \right\| \\ &= \left\| \left(\sum_{n=N}^{\infty} C_n^* C_n \right)^{\frac{1}{2}} A X B \left(\sum_{n=N}^{\infty} D_n^* D_n \right)^{\frac{1}{2}} \right\| \\ (2.9) \quad &\leq \frac{1}{2} \left\| \left(\sum_{n=N}^{\infty} C_n^* C_n \right) A X B + A X B \left(\sum_{n=N}^{\infty} D_n^* D_n \right) \right\|. \end{aligned}$$

We see by (2.8) that $\{\sum_{n=N}^{\infty} C_n^* C_n\}_{N=1}^{\infty}$ and $\{\sum_{n=N}^{\infty} D_n^* D_n\}_{N=1}^{\infty}$ represent bounded sequences of selfadjoint operators which strongly converge to 0 as $N \rightarrow \infty$. As $AXB \in \mathcal{C}_{\|\cdot\|}$, which is separable, then the right-hand side of (2.9) tends to 0 as $N \rightarrow \infty$ by Theorem 3.6.3. of [GK]. The conclusion follows. $\square$

Corollary 2.1. *For normal A and B in $B(H)$ and for all real $r \geq 2$,*

$$(2.10) \quad \left\| \frac{AX + XB}{2} \right\| \leq \left\| \left(\frac{1 + |A|^r}{2} \right)^{\frac{1}{r}} X \left(\frac{1 + |B|^r}{2} \right)^{\frac{1}{r}} \right\|$$

as well as

$$(2.11) \quad \left\| \frac{X + AXB}{2} \right\| \leq \left\| \left(\frac{1 + |A|^r}{2} \right)^{\frac{1}{r}} X \left(\frac{1 + |B|^r}{2} \right)^{\frac{1}{r}} \right\|$$

for all $X \in B(H)$ and for all u.i. norms $\|\cdot\|$.

Proof. $\{A, I\}$ and $\{I, B\}$ are families of normal commuting operators, and so for $r = 2$ the desired conclusion follows by Theorem 2.2. For $r > 2$ the mapping $t \rightarrow t^{\frac{2}{r}}$ is operator monotone by a well known Heinz theorem, and therefore this is an operator concave mapping (see [BSh]). Specifically, $\frac{1+|A|^2}{2} \leq \left(\frac{1+|A|^r}{2} \right)^{\frac{2}{r}}$, from which we obtain $\left\| \left(\frac{1+|A|^2}{2} \right)^{\frac{1}{2}} \left(\frac{1+|A|^r}{2} \right)^{-\frac{1}{r}} \right\| \leq 1$ and similarly $\left\| \left(\frac{1+|B|^2}{2} \right)^{\frac{1}{2}} \left(\frac{1+|B|^r}{2} \right)^{-\frac{1}{r}} \right\| \leq 1$. Therefore

$$\left\| \left(\frac{1+|A|^2}{2} \right)^{\frac{1}{2}} X \left(\frac{1+|B|^2}{2} \right)^{\frac{1}{2}} \right\| \leq \left\| \left(\frac{1+|A|^r}{2} \right)^{\frac{1}{r}} X \left(\frac{1+|B|^r}{2} \right)^{\frac{1}{r}} \right\|,$$

which completes the proof. $\square$

Corollary 2.2. *For normal A and B in $B(H)$ the inequality*

$$(2.12) \quad \left\| \frac{AX + XB}{2} \right\| \leq \|X\|^{1-\frac{1}{r}} \left\| \frac{|A|^r X + X|B|^r}{2} \right\|^{\frac{1}{r}}$$

holds for all real $r \geq 2$, for all u.i. norms $\|\cdot\|$ and for all $X \in \mathcal{C}_{\|\cdot\|}$.

Proof. By Corollary 2.1, for all $t > 0$,

$$\begin{aligned} \left\| \frac{AX + XB}{2} \right\| &= t^{-1} \left\| \frac{tAX + XtB}{2} \right\| \\ &\leq t^{-1} \left\| \left(\frac{1 + |tA|^r}{2} \right)^{\frac{1}{r}} X \left(\frac{1 + |tB|^r}{2} \right)^{\frac{1}{r}} \right\|, \end{aligned}$$

and therefore

$$\left\| \frac{AX + XB}{2} \right\| \leq t^{-1} \|X\|^{1-\frac{2}{r}} \left\| \left(\frac{1 + |tA|^r}{2} \right)^{\frac{1}{2}} X \left(\frac{1 + |tB|^r}{2} \right)^{\frac{1}{2}} \right\|^{\frac{2}{r}},$$

by [Ki], because $\frac{2}{r} < 1$. Therefore, the arithmetic-geometric mean inequality implies

$$\begin{aligned} \left\| \frac{AX + XB}{2} \right\| &\leq \frac{1}{2t} \|X\|^{1-\frac{2}{r}} \left\| \frac{1 + |tA|^r}{2} X + X \frac{1 + |tB|^r}{2} \right\|^{\frac{2}{r}} \\ (2.13) \quad &\leq \frac{1}{2} \|X\|^{1-\frac{2}{r}} \left(t^{-\frac{r}{2}} \|X\| + t^{\frac{r}{2}} \left\| \frac{|A|^r X + X|B|^r}{2} \right\|^{\frac{2}{r}} \right). \end{aligned}$$

As the right-hand side equals $\|X\|^{1-\frac{1}{r}} \left\| \frac{|A|^r X + X|B|^r}{2} \right\|^{\frac{1}{r}}$, which attains its minimum for $t = \|X\|^{\frac{1}{r}} \left\| \frac{|A|^r X + X|B|^r}{2} \right\|^{-\frac{1}{r}}$, the conclusion follows. $\square$

Theorem 2.3. *For normal contractions A and B the inequality*

$$(2.14) \quad \left\| (I - A^*A)^{\frac{1}{2}} X (I - B^*B)^{\frac{1}{2}} \right\| \leq \|X - AXB\|,$$

holds for all $X \in B(H)$ and for all unitarily invariant norms $\|\cdot\|$.

Proof. First, we note that $s\text{-}\lim_{n \rightarrow \infty} A^n(I - A^*A)^{\frac{1}{2}} = 0$. Indeed, by a spectral theorem, for every $f \in H$ there is a positive, finite Borel measure μ concentrated on $D = \{z \in \mathbb{C} : |z| \leq 1\}$ such that $\|A^n(I - A^*A)^{\frac{1}{2}} f\|^2 = \int_D |z|^{2n}(1 - |z|^2) d\mu_f(z)$, whence the desired conclusion follows by Lebesgue's dominating convergence theorem. Therefore

$$w\text{-}\lim_{n \rightarrow \infty} (I - A^*A)^{\frac{1}{2}} (X - A^n X B^n) (I - B^*B)^{\frac{1}{2}} = (I - A^*A)^{\frac{1}{2}} X (I - B^*B)^{\frac{1}{2}}.$$

So by Theorem 2.2 we get

$$\begin{aligned}
 & \left\| (I - A^*A)^{\frac{1}{2}} X (I - B^*B)^{\frac{1}{2}} \right\| \\
 &= \left\| \lim_{n \rightarrow \infty} (I - A^*A)^{\frac{1}{2}} (X - A^n X B^n) (I - B^*B)^{\frac{1}{2}} \right\| \\
 &= \left\| \sum_{k=0}^{\infty} (I - A^*A)^{\frac{1}{2}} A^k (X - AXB) B^k (I - B^*B)^{\frac{1}{2}} \right\| \\
 &\leq \left\| \left(\sum_{k=0}^{\infty} (I - |A|^2) |A|^{2k} \right)^{\frac{1}{2}} (X - AXB) \left(\sum_{k=0}^{\infty} |B|^{2k} (I - |B|^2) \right)^{\frac{1}{2}} \right\| \\
 (2.15) \quad &= \left\| (I - P)(X - AXB)(I - Q) \right\| \leq \|X - AXB\|,
 \end{aligned}$$

where P and Q are the orthogonal projections on $\text{Ker}(I - A^*A)$ and $\text{Ker}(I - B^*B)$ respectively. This concludes the proof. $\square$

REFERENCES

- [BSh] J. Bendat and S. Sherman, *Monotone and Convex Operator Functions*, Trans. Amer. Math. Soc. **79** (1955), 58–71. MR **18**:588b
- [BhD] R. Bhatia and Ch. Davis, *More matrix forms of the arithmetic-geometric mean inequality*, SIAM J. Matrix. Anal. Appl. **14** (1993), 132–136. MR **94b**:15017
- [F83] L. Fialkow, *Spectral properties of elementary operators*, Acta Sci. Math. (Szeged), **46** (1983), 269–282. MR **85h**:47003
- [F85] L. Fialkow, *Spectral properties of elementary operators II*, Trans. Amer. Math. Soc. **290** (1985), 415–429. MR **86j**:47005
- [F87] L. Fialkow, *The range inclusion problem for elementary operators*, Michigan J. Math. **34** (1987), 451–459. MR **89a**:47052
- [FL] L. Fialkow and R. Loeb, *Elementary mappings into ideals of operators*, Illinois Journal Math. Vol **28** (1984), 555–578. MR **86g**:47054
- [GK] I.C. Gohberg and M.G. Krein, *Introduction to the theory of linear nonselfadjoint operators*, Transl. Math. Monographs, vol. **18**, Amer. Math. Soc. Providence, R.I. 1969. MR **39**:7447
- [Ki] F. Kittaneh, *Norm inequalities for fractional powers of positive operators*, Lett. Math. Phys. **27** (1993), 279–285. MR **94e**:47011
- [LR] G. Lumer and M. Rosenblum, *Linear operator equations*, Proc. Amer. Math. Soc. **10** (1959), 32–41. MR **21**:2927
- [McI] A. McIntosh, A. Pryde and W. Ricker, *Estimates for the solution of operator equation $\sum_{j=1}^m A_j Q B_j = U$* , Operator Theory, Adv. & Appl., Vol. **20**, pp. 197–207. Birkhauser Verlag Basel 1988. MR **89k**:47024
- [Sch] R. Schatten, *Norm Ideals of completely continuous operators*, Springer-Verlag, Berlin, 1960. MR **22**:9878
- [Si] B. Simon, *Trace Ideals and their applications*, Cambridge University Press, 1979. MR **80k**:47048
- [W83] G. Weiss, *An extension of the Fuglede commutativity theorem modulo the Hilbert-Schmidt class to the operators of the form $\sum M_n X N_n$* , Trans. Amer. Math. Soc. **278** (1983), 1–20. MR **84e**:47026

UNIVERSITY OF BELGRADE, FACULTY OF MATHEMATICS, STUDENSKI TRG 16, P. O. Box 550,
11000 BELGRADE, YUGOSLAVIA

E-mail address: jovic@matf.bg.ac.yu