MULTIPLIER THEOREMS FOR HERZ TYPE HARDY SPACES

SHANZHEN LU AND DACHUN YANG

(Communicated by J. Marshall Ash)

ABSTRACT. In this paper, the authors establish a multiplier theorem for Herz type Hardy spaces.

Let T_m be a multiplier operator defined in terms of Fourier transforms by

$$\widehat{T_m f}(\xi) = m(\xi)\widehat{f}(\xi)$$

for suitable functions f . It is well-known that there is a multiplier theorem for $H^1(\mathbb{R}^n)$ (see [FS]): if $\alpha > n/2$ and

$$(1) \quad \int_{R < |\xi| < 2R} |D^\beta m(\xi)|^2 d\xi \leq CR^{n-2|\beta|}, \quad 0 < R < \infty,$$

for all $|\beta| \leq \alpha$, then T_m can be extended to be a bounded operator on $H^1(\mathbb{R}^n)$. That is, m is a bounded multiplier of $H^1(\mathbb{R}^n)$.

Fix a function $\eta \in C_0^\infty(\mathbb{R}^n)$ with $0 \leq \eta \leq 1$, $\eta = 1$ on $1/2 \leq |\xi| \leq 2$ and $\text{supp } \eta \subset \{1/4 \leq |\xi| \leq 4\}$. For $\delta > 0$, let us denote

$$m_\delta(\xi) = m(\delta\xi)\eta(\xi).$$

It is easy to check that (1) is equivalent to

$$(2) \quad \sup_{\delta} \|\widehat{m_\delta}\|_{K_2^{\alpha,2}(\mathbb{R}^n)} < \infty,$$

where $K_2^{\alpha,2}(\mathbb{R}^n)$ is a non-homogeneous Herz space (see [BS]). By using some embedding relations on Herz spaces, A. Baernstein II and E. T. Sawyer [BS] weakened (2) into

$$(3) \quad \sup_{\delta} \|\widehat{m_\delta}\|_{K_1^{\varepsilon,1}(\mathbb{R}^n)} < \infty,$$

where $0 < \varepsilon < \alpha - \frac{n}{2}$. In fact, this is just a special case of their theorem. In [BS], Baernstein and Sawyer showed that m is a bounded multiplier of $H^1(\mathbb{R}^n)$ under an even weaker condition than (3); see Theorem 3b in [BS, page 21].

By using the technique of Herz type Hardy spaces developed by the authors in [LY1]-[LY3] and [Y], in this paper, we shall first establish a multiplier theorem for the homogeneous Herz type Hardy space $\dot{H}K_q^{n(1-1/q),1}(\mathbb{R}^n)$ which is introduced by the authors of this paper in [LY1]. Then as simple consequences of this theorem, a multiplier theorem for the corresponding non-homogeneous version of the space

Received by the editors April 13, 1995 and, in revised form, April 5, 1997.

1991 *Mathematics Subject Classification*. Primary 42B15; Secondary 42B30.

Key words and phrases. Herz space, Hardy space, multiplier.

Research was supported by NNSF and SEDF of China.

$\dot{H}\dot{K}_q^{n(1-1/q),1}(\mathbb{R}^n)$ and the special case mentioned above of the multiplier theorem of Baernstein and Sawyer for $H^1(\mathbb{R}^n)$ will be deduced.

Now, for the reader's convenience, let us recall the definition of the Herz spaces here. For $k \in \mathbb{Z}$, let $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ and $A_k = B_k \setminus B_{k-1}$. We also denote by χ_k the characteristic function of the set A_k .

Definition 1. Let $\alpha \in \mathbb{R}$ and $0 < p, q \leq \infty$.

- (i) The homogeneous Herz space $\dot{K}_q^{\alpha,p}(\mathbb{R}^n)$ is defined in terms of

$$\|f\|_{\dot{K}_q^{\alpha,p}(\mathbb{R}^n)} \equiv \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|f\chi_k\|_{L^q(\mathbb{R}^n)}^p \right\}^{1/p}$$

by letting

$$\dot{K}_q^{\alpha,p}(\mathbb{R}^n) \equiv \{f \in L_{loc}^q(\mathbb{R}^n \setminus \{0\}) : \|f\|_{\dot{K}_q^{\alpha,p}(\mathbb{R}^n)} < \infty\}.$$

- (ii) The non-homogeneous Herz space $K_q^{\alpha,p}(\mathbb{R}^n)$ is defined in terms of

$$\|f\|_{K_q^{\alpha,p}(\mathbb{R}^n)} \equiv \left\{ \|f\chi_{B_0}\|_{L^q(\mathbb{R}^n)}^p + \sum_{k=1}^{\infty} 2^{k\alpha p} \|f\chi_k\|_{L^q(\mathbb{R}^n)}^p \right\}^{1/p}$$

by letting

$$K_q^{\alpha,p}(\mathbb{R}^n) \equiv \{f \in L_{loc}^q(\mathbb{R}^n) : \|f\|_{K_q^{\alpha,p}(\mathbb{R}^n)} < \infty\}.$$

Here the usual modification was made when $p = \infty$.

In what follows, when $p = 1$, $1 < q < \infty$ and $\alpha = n(1 - 1/q)$, we shall abbreviate $\dot{K}_q^{\alpha,p}(\mathbb{R}^n)$ and $K_q^{\alpha,p}(\mathbb{R}^n)$, respectively, as $\dot{K}_q(\mathbb{R}^n)$ and $A^q(\mathbb{R}^n)$. The latter is also said to be the Beurling algebras; see [CL] and [GR].

Definition 2. Let $1 < q < \infty$. For $f \in \mathcal{S}'(\mathbb{R}^n)$, let Gf be the grand maximal function of f (see [FS] for its definition).

- (i) The Hardy space $H\dot{K}_q(\mathbb{R}^n)$ associated with the Herz space $\dot{K}_q(\mathbb{R}^n)$ is defined by

$$H\dot{K}_q(\mathbb{R}^n) \equiv \{f \in \mathcal{S}'(\mathbb{R}^n) : Gf \in \dot{K}_q(\mathbb{R}^n)\}.$$

In this case, we also define $\|f\|_{H\dot{K}_q(\mathbb{R}^n)} \equiv \|Gf\|_{\dot{K}_q(\mathbb{R}^n)}$.

- (ii) The Hardy space $HA^q(\mathbb{R}^n)$ associated with the Beurling algebra $A^q(\mathbb{R}^n)$ is defined by

$$HA^q(\mathbb{R}^n) \equiv \{f \in \mathcal{S}'(\mathbb{R}^n) : Gf \in A^q(\mathbb{R}^n)\}.$$

In this case, we also define $\|f\|_{HA^q(\mathbb{R}^n)} \equiv \|Gf\|_{A^q(\mathbb{R}^n)}$.

We remark that $HA^q(\mathbb{R}^n)$ was first introduced by Chen and Lau in [CL] for $n = 1$, and then by García-Cuerva in [GR] for $n > 1$. Obviously, $H\dot{K}_q(\mathbb{R}^n)$ is a homogeneous version of $HA^q(\mathbb{R}^n)$. Moreover, in [LY1], the authors proved that

$$(4) \quad HA^q(\mathbb{R}^n) = H\dot{K}_q(\mathbb{R}^n) \cap L^q(\mathbb{R}^n)$$

and

$$(5) \quad \|f\|_{HA^q(\mathbb{R}^n)} \sim \|f\|_{H\dot{K}_q(\mathbb{R}^n)} + \|f\|_{L^q(\mathbb{R}^n)}.$$

It is also well-known that $HA^q(\mathbb{R}^n) \subsetneq \dot{HK}_q(\mathbb{R}^n) \subsetneq H^1(\mathbb{R}^n)$ for any $q \in (1, \infty)$.

Let us now formulate our multiplier theorem for $\dot{HK}_q(\mathbb{R}^n)$.

Theorem 1. *Let $q \in (1, \infty)$ and m satisfy*

$$(6) \quad M \equiv \sup_{\delta} \|\widehat{m}_{\delta}\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)} < \infty.$$

Then m is a bounded multiplier of $\dot{HK}_q(\mathbb{R}^n)$.

By Corollary 2 in [BS, page 22], we know that if m satisfies the condition of Theorem 1, then m is a bounded multiplier of $L^q(\mathbb{R}^n)$ for $1 < q < \infty$. Therefore, from (4), (5) and Theorem 1, we have the following simple corollary.

Corollary 1. *Let $q \in (1, \infty)$ and m satisfy (6). Then m is a bounded multiplier of $HA^q(\mathbb{R}^n)$.*

The proof of Theorem 1 is based on the decomposition characterizations of Herz spaces and Herz type Hardy spaces in terms of central units and central atoms respectively. Let us recall that a function $e(x)$ is said to be a central (α, q) unit of restrict type if e satisfies

- i) $\text{supp } e \subset B(0, r), r \geq 1$;
- ii) $\|e\|_{L^q(\mathbb{R}^n)} \leq |B(0, r)|^{-\alpha/n}$.

Lemma 1 ([LY2]). *Let $0 < \alpha < \infty$, $0 < p < \infty$ and $1 \leq q < \infty$. Then $f \in K_q^{\alpha,p}(\mathbb{R}^n)$ if and only if f can be expressed as*

$$f(x) = \sum_{k=0}^{\infty} \lambda_k e_k(x),$$

where each e_k is a central (α, q) unit of restrict type supported on B_k and $\sum_{k=0}^{\infty} |\lambda_k|^p < \infty$. Moreover,

$$\inf \left\{ \left(\sum_k |\lambda_k|^p \right)^{1/p} \right\} \sim \|f\|_{K_q^{\alpha,p}(\mathbb{R}^n)},$$

where the infimum is taken over all of the above decompositions of f .

Let us now turn to the definition of central atoms. A function $a(x)$ is said to be a central $(1, q)$ atom if a satisfies

- (i) $\text{supp } a \subset B(0, r), r > 0$;
- (ii) $\|a\|_{L^q(\mathbb{R}^n)} \leq |B(0, r)|^{1/q-1}$;
- (iii) $\int_{\mathbb{R}^n} a(x) dx = 0$.

Lemma 2 ([LY3]). *Let $1 < q < \infty$. Then $f \in \dot{HK}_q(\mathbb{R}^n)$ if and only if f can be expressed as*

$$f(x) = \sum_{k=-\infty}^{\infty} \lambda_k a_k(x),$$

where each a_k is a central $(1, q)$ atom supported on B_k and $\sum_{k=-\infty}^{\infty} |\lambda_k| < \infty$. Moreover,

$$\inf \left\{ \sum_{k=-\infty}^{\infty} |\lambda_k| \right\} \sim \|f\|_{\dot{HK}_q(\mathbb{R}^n)},$$

where the infimum is taken over all of the above decompositions of f .

To prove Theorem 1, we still need a lemma. Let $\psi \in \mathcal{S}(\mathbb{R}^n)$, the Schwartz space of functions. In what follows, we let $\widehat{a_\delta}(\xi) \equiv \widehat{a}(\delta\xi)\psi(\xi)$.

Lemma 3. *Let a be a central $(1, q)$ atom supported on $B(0, 1)$ and $b_j = (\widehat{a_{2^j}})^\vee$. Then for any given $d > 0$, we have the following three facts:*

- (i) $\|b_j\|_{L^q(\mathbb{R}^n)} \leq C2^{-nj(1-1/q)}$.
- (ii) $|b_j(x)| \leq C_d 2^{-nj(1-1/q)} |x|^{-d}$, for $|x| \geq 2^{j+1}$.
- (iii) $|b_j(x)| \leq C_d 2^j (1 + |x|)^{-d}$, for all x and $j \leq 0$.

Proof. Since $1 < q < \infty$ and

$$b_j(x) = 2^{-nj} \int_{|x-y| < 2^j} a(2^{-j}(x-y)) \widehat{\psi}(y) dy,$$

it follows from the generalized Minkowski inequality that

$$\begin{aligned} \|b_j\|_{L^q(\mathbb{R}^n)} &\leq 2^{-nj} \int_{\mathbb{R}^n} \left\{ \int_{\mathbb{R}^n} |a(2^{-j}(x-y))|^q dx \right\}^{1/q} |\widehat{\psi}(y)| dy \\ &\leq C 2^{-nj} 2^{nj/q} \|a\|_{L^q(\mathbb{R}^n)} \|\widehat{\psi}\|_{L^1(\mathbb{R}^n)} \leq C 2^{-nj(1-1/q)}. \end{aligned}$$

Thus, (i) holds. Let us now assume $|x| \geq 2^{j+1}$. Note that $|y| \geq |x|/2$ and $\widehat{\psi} \in \mathcal{S}(\mathbb{R}^n)$. Then we have

$$\begin{aligned} |b_j(x)| &\leq 2^{-nj} \int_{|x-y| < 2^j} |a(2^{-j}(x-y))| \cdot |\widehat{\psi}(y)| dy \\ &\leq 2^{-nj} \left(\int_{|x-y| < 2^j} |a(2^{-j}(x-y))|^q dy \right)^{1/q} \left(\int_{|x-y| < 2^j} |\widehat{\psi}(y)|^{q'} dy \right)^{1/q'} \\ &= 2^{nj(1/q-1)} \|a\|_{L^q(\mathbb{R}^n)} \left(\int_{|y| \geq |x|/2} |\widehat{\psi}(y)|^{q'} dy \right)^{1/q'} \\ &\leq C_d 2^{nj(1/q-1)} |x|^{-d}. \end{aligned}$$

Thus, (ii) also holds. Finally, let us assume $j \leq 0$. Since $\int_{\mathbb{R}^n} a(y) dy = 0$, we have

$$b_j(x) = \int_{|y| < 1} a(y) \{\widehat{\psi}(x - 2^j y) - \widehat{\psi}(x)\} dy.$$

It follows from the mean value theorem that there exists a $\theta \in (0, 1)$ such that

$$|b_j(x)| \leq \int_{|y| < 1} |a(y)| \cdot |(\nabla_x \widehat{\psi})(x - \theta 2^j y)| 2^j |y| dy,$$

where $\nabla_x = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right)$. Note that $1 + |x - \theta 2^j y| \geq (1 + |x|)/2$ and $\widehat{\psi} \in \mathcal{S}(\mathbb{R}^n)$. Then we have

$$|b_j(x)| \leq C_d 2^j (1 + |x|)^{-d} \int_{|y| < 1} |y| \cdot |a(y)| dy \leq C_d 2^j (1 + |x|)^{-d}.$$

This completes the proof of the lemma. $\square$

Proof of Theorem 1. By the decompositions of $H\dot{K}_q(\mathbb{R}^n)$ in terms of central atoms, it suffices to prove that the inequality

$$(7) \quad \|T_m a\|_{H\dot{K}_q(\mathbb{R}^n)} \leq C$$

holds for all central $(1, q)$ atoms $a(x)$. Let $a(x)$ be a central $(1, q)$ atom. Since M is invariant for all dilations of m , we may assume $\text{supp } a \subset B(0, 1)$. In Lemma 3,

if we take $\psi \in C_0^\infty(\mathbb{R}^n)$ such that $\text{supp } \psi \subset \{\xi : 1/2 \leq |\xi| \leq 2\}$, $0 \leq \psi \leq 1$, and $\sum_{j \in \mathbb{Z}} \psi(\xi 2^{-j}) = 1$, then $\psi\eta = \psi$ and

$$\begin{aligned} m(\xi)\widehat{a}(\xi) &= \sum_{j=-\infty}^{\infty} m(\xi)\eta(2^{-j}\xi)\widehat{a}(\xi)\psi(2^{-j}\xi) \\ &= \sum_{j=-\infty}^{\infty} m_{2^j}(2^{-j}\xi)\widehat{a_{2^j}}(2^{-j}\xi), \end{aligned}$$

where $\widehat{a_\delta}(\xi) = \widehat{a}(\delta\xi)\psi(\xi)$. By letting $N_j \equiv (m_{2^j})^\vee$, we have

$$(8) \quad T_m a(x) = \sum_{j=-\infty}^{\infty} 2^{nj}(N_j * b_j)(2^j x).$$

Without loss of generality, we may assume $M = 1$. Thus, the inequality $\|\widehat{m_\delta}\|_{L^1(\mathbb{R}^n)} \leq 1$ holds for any $\delta > 0$. Therefore, it follows from the Hausdorff-Young inequality that $\|m\|_{L^\infty(\mathbb{R}^n)} \leq 1$ and

$$\|N_j\|_{L^1(\mathbb{R}^n)} \leq \|N_j\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)} \leq 1.$$

Let us first prove

$$(9) \quad \|T_m a\|_{\dot{K}_q(\mathbb{R}^n)} \leq C,$$

where C is independent of a and a is a central $(1, q)$ atom with $\text{supp } a \subset B(0, 1)$. We write

$$\begin{aligned} \|T_m a\|_{\dot{K}_q(\mathbb{R}^n)} &= \sum_{k=-\infty}^{\infty} 2^{kn(1-1/q)} \|(T_m a)\chi_k\|_{L^q(\mathbb{R}^n)} \\ &= \sum_{k=-\infty}^2 \cdots + \sum_{k=3}^{\infty} \cdots \equiv I_1 + I_2. \end{aligned}$$

Since m is a bounded multiplier of $L^q(\mathbb{R}^n)$ by Corollary 2 in [BS, page 22], we have

$$I_1 \leq C \sum_{k=-\infty}^2 2^{kn(1-1/q)} \|a\|_{L^q(\mathbb{R}^n)} \leq C \sum_{k=-\infty}^2 2^{nk(1-1/q)} \leq C.$$

On the other hand, by (8), we have

$$\begin{aligned} I_2 &= \sum_{k=3}^{\infty} 2^{kn(1-1/q)} \|(T_m a)\chi_k\|_{L^q(\mathbb{R}^n)} \\ &\leq \sum_{k=3}^{\infty} \sum_{j=-\infty}^{\infty} 2^{nj} 2^{kn(1-1/q)} \|(N_j * b_j)(2^j \cdot)\chi_k(\cdot)\|_{L^q(\mathbb{R}^n)} \\ &= \sum_{j=-\infty}^{\infty} \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \|(N_j * b_j)(\cdot)\chi_l(\cdot)\|_{L^q(\mathbb{R}^n)} \\ &= \sum_{j=-\infty}^0 \sum_{l=j+3}^{\infty} \cdots + \sum_{j=1}^{\infty} \sum_{l=j+3}^{\infty} \cdots \equiv I_{2,1} + I_{2,2}. \end{aligned}$$

Let us first estimate $I_{2,1}$. By Lemma 1, N_j can be expressed as

$$N_j(x) = \sum_{k=0}^{\infty} \lambda_k^j e_k^j(x),$$

where each e_k^j is a central $(n(1-1/q), 1)$ unit of restrict type supported on B_k and

$$\inf \left(\sum_{k=0}^{\infty} |\lambda_k^j| \right) \sim \|N_j\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)}.$$

Thus,

$$\begin{aligned} I_{2,1} &= \sum_{j=-\infty}^0 \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \|(N_j * b_j)(\cdot) \chi_l(\cdot)\|_{L^q(\mathbb{R}^n)} \\ &\leq \sum_{j=-\infty}^0 \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \sum_{k=0}^{\infty} |\lambda_k^j| \cdot \|(e_k^j * b_j) \chi_l\|_{L^q(\mathbb{R}^n)} \\ &= \sum_{j=-\infty}^0 \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \sum_{k=0}^{\max\{l-2,0\}} |\lambda_k^j| \cdot \|(e_k^j * b_j) \chi_l\|_{L^q(\mathbb{R}^n)} \\ &\quad + \sum_{j=-\infty}^0 \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \sum_{k=\max\{l-1,0\}}^{\infty} |\lambda_k^j| \cdot \|(e_k^j * b_j) \chi_l\|_{L^q(\mathbb{R}^n)} \\ &\equiv I_{2,1}^1 + I_{2,1}^2. \end{aligned}$$

By (iii) in Lemma 3 with $d = n + \varepsilon$, $0 < \varepsilon < 1$, we have

$$\begin{aligned} \|(e_k^j * b_j) \chi_l\|_{L^q(\mathbb{R}^n)} &\leq C 2^j \|e_k^j\|_{L^1(\mathbb{R}^n)} \left(\int_{A_l} |x|^{-dq} dx \right)^{1/q} \\ &\leq C 2^j 2^{-l(n+\varepsilon)} 2^{ln/q}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} I_{2,1}^1 &\leq C \sum_{j=-\infty}^0 2^j \sum_{l=j+3}^{\infty} 2^{-l\varepsilon} \sum_{k=0}^{\infty} |\lambda_k^j| \\ &\leq C \sum_{j=-\infty}^0 2^j 2^{-j\varepsilon} \|N_j\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)} \\ &\leq C \sum_{j=-\infty}^0 2^{j(1-\varepsilon)} \leq C. \end{aligned}$$

By (iii) in Lemma 3 again, we have

$$\begin{aligned} \|(e_k^j * b_j) \chi_l\|_{L^q(\mathbb{R}^n)} &\leq \|e_k^j\|_{L^1(\mathbb{R}^n)} \|b_j\|_{L^q(\mathbb{R}^n)} \leq C 2^j |B(0, 2^k)|^{-(1-1/q)} \\ &\leq C 2^j 2^{-kn(1-1/q)}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} I_{2,1}^2 &\leq C \sum_{j=-\infty}^0 2^j \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \sum_{k=\max\{l-1,0\}}^{\infty} |\lambda_k^j| 2^{-kn(1-1/q)} \\ &\leq C \sum_{j=-\infty}^0 2^j \sum_{k=0}^{\infty} |\lambda_k^j| 2^{-kn(1-1/q)} \sum_{l=-\infty}^{k+1} 2^{nl(1-1/q)} \\ &\leq C \sum_{j=-\infty}^0 2^j \|N_j\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)} \leq C. \end{aligned}$$

Hence, we obtain $I_{2,1} \leq C$.

We now estimate $I_{2,2}$. Let $x \in A_l$, $l \geq j+3$. Then,

$$\begin{aligned} (N_j * b_j)(x) &= \int_{|y| \leq 2^{l-1}} N_j(y) b_j(x-y) dy \\ &\quad + [(N_j \chi_{\tilde{A}_l}) * b_j](x) + \int_{|y| > 2^{l+2}} N_j(y) b_j(x-y) dy, \end{aligned}$$

where $\tilde{A}_l = A_{l-1} \cup A_l \cup A_{l+1}$. Note that if $|y| \leq 2^{l-2}$ and $x \in A_l$, $l \geq j+3$, then $|x-y| \geq 2^{j+1}$ and $|x-y| \geq |x|/2$. Thus, it follows from (ii) in Lemma 3 that

$$|b_j(x-y)| \leq C_d 2^{-nj(1-1/q)} |x|^{-d} \leq C_d 2^{-nj(1-1/q)} 2^{-ld}.$$

Also, note that if $|y| > 2^{l+3}$ and $x \in A_l$, $l \geq j+3$, then $|x-y| \geq 2^{j+1}$ and $|x-y| \geq |y|/2$. Also, it follows from (ii) in Lemma 3 that

$$|b_j(x-y)| \leq C_d 2^{-nj(1-1/q)} |y|^{-d} \leq C_d 2^{-nj(1-1/q)} 2^{-ld}.$$

Thus, when $x \in A_l$, $l \geq j+3$, we have

$$\begin{aligned} |(N_j * b_j)(x)| &\leq C_d 2^{-nj(1-1/q)} 2^{-ld} \|N_j\|_{L^1(\mathbb{R}^n)} + |(N_j \chi_{\tilde{A}_l}) * b_j(x)| \\ &\leq C_d 2^{-nj(1-1/q)-ld} + |(N_j \chi_{\tilde{A}_l}) * b_j(x)|. \end{aligned}$$

Applying these estimates and (i) in Lemma 3 with $d = n + \varepsilon$ to $I_{2,2}$, we obtain

$$\begin{aligned} I_{2,2} &= \sum_{j=1}^{\infty} \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} \|(N_j * b_j)(\cdot) \chi_l(\cdot)\|_{L^q(\mathbb{R}^n)} \\ &\leq C \sum_{j=1}^{\infty} \sum_{l=j+3}^{\infty} 2^{nl(1-1/q)} 2^{-nj(1-1/q)} 2^{-ld} 2^{ln/q} \\ &\quad + \sum_{j=1}^{\infty} \sum_{l=j+3}^{\infty} 2^{ln(1-1/q)} \|(N_j \chi_{\tilde{A}_l}) * b_j(\cdot) \chi_l(\cdot)\|_{L^q(\mathbb{R}^n)} \\ &\leq C \sum_{j=1}^{\infty} 2^{-j(d-n/q)} + \sum_{j=1}^{\infty} \sum_{l=j+3}^{\infty} 2^{ln(1-1/q)} \|N_j \chi_{\tilde{A}_l}\|_{L^1(\mathbb{R}^n)} \|b_j\|_{L^q(\mathbb{R}^n)} \\ &\leq C + C \sum_{j=1}^{\infty} 2^{-nj(1-1/q)} \|N_j\|_{K_1^{n(1-1/q),1}(\mathbb{R}^n)} \leq C + C \sum_{j=1}^{\infty} 2^{-nj(1-1/q)} \leq C. \end{aligned}$$

Now, (9) follows from the above estimates on I_1 , $I_{2,1}$ and $I_{2,2}$.

Actually, it is easy to prove that (9) is true for any central $(1, q)$ atom. That is, the inequality

$$(10) \quad \|T_m \tilde{a}\|_{\dot{K}_q(\mathbb{R}^n)} \leq C$$

holds for any central $(1, q)$ atom $\tilde{a}$. In fact, let us assume that $\text{supp } \tilde{a} \subset B(0, r)$. Obviously, there exists a $k_0 \in \mathbb{Z}$ such that $2^{k_0} < r \leq 2^{k_0+1}$. If I_1 and I_2 in the proof of (9) are now replaced by

$$\sum_{k=-\infty}^{k_0+2} 2^{kn(1-1/q)} \|(T_m \tilde{a}) \chi_k\|_{L^q(\mathbb{R}^n)}$$

and

$$\sum_{k=k_0+3}^{\infty} 2^{kn(1-1/q)} \|(T_m \tilde{a}) \chi_k\|_{L^q(\mathbb{R}^n)}$$

respectively, then (10) can be proved by a method similar to that of proving (9).

To prove (7), by the characterization of $\dot{H}\dot{K}_q(\mathbb{R}^n)$ in terms of Riesz transforms (see [Y]), it suffices to show

$$(11) \quad \sum_{j=1}^n \|R_j(T_m a)\|_{\dot{K}_q(\mathbb{R}^n)} \leq C,$$

where C is independent of a and R_j is the j -th Riesz transform. Since Riesz transforms are bounded on $\dot{H}\dot{K}_q(\mathbb{R}^n)$ (see [Y]), we have

$$R_j a(x) = \sum_k \lambda_k^j a_k^j(x)$$

and

$$\sum_k |\lambda_k^j| \leq C \|R_j a\|_{\dot{H}\dot{K}_q(\mathbb{R}^n)} \leq C,$$

where each a_k^j is a central $(1, q)$ atom and C is independent of a . Thus, it follows from (10) that

$$\begin{aligned} \sum_{j=1}^n \|R_j(T_m a)\|_{\dot{K}_q(\mathbb{R}^n)} &= \sum_{j=1}^n \left\| \sum_k \lambda_k^j (T_m a_k^j) \right\|_{\dot{K}_q(\mathbb{R}^n)} \\ &\leq \sum_{j=1}^n \sum_k |\lambda_k^j| \cdot \|T_m a_k^j\|_{\dot{K}_q(\mathbb{R}^n)} \leq C \sum_{j=1}^n \sum_k |\lambda_k^j| \leq C. \end{aligned}$$

Thus, (11) holds. This completes the proof of Theorem 1. $\square$

Let us now point out that if a linear operator T commutes with translations, then the boundedness of T on $\dot{H}\dot{K}_q(\mathbb{R}^n)$ implies its boundedness on $H^1(\mathbb{R}^n)$. Precisely, we have

Theorem 2. *Let T be a linear operator that commutes with translations. If T is bounded on $\dot{H}\dot{K}_q(\mathbb{R}^n)$, $1 < q < \infty$, then T is also bounded on $H^1(\mathbb{R}^n)$.*

Proof. By the atomic decomposition of $H^1(\mathbb{R}^n)$ (see [CW]), it suffices to prove that the inequality

$$\|Ta\|_{H^1(\mathbb{R}^n)} \leq C$$

holds for any $(1, q)$ atom a . Let $a(x)$ be a $(1, q)$ atom supported on $B(x_0, r)$. That is, $a(x)$ satisfies the following conditions: $\text{supp } a \subset B(x_0, r)$, $\|a\|_{L^q(\mathbb{R}^n)} \leq |B(x_0, r)|^{1/q-1}$, and $\int_{\mathbb{R}^n} a(x)dx = 0$. Let $\bar{a}(x) \equiv \tau_{-x_0}a(x) = a(x+x_0)$. It is easy to see that $\bar{a}$ is a central $(1, q)$ atom supported on $B(0, r)$. Thus, from the conditions of the theorem, it follows that

$$\|T\bar{a}\|_{\dot{HK}_q(\mathbb{R}^n)} \leq C,$$

where C is independent of $\bar{a}$. Since T commutes with translations, we have

$$\begin{aligned} \|\tau_{-x_0}Ta\|_{H^1(\mathbb{R}^n)} &\leq \|\tau_{-x_0}Ta\|_{\dot{HK}_q(\mathbb{R}^n)} = \|T\tau_{-x_0}a\|_{\dot{HK}_q(\mathbb{R}^n)} \\ &= \|T\bar{a}\|_{\dot{HK}_q(\mathbb{R}^n)} \leq C. \end{aligned}$$

Thus, $\tau_{-x_0}Ta \in H^1(\mathbb{R}^n)$ and

$$\tau_{-x_0}Ta(x) = \sum_j \lambda_j a_j(x),$$

where each a_j is a $(1, q)$ atom, and $\sum_j |\lambda_j| \sim \|\tau_{-x_0}Ta\|_{H^1(\mathbb{R}^n)}$. Since $H^1(\mathbb{R}^n)$ is translation invariant, we then have $Ta \in H^1(\mathbb{R}^n)$ and

$$\|Ta\|_{H^1(\mathbb{R}^n)} \leq \sum_j |\lambda_j| \leq C.$$

This finishes the proof of Theorem 2. $\square$

Note that if $q \in (1, \infty)$, then $\alpha = n(1 - 1/q) \in (0, n)$. As a simple corollary of Theorem 1 and Theorem 2, we have

Corollary 2. *Let $0 < \varepsilon < n$. If m satisfies*

$$\sup_{\delta} \|\widehat{m_{\delta}}\|_{K_1^{\varepsilon, 1}(\mathbb{R}^n)} < \infty,$$

then m is a bounded multiplier of $H^1(\mathbb{R}^n)$.

Finally, we point out that it is still an open problem whether (6) is a necessary condition for an $L^\infty(\mathbb{R}^n)$ function m to be a bounded multiplier of $\dot{HK}_q(\mathbb{R}^n)$ in any sense (see [BS]). And we will discuss the similar problems of multipliers on general Herz type Hardy spaces $\dot{HK}_q^{\alpha, p}(\mathbb{R}^n)$ in a future paper.

ACKNOWLEDGMENT

We thank the referee very much for his many helpful comments. We also thank Professor J. Marshall Ash very much for his kindness and enthusiasm.

REFERENCES

- [BS] A. Baernstein II and E. T. Sawyer, *Embedding and Multiplier Theorem for $H^p(\mathbb{R}^n)$* , Memoirs of Amer. Math. Soc., vol. 53, no. 318, Amer. Math. Soc., Providence R. I., 1985. MR **86g**:42036
- [CL] Y. Z. Chen and K. S. Lau, *On some new classes of Hardy spaces*, J. Funct. Anal. **84** (1989), 255-278. MR **90f**:46059
- [CW] R. R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569-645. MR **56**:6264
- [FS] C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), 137-193. MR **56**:6263
- [GR] J. García-Cuerva, *Hardy spaces and Beurling algebras*, J. London Math. Soc. (2) **39** (1989), 499-513. MR **90i**:42032
- [LY1] S. Z. Lu and D. C. Yang, *The Littlewood-Paley function and φ -transform characterizations of a new Hardy space HK_2 associated with the Herz space*, Studia Math. **101** (1992), 285-298. MR **93b**:42031
- [LY2] S. Z. Lu and D. C. Yang, *The decomposition of the weighted Herz spaces on $\mathbb{R}^n$ and its applications*, Sci. in China (Ser. A) **38** (1995), 147-158. MR **96c**:46026
- [LY3] S. Z. Lu and D. C. Yang, *The weighted Herz-type Hardy space and its applications*, Sci. in China (Ser. A) **38** (1995), 662-673. MR **96i**:42018
- [Y] D. C. Yang, *Some Hardy Spaces Associated with the Herz Spaces*, Doctoral Dissertation, Beijing Normal University, Beijing, 1992.

DEPARTMENT OF MATHEMATICS, BEIJING NORMAL UNIVERSITY, BEIJING 100875, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `lusz@bnu.edu.cn`

E-mail address: `dcyang@bnu.edu.cn`