ABSOLUTE FIXED POINT SETS FOR MULTI-VALUED MAPS

ERIC L. MCDOWELL

(Communicated by Alan Dow)

ABSTRACT. The notion of a multi-valued absolute fixed point set (MAFS) will be defined and characterized in the setting of set-valued maps with images containing multiple components.

1. INTRODUCTION

The notion of an absolute fixed point set, in the setting of single-valued maps, was first studied in [1], [2], and [3]; in [3], Martin characterized the class of finite dimensional absolute fixed point sets as the class of finite dimensional absolute retracts. The notion of an absolute fixed point set was extended in [4] to the context of continuum-valued maps and a completely inherent characterization for the notion was obtained. In this paper, we relax the requirement that the maps be continuum-valued and we obtain a characterization for the resulting notion.

A *compactum* is a nonempty compact metric space, and a *continuum* is a connected compactum. If Z is a compactum, then $C(Z)$ denotes the space of all subcontinua of Z with the Hausdorff metric and 2^Z denotes the space of all subcompacta of Z with the Hausdorff metric [5]. A *map* is a continuous function, and a map into some 2^Z will be referred to as a *multi-valued map*; in particular, a map into some $C(Z)$ will be said to be *continuum-valued*.

A *fixed point of a multi-valued map* F is a point, p , such that $p \in F(p)$; the *fixed point set of* F is the set of all fixed points of F and is denoted by $\mathcal{F}(F)$. For a compactum Z , we use the phrase *multi-valued fixed point set of* Z to refer to a subset, A , of Z such that A is the fixed point set of some multi-valued map $F : Z \rightarrow 2^Z$. In case that F is continuum-valued, we call A a *continuum-valued fixed point set of* Z .

The notion that was studied in [4] is defined as follows. A *multi-valued absolute fixed point set* (MAFS) is a compactum, A , such that whenever A is embedded as a subspace, A' , of any compactum, Z , then A' is a continuum-valued fixed point set of Z . It was shown that a compactum A is a MAFS if and only if A contains only finitely many components, all but at most one of which are locally connected [4, 1.1].

Received by the editors January 9, 1997 and, in revised form, April 17, 1997.

1991 *Mathematics Subject Classification*. Primary 54F15, 54C60; Secondary 54C15, 54B20, 54D05.

Key words and phrases. Compactum, continuum, hyperspace, locally connected, multi-valued absolute fixed point set, MAFS, multi-valued map.

Let Z be a compactum, and let $n \in \{1, 2, \dots, \omega\}$ where ω denotes the first infinite cardinal number. We define $C_n(Z)$ for each n as follows:

$$C_n(Z) = \{A \in 2^Z : A \text{ has at most } n \text{ components}\}.$$

In particular, if $A \in C_\omega(Z)$, then A has at most countably many components.

The notion that we study in this paper may now be defined as follows. An n -MAFS is a compactum, A , such that whenever A is embedded as a subspace, A' , of any compactum, Z , then there is some map $F : Z \rightarrow C_n(Z)$ for which $\mathcal{F}(F) = A'$.

We now state the characterizations that we will prove.

1.1 Theorem. *Let A be a compactum, and let $n \in \{1, 2, \dots\}$. Then, A is an n -MAFS if and only if A has only finitely many components, all but at most n of which are locally connected.*

1.2 Theorem. *Let A be a compactum. Then, A is an ω -MAFS if and only if A has at most countably many components.*

We remark that the notions of MAFS and 1-MAFS are identical, and that the characterization given in 1.1 for every finite n is analogous to that given for the class of MAFS. Therefore, it seems natural to consider an inductive argument for 1.1. However, such a proof seems to be elusive.

2. PROOF OF THEOREM 1.1

The following lemmas will be used several times throughout this paper; the proof of 2.1 is similar that given for 1.43 in [5] and will be omitted.

2.1 Lemma. *Let Z be a compactum and let $\Lambda \subseteq 2^Z$. If Λ is connected and $\Lambda \cap C_n(Z) \neq \emptyset$, then $\bigcup \Lambda \in C_n(Z)$, where $n \in \{1, 2, \dots, \omega\}$.*

The proof of the following lemma is similar to that given for 3.1 in [4] and will be omitted.

2.2 Lemma. *Every retract of an n -MAFS is an n -MAFS for $n \in \{1, 2, \dots, \omega\}$.*

The result in 2.3 will be used in 2.4 to provide a necessary condition for a compactum to be an n -MAFS for any finite n . A second necessary condition will be given in 2.5.

2.3 Proposition. *A simple sequence is not an n -MAFS for any $n \in \{1, 2, \dots\}$.*

Proof. Let $J = \{(0, 0, z) \in \mathbf{R}^3 : 0 \leq z \leq 2\}$ and $J' = \{(0, 0, z) \in \mathbf{R}^3 : 0 \leq z \leq 1\}$. Define $W_0 = \{(x, 0, 1 + \sin \frac{1}{x}) : 0 < x \leq 1\}$. For each $i = 1, 2, \dots$, define W_i to be a $\sin \frac{1}{x}$ curve in $\mathbf{R}^3$ whose height near the z -axis is half its height elsewhere, and arrange these curves around the z -axis in such a way that they converge to W_0 . Specifically, define W_i to be an isometric copy of

$$\left\{ \left(x, \frac{1}{2} \left(1 + \sin \frac{1}{x} \right) \right) : 0 < x \leq \frac{2}{(4i+3)\pi} \right\} \cup \left\{ \left(x, 1 + \sin \frac{1}{x} \right) : \frac{2}{(4i+3)\pi} \leq x \leq 1 \right\}$$

in $\mathbf{R}^3$ such that

- (i) $W_i \cap W_j = \emptyset$ whenever $i \neq j$ ($i, j \in \{0, 1, 2, \dots\}$),
- (ii) $\overline{W_i} - W_i = J'$ ($i \in \{1, 2, \dots\}$),
- (iii) $\lim W_i = W_0$.

Define $Z = J \cup \bigcup_{i=0}^{\infty} W_i$. For convenience, we will refer to a point $(0, 0, a)$ as a , and we will use interval notation in the natural way to denote connected subsets of J ; in particular, 2 denotes $(0, 0, 2)$, and $[0, 2] = J$.

Let p_i denote the point of W_i that corresponds to $\frac{2}{(4i-1)\pi}$ for each $i = 1, 2, \dots$; observe that there is a subset of W_i with height 2 between p_i and the z -axis. Set $A = \{0\} \cup \{p_1, p_2, \dots\}$. Then A is a simple sequence in Z with limit point 0. Let $F : Z \rightarrow C_n(Z)$ be any map from Z to $C_n(Z)$. We will show that A cannot be the fixed point set of F .

For the purpose of contradiction, suppose that $\mathcal{F}(F) = A$. Observe that if there was some $x \in (0, 2]$ for which some component of $F(x)$ was contained above x in J , then the continuity of F would imply that some point above x is fixed by F . However, this is impossible by the assumption that $\mathcal{F}(F) \cap J = \{0\}$. It follows that

- (1) no component of $F(x)$ lies above x in J for all $x \in (0, 2]$.

Now, suppose that

$$(*) \quad J \subseteq F(0).$$

Then, by the continuity of F , there is some $\epsilon \in (0, 1)$ for which

- (2) some subset of $F(x)$ lies near J for each $x \in [0, \epsilon]$.

Note also that since $(0, \epsilon) \cap A = \emptyset$, we have that

- (3) $J' \not\subseteq F(x)$ for any $x \in (0, \epsilon)$.

Choose any $x \in (0, \epsilon)$. By (1) and (2), at least one component of $F(x)$ lies in $\bigcup_{i=0}^{\infty} W_i$. Also, since $F(x) \in C_n(Z)$, (3) gives that there are only $k \leq n$ many members of $\{W_i : i = 0, 1, 2, \dots\}$ which meet $F(x)$, say $W_{i_1}, W_{i_2}, \dots, W_{i_k}$. Let y be any other member of $(0, \epsilon)$. Using the continuity of F , we see that if $F(y) \cap W_j \neq \emptyset$ for some $j \notin \{i_1, \dots, i_k\}$, then $J' \subseteq F(z)$ for some $z \in [x, y]$, contradicting (3). It follows that $F(x) \subseteq J' \cup W_{i_1} \cup \dots \cup W_{i_k}$ for every $x \in (0, \epsilon)$. Therefore, we have by continuity that $F(0) \subseteq J' \cup \overline{W_{i_1}} \cup \dots \cup \overline{W_{i_k}}$. So, from $(*)$ we conclude that $i_j = 0$ for some $j \in \{1, \dots, k\}$. Thus, (3) gives that for each $x \in (0, \epsilon)$, some component of $F(x)$ is properly contained in W_0 . It follows from the continuity of F that there is some $x_0 \in W_0$ that lies between J and some component of $F(x_0)$ in W_0 ; we conclude that W_0 contains a fixed point of F , contrary to our supposition that $\mathcal{F}(F) = A$. Therefore, $(*)$ cannot occur.

Suppose on the other hand that

$$(**) \quad J \not\subseteq F(0).$$

Observe that if $2 \in F(0)$, then $(**)$ implies that some component of $F(0)$ lies strictly above 0 in J ; however, it would then follow by continuity that some component of $F(x)$ lies strictly above x for some $x > 0$. Since this would contradict (1), we must have that $2 \notin F(x)$ for every $x \in J$. Therefore, the continuity of F gives that each $x \in J$ is contained in an open subset, U_x , of Z for which $2 \notin F(y)$ for every $y \in U_x$. Thus, $U = \bigcup_{x \in J} U_x$ is an open subset of Z containing J for which

- (4) $2 \notin F(y)$ for every $y \in U$.

Observe that U contains a subcontinuum, Y , such that $J \subseteq Y$ and $p_i \in Y$ for all i sufficiently large. Since F is continuous, we have that $F(Y)$ is connected; thus, $\bigcup F(Y)$ belongs to $C_n(Z)$ by 2.1. Furthermore, our assumption that $\mathcal{F}(F) = A$ implies that $\bigcup F(Y)$ contains p_i for i sufficiently large. Thus, since $\bigcup F(Y)$ contains at most $n < \infty$ components, some component, K , of $\bigcup F(Y)$ must contain p_i for

infinitely many i . Since K is a closed connected subset of $\bigcup F(Y)$ that contains infinitely many members of $\{p_i : i = 1, 2, \dots\}$, it follows from our choice of p_i that $[0, 2] \subseteq K$, contrary to (4). Hence, $(**)$ cannot occur. Since we can have neither $(*)$ nor $(**)$, we must conclude that our supposition that $\mathcal{F}(F) = A$ is false. So, A cannot be the fixed point set of any $F : Z \rightarrow C_n(Z)$. This proves 2.3.

2.4 Corollary. *Every n -MAFS contains at most finitely many components for each $n \in \{1, 2, \dots\}$.*

Proof. Let $n \in \{1, 2, \dots\}$. If A is a compactum with infinitely many components, then A contains a simple sequence as a retract [4, 3.2]. It follows immediately from 2.2 and 2.3 that A is not an n -MAFS.

2.5 Proposition. *Let $n \in \{1, 2, \dots\}$. If A is an n -MAFS, then all but at most n components of A must be locally connected.*

Proof. By 2.2 and 2.4, it suffices to show that

- (*) if A is the disjoint union of $n + 1$ nonlocally connected continua, then A is not an n -MAFS.

Let $A_1, \dots, A_{n+1}$ denote the components of A . Since A_1 is not locally connected at some $a_\infty \in A_1$, there exists a neighborhood, V , of a_∞ , and points $a_i \in V$, $i = 1, 2, \dots$, for which $a_\infty = \lim a_i$ and such that

- (1) no two of $a_\infty, a_1, a_2, \dots$ lie in the same component of V ,
- (2) $C_\infty = \lim C_i$, where C_i denotes the component of a_i in V for each $i \in \{1, 2, \dots, \infty\}$.

Let $S = \{(x, \sin \frac{1}{x}) : 0 < x \leq 1\}$. For each $i \in \{1, 2, \dots\}$ and each $j \in \{1, 2, \dots, i\}$, let $f_{ij} : S \rightarrow \mathbf{R}^3$ be an embedding for which

- (3) $\overline{f_{ij}(S)} - f_{ij}(S) = \{0\} \times \{0\} \times [0, \frac{1}{j}] = I_j$,
- (4) $f_{ij}(S) \subset \{(x, y) : x^2 + y^2 \leq \frac{1}{k}\} \times I_j$, where $k = \frac{i(i-1)}{2} + j$,
- (5) $f_{ij}(S) \cap f_{i'j'}(S) \neq \emptyset$ if and only if $i = i'$ and $j = j'$.

In other words, $\{f_{ij}(S)\}$ is a sequence of $\sin \frac{1}{x}$ curves with decreasing widths and alternating heights. The first term, $f_{11}(S)$, fits inside the tube centered on the z -axis with radius 1, while the fifth term, $f_{32}(S)$ ($5 = \frac{3(3-1)}{2} + 2$), fits inside the tube centered on the z -axis with radius $\frac{1}{5}$; also, $f_{11}(S)$ has height 1, and the heights of $f_{31}(S)$, $f_{32}(S)$, and $f_{33}(S)$ have heights 1, $\frac{1}{2}$, and $\frac{1}{3}$ respectively. Define $p = (0, 0, 0)$ and $q_j = (0, 0, \frac{1}{j})$ for each $j \in \{1, 2, \dots\}$. For each $i \in \{1, 2, \dots\}$ and each $j \in \{1, \dots, i\}$, let $p_{ij} = p_k$ be a point of $f_{ij}(S)$ such that $p = \lim p_i$, where $k = \frac{i(i-1)}{2} + j$. Define $Y_1 = \bigcup_{j \leq i} \overline{f_{ij}(S)}$. Without loss of generality, we

may assume that A_1 and Y_1 are disjoint. Let $h_i : Y_1 \rightarrow \mathbf{R}^3$ be an embedding for each $i = 1, 2, \dots, n + 1$ such that $h_1(Y_1) = Y_1$, $h_i(Y_1) \cap h_j(Y_1) = \emptyset$ whenever $i \neq j$, and $h_i(Y_1) \cap A = \emptyset$ for all $i = 1, 2, \dots, n + 1$. Set $Y_i = h_i(Y_1)$ for each $i = 1, 2, \dots, n + 1$. Form Z_1 from A_1 and Y_1 by identifying a_∞ with p , and a_k with p_k for each $k = 1, 2, \dots$ and define Z_i from A_i and Y_i in a similar manner. Let T be a simple $(n + 1)$ -od [6, 9.8] with vertex t and endpoints $t_1, \dots, t_{n+1}$ such that $T \cap \bigcup_{i=1}^{n+1} Z_i = \emptyset$. Define Z to be the continuum obtained by identifying each t_i with the image of $h_i(p) \in Z_i$. Then Z is a continuum that contains a homeomorphic copy of A . We will use A , V , C_i , I_i , Y_i , and T to denote the natural copies of these spaces in Z .

Suppose that there is some map $F : Z \rightarrow C_n(Z)$ with $\mathcal{F}(F) = A$. Note that since $F(t)$ has at most n components, we may assume, without loss of generality, that $F(t)$ shares no point in common with the component of Z_1 in $Z - \{t\}$. It follows from the continuity of F , and the supposition that $\mathcal{F}(F) = A$, that

$$(6) \quad F(p) \cap Z_1 = \{p\}.$$

By (6) and the continuity of F , there is some open set U of Z with $p \in U$ for which

$$(7) \quad Z_1 \cap F(x) \subseteq V \cup Y_1 \text{ for each } x \in U.$$

It is clear from (3) that U contains I_{j_0} for some $j_0 \in \{1, 2, \dots\}$; indeed, it is clear from (4) that U contains $\overline{f_{ij_0}(S)}$ for all sufficiently large i . Note that from (6), the continuity of F , and the supposition that $\mathcal{F}(F) = A$, we have that $q_{j_0} \notin F(x)$ for any $x \in I_{j_0}$. Thus, for each $x \in I_{j_0}$, there is an open set, $V_x \subseteq U$, such that $q_{j_0} \notin F(y)$ for every $y \in V_x$. So, $U' = \bigcup_{x \in I_{j_0}} V_x$ is an open subset of U for which

$$(8) \quad q_{j_0} \notin F(y) \text{ for every } y \in U'.$$

Let E be a subcontinuum of U' that contains p_{ij_0} for all sufficiently large i . Then $F(E)$ is a subcontinuum of $C_n(Z)$, and so $\bigcup F(E)$ belongs to $C_n(Z)$ by 2.1. Since each p_{ij_0} belongs to $\mathcal{F}(F)$, it follows that some component, K , of $\bigcup F(E)$ must contain infinitely many members of $\{p_{ij_0} : i = 1, 2, \dots\}$. Hence, K must also contain p . Observe that $K \subseteq V \cup Y_1$ by (7). So by (1), $\overline{K - V}$ is a subcontinuum of Y_1 that contains p and infinitely many members of $\{p_{ij_0} : i = 1, 2, \dots\}$. But, by (5), any subcontinuum of Y_1 that contains both p and some p_{ij_0} must also contain I_{j_0} . Thus, $q_{j_0} \in K$, contrary to (8). So, we must conclude that our supposition that $\mathcal{F}(F) = A$ is false. This shows (*), as required. Thus, we have proven 2.5.

The following proposition asserts that the properties described in 2.4 and 2.5 are, together, sufficient for a compactum to be an n -MAFS.

2.6 Proposition. *Let Z be a compactum, and let $n \in \{1, 2, \dots\}$. If A is a compactum in Z that contains only finitely many components, all but at most n of which are locally connected, then $A = \mathcal{F}(F)$ for some map $F : Z \rightarrow C_n(Z)$.*

Proof. Let $A = A_1 \cup A_2 \cup \dots \cup A_m$, where A_i is a component of A for each $i \in \{1, \dots, m\}$. Note that if $m \leq n$, then $F : Z \rightarrow C_n(Z)$, given by $F(z) = A$ for every $x \in Z$, is a map with $\mathcal{F}(F) = A$. So we assume that $m > n$. Without loss of generality, we also assume that $A_{n+1}, \dots, A_m$ are locally connected. It follows from the characterization of the class of MAFS [4, 1.1] that there is a map $G : Z \rightarrow C(Z)$ with $\mathcal{F}(G) = A_n \cup A_{n+1} \cup \dots \cup A_m$. Define $F : Z \rightarrow C_n(Z)$ by

$$F(x) = G(x) \cup A_1 \cup \dots \cup A_{n-1}$$

for each $x \in Z$. Then F is a map with $\mathcal{F}(F) = A$. This proves 2.6.

The theorem in 1.1 has now been proved by 2.4, 2.5, and 2.6.

3. PROOF OF THEOREM 1.2

3.1 Proposition. *Every compactum with at most countably many components is an ω -MAFS.*

Proof. Let Z be a compactum, and let A be a compact subset of Z with at most countably many components. Then $F : Z \rightarrow C_\omega(Z)$ defined by $F(x) = A$ for all $x \in Z$ is a map with $\mathcal{F}(F) = A$. This proves 3.1.

The rest of the work in this section is directed toward demonstrating that having only countably many components is also a necessary condition for a compactum to be an ω -MAFS.

3.2 Lemma. *An ω -MAFS cannot be both totally disconnected and uncountable.*

Proof. Let A be an uncountable, totally disconnected compactum. Without loss of generality, we may assume that A is a subset of the circle in $\mathbf{R}^3$ given by

$$\{(x, y, 1 + \sin 1) : x^2 + y^2 = 1\}.$$

Let $U_1 \supset U_2 \supset \cdots$ be a strictly decreasing sequence of uncountable, simultaneously open and closed subsets of A such that $U_1 = A$ and $\bigcap_{i=1}^{\infty} U_i = \{p\}$ for some $p \in A$. Define J , J' and W_i ($i \in \{1, 2, \dots\}$) as we did in the proof of 2.3. Now, for each $a \in A - \{p\}$, observe that there is precisely one $i_a \in \{1, 2, \dots\}$ for which $a \in U_{i_a} - U_{i_a+1}$; let $h_a : W_{i_a} \rightarrow \mathbf{R}^3$ be the isometric embedding for which

$$\begin{aligned} h_a((1, 1 + \sin 1)) &= a, \\ \overline{h_a(W_{i_a})} - h_a(W_{i_a}) &= J'. \end{aligned}$$

Let $S = \{(x, 1 + \sin \frac{1}{x} : 0 < x \leq 1\}$, and let $h_p : S \rightarrow \mathbf{R}^3$ be the isometric embedding for which

$$\begin{aligned} h_p((1, 1 + \sin 1)) &= p, \\ \overline{h_p(S)} - h_p(S) &= J. \end{aligned}$$

Let $Z = \overline{h_p(S)} \cup \bigcup_{a \in A - \{p\}} h_a(W_{i_a})$. Now, for each $a \in A$, define

$$x_a = \begin{cases} h_a\left(\frac{2}{(4i_a-1)\pi}\right), & \text{if } a \neq p, \\ (0, 0, 0), & \text{if } a = p. \end{cases}$$

Observe that the map given by $a \mapsto x_a$ is an embedding of A into Z . Set $A' = \{x_a : a \in A\}$, and suppose that there exists some $F : Z \rightarrow C_\omega(Z)$ with $\mathcal{F}(F) = A'$. This supposition leads to a contradiction by an argument similar to the one given for 2.3. This proves 3.2.

The following technical lemma will facilitate the proof of 3.4.

3.3 Lemma. *Let Z be a compactum, and let V be an open subset of Z which meets uncountably many components of Z . Then, for each $n \in \{1, 2, \dots\}$ there exist points $p_1, \dots, p_n$ of V which lie in n distinct components of Z , and for which any open set containing p_i meets uncountably many components of Z for each $i = 1, \dots, n$.*

Proof. Suppose that the conclusion fails for $n = 1$; that is, suppose that

(*) for every $p \in V$, there is some open subset of V that contains p and meets at most countably many components of Z .

Let $\mathcal{C}$ denote the collection of all components of Z that meet V . For each $C \in \mathcal{C}$, let $z_C \in C \cap V$; define $A = \{z_C : C \in \mathcal{C}\}$. Then, by hypothesis, we have that

(1) A is uncountable.

Now, let $\{V_i\}_{i=1}^{\infty}$ be a sequence of open subsets of V for which $cl_Z(V_i) \subseteq V$ and for which $\bigcup_{i=1}^{\infty} V_i = V$. Clearly we have that

(2) $\bigcup_{i=1}^{\infty} (cl_Z(V_i) \cap A) = A$.

Therefore, since the union of countably many countable sets is countable, it follows immediately from (1) and (2) that $cl_Z(V_k) \cap A$ is uncountable for some $k \in \{1, 2, \dots\}$. Setting $E = cl_Z(V_k)$ shows that

- (3) E is a closed subset of Z in V that meets uncountably many components of Z .

By (*) and the compactness of E , there exist finitely many open subsets, $U_1, \dots, U_m$, of V for which $E \subseteq \bigcup_{i=1}^m U_i$ and such that each U_i meets at most countably many components of Z . But this implies that E meets at most countably many components of Z , contrary to (3). Hence, the supposition in (*) is false. Therefore, the lemma holds for $n = 1$.

Now suppose that the lemma holds for $n = j$. Let $p_1, \dots, p_j$ be points of V that satisfy the conclusion of the lemma, and let K_i denote the component of p_i in Z for each $i \in \{1, \dots, j\}$. Set $K = \bigcup_{i=1}^j K_i$. Then $V - K$ is an open subset of Z that meets uncountably many components of Z . So, by the case for $n = 1$, there is some point $p_{j+1} \in V - K$ for which any open set containing p_{j+1} meets uncountably many components of Z . This completes the proof of 3.3.

The following notation will facilitate the proof of our next result. Let $\{0, 1\}^{<\omega}$ denote the set of all finite sequences with range in $\{0, 1\}$, that is,

$$\{0, 1\}^{<\omega} = \{\langle a_1, \dots, a_n \rangle : n = 1, 2, \dots; a_i = 0, 1; i = 1, \dots, n\} \cup \{\emptyset\}.$$

For any $s \in \{0, 1\}^{<\omega}$, let $|s|$ denote the length of s . Let $s0$ and $s1$ denote those elements of $\{0, 1\}^{<\omega}$ that have length $|s| + 1$, that agree with s in the first $|s|$ positions, and that have 0 and 1, respectively, in the final position. If $|s| = 1$, set $s^- = \emptyset$; otherwise, let s^- denote that element of $\{0, 1\}^{<\omega}$ of length $|s| - 1$ that agrees with s in the first $|s| - 1$ positions. Finally, let $\{0, 1\}^\omega$ denote the set of all countably infinite sequences with range in $\{0, 1\}$.

3.4 Lemma. *Let Z be a compactum with uncountably many components. Then Z has a retract which is both uncountable and totally disconnected.*

Proof. Using 3.3, there exist distinct components, $K_{\langle 0 \rangle}$ and $K_{\langle 1 \rangle}$, of Z , such that for some $p_{\langle i \rangle} \in K_{\langle i \rangle}$, every open set about $p_{\langle i \rangle}$ meets uncountably many components of Z . Using the fact that every component of Z is also a quasicomponent of Z [6, 5.18], let $U_{\langle i \rangle}$ be a simultaneously closed and open (clopen) subset of Z for each $i = 0, 1$ such that $K_{\langle i \rangle} \subseteq U_{\langle i \rangle}$, $U_{\langle 0 \rangle} \cap U_{\langle 1 \rangle} = \emptyset$, and $U_{\langle 0 \rangle} \cup U_{\langle 1 \rangle} \neq Z$. Choose any point, $p_\emptyset$, in $Z - (U_{\langle 0 \rangle} \cup U_{\langle 1 \rangle})$, and define $f_\emptyset : Z - (U_{\langle 0 \rangle} \cup U_{\langle 1 \rangle}) \rightarrow Z$ by

$$f_\emptyset(x) = p_\emptyset \text{ for all } x \in Z - (U_{\langle 0 \rangle} \cup U_{\langle 1 \rangle}).$$

We will now construct U_s , V_s , K_s , p_s , and f_s for each $s \in \{0, 1\}^{<\omega}$ by induction on $\{0, 1\}^{<\omega}$. Set $V_\emptyset = U_\emptyset = Z$, and let $s \in \{0, 1\}^{<\omega}$. Let V_s be an open set about p_s such that

- (1) the diameter of V_s is $\leq 2^{-|s|}$,
- (2) $V_s \subseteq U_s \cap V_{s^-}$.

By applying 3.3 to U_s , we can find distinct components, K_{s0} and K_{s1} , of U_s (hence, of Z), for which

- (3) $K_{si} \neq K_s$ for $i = 0, 1$,
- (4) there is some $p_{si} \in K_{si} \cap V_s$ such that every open set about p_{si} meets uncountably many components of Z .

Let U_{si} be a clopen subset of Z for each $i = 0, 1$ such that

- (5) $U_{si} \subseteq U_s$,
- (6) $K_{si} \subseteq U_{si}$,
- (7) $U_{s0} \cap U_{s1} = \emptyset$,
- (8) $U_{si} \cap K_s = \emptyset$.

Then, define $f_s : U_s - (U_{s0} \cup U_{s1}) \rightarrow Z$ by

$$f_s(x) = p_s \text{ for all } x \in U_s - (U_{s0} \cup U_{s1}).$$

Observe that since each U_{si} is open, (4) and (6) give that U_{si} contains uncountably many components of Z for each $i = 0, 1$. This completes our construction. Now, for each $r \in \{0, 1\}^\omega$, let r_n denote the unique element of $\{0, 1\}^{<\omega}$ of length n for which $r_n(k) = r(k)$ for all $k \in \{1, \dots, n\}$. Let $Z' = \bigcup \{\bigcap_{n=1}^\infty U_{r_n} : r \in \{0, 1\}^\omega\}$. It is easy to verify that

- (9) $x \in Z - Z'$ if and only if $x \in U_s - (U_{s0} \cup U_{s1})$ for some $s \in \{0, 1\}^{<\omega}$.

Since $U_s - (U_{s0} \cup U_{s1})$ is open for every $s \in \{0, 1\}^{<\omega}$, (9) gives that

- (10) Z' is a closed subset of Z .

Define $f' : Z - Z' \rightarrow Z$ by

$$f'(x) = f_s(x) \text{ if } x \in U_s - (U_{s0} \cup U_{s1}), \quad s \in \{0, 1\}^{<\omega}.$$

Note that f' is well defined by (5), (7), and (9). Furthermore, since f_s is constant for each $s \in \{0, 1\}^{<\omega}$ and each $U_s - (U_{s0} \cup U_{s1})$ is open in Z , it is clear that

- (11) f' is continuous.

For each $r \in \{0, 1\}^\omega$, use (1) and (2) to define $p_r = \bigcap_{n=1}^\infty \overline{V_{r_n}}$. Then, define $f : Z \rightarrow Z$ by

$$f(x) = \begin{cases} f'(x) & \text{if } x \in Z - Z', \\ p_r, & \text{if } x \in \bigcap_{n=1}^\infty U_{r_n} \text{ for some } n. \end{cases}$$

Again, f is well defined by (5), (7), and (9).

By (10) and (11), we have that f is continuous at each point of $Z - Z'$. So, to see that f is continuous on Z , it is enough to show that f is continuous at each $x \in Z'$. Let $x_0 \in Z'$, and let $\{x_i\}_{i=1}^\infty$ be a sequence in Z with $\lim x_i = x_0$. By the definitions of Z' and f , there is some $r \in \{0, 1\}^\omega$ for which $f(x_0) = p_r$. Let V be an open subset of Z about $f(x_0) = p_r$. Then, by (1), (2), and the definition of p_r , V contains V_{r_N} for some $N \in \{1, 2, \dots\}$. By the definition of f , we have that $x_0 \in U_{r_{N+1}}$; hence, $x_i \in U_{r_{N+1}}$ for all i sufficiently large. But, by (2), (4), and the definitions of f_s , p_r , and f , it follows that $f(U_{r_{N+1}}) \subseteq V_{r_N}$. So $f(x_i) \in V$ for all i sufficiently large; therefore, f is continuous at x_0 , as required.

Finally, observe that by (2), (3), (4), and (8), $f(Z) \cap K$ contains at most one point from each component, K , of Z ; hence, we have that $f(Z)$ is totally disconnected. Furthermore, it is clear from our construction that $f(Z)$ is a retract of Z with uncountably many elements. This proves 3.4.

3.5 Proposition. *Every ω -MAFS contains at most countably many components.*

Proof. This follows immediately from 2.2, 3.2 and 3.4.

The theorem in 1.2 has now been proved by 3.1 and 3.5.

3.6 Remark. Let $\mathfrak{c}$ denote the cardinality of $\mathbf{R}$. By defining a $\mathfrak{c}$ -MAFS in the natural way, it is easily seen that every compactum is a $\mathfrak{c}$ -MAFS since every compactum has at most $\mathfrak{c}$ components.

3.7 Problem. Characterize the class of n -MAFS, $n \in \{1, 2, \dots, \omega\}$, in the setting of compact Hausdorff spaces.

ACKNOWLEDGMENTS

The results in this paper are a part of the author's doctoral dissertation, *Absolute Fixed Point Sets for Multi-Valued Maps*, completed at West Virginia University under the direction of Sam B. Nadler, Jr. The author wishes to thank Professor Nadler for his enthusiastic support and assistance.

REFERENCES

1. John R. Martin, *On absolute fixed-point sets*, Colloq. Math. **35** (1976), 67–71. MR **53**:4033
2. ———, *Absolute fixed-point sets in compacta*, Colloq. Math. **39** (1977), 41–44. MR **80f**:54028
3. ———, *Absolute fixed point sets and AR-spaces*, Fund. Math. **112** (1981), 159–164. MR **82j**:54098
4. Eric L. McDowell and Sam B. Nadler, Jr., *Absolute fixed point sets for continuum-valued maps*, Proc. Amer. Math. Soc. **124** (1996), 1271–1276. MR **96g**:54042
5. Sam B. Nadler, Jr., *Hyperspaces of sets*, Marcel Dekker Inc., New York, 1978. MR **58**:18330
6. ———, *Continuum theory, an introduction*, Marcel Dekker, Inc., New York, 1992. MR **93m**:54002

DEPARTMENT OF MATHEMATICS, BETHANY COLLEGE, BETHANY, WEST VIRGINIA 26032

E-mail address: e.mcdowell@mail.bethanywv.edu

Current address: Department of Mathematical Sciences, Berry College, Mount Berry, Georgia 30149

E-mail address: emcdowell@berry.edu