A MAXIMUM PRINCIPLE FOR P -HARMONIC MAPS WITH L^q FINITE ENERGY

KENSHO TAKEGOSHI

(Communicated by Peter Li)

ABSTRACT. We show a maximum principle for P -harmonic maps with L^q -finite energy. As an application we can generalize a non-existence theorem for harmonic maps with finite Dirichlet integral by Schoen and Yau to those maps.

1. INTRODUCTION

Let $f : (M, g) \rightarrow (N, h)$ be a smooth map of Riemannian manifolds. We are interested in determining under which condition the boundedness of the length $|df|$ of df can be induced. This problem is related to the maximum principle for solutions of an elliptic operator of second order in a sense. For instance several Schwarz type lemmas for harmonic maps, which state the boundedness of $|df|$ under a certain negative curvature condition of the target manifold (N, h) , are known and a few Liouville type theorems are induced from them (cf. [C-Y], [E-L], [G-H], [Sh], [Y-2]). As a degenerate case of the negativity condition of the curvature of (N, h) the following observation is interesting in view of our aspect (see also [C]).

Theorem*. *Let $f : (M, g) \rightarrow (N, h)$ be a smooth map from a non-compact complete (connected) Riemannian manifold (M, g) of dimension m into a Riemannian manifold (N, h) . Let Γ_M (resp. Γ_N) denote the set of points of M (resp. N) where Ric_M (resp. Ric_N) is not non-negative (resp. not non-positive). Let Σ and E be subsets of M such that $\Sigma := \Gamma_M \cup f^{-1}(\Gamma_N) \cup E$ and $M \setminus \overline{\Sigma} \neq \emptyset$. Suppose $f : (M \setminus \overline{\Sigma}, g) \rightarrow (N, h)$ is harmonic, i.e. $\text{Trace}_g \nabla(df) = 0$ on $M \setminus \overline{\Sigma}$, and $\int_{M \setminus \overline{\Sigma}} |df|^q < +\infty$ for $q > 1$. Then the following holds:*

$$|df|(x) \leq \sup_{y \in \Sigma} |df|(y) \quad \text{for any } x \in M,$$

where the right hand side of the above inequality may take infinity and is defined to be zero if $\Sigma = \emptyset$.

To show this result the subharmonicity of $|df|$ on $M \setminus \overline{\Sigma}$ plays an essential role in view of the Weitzenböck formula. In fact Theorem* can be obtained by [Sc-Y], Theorem 1, and applying the following general statement to $|df|$ (see also [L-S]).

Received by the editors April 21, 1997.

1991 *Mathematics Subject Classification.* Primary 58D15, 58E20.

Proposition. *Let (M, g) be as above and let v be a smooth function on M satisfying $\Delta v := \text{Trace}_g \nabla^2 v \geq 0$ outside a proper subset A of M . Suppose the interior of $\{v \leq \sup_A v\}$ is not empty and $\int_{M \setminus A} |dv^q| < +\infty$ with $q \geq 1$. Then $v(x) \leq \sup_A v$ for any $x \in M$.*

For the proof the reader should see [Y-1], Corollary and the proof of Theorem 3 (see also §2). Here the L^1 -integrability condition of dv^q is essential and cannot be removed.

For a positive number $p > 1$ $f : (M, g) \rightarrow (N, h)$ is said to be p -harmonic if f satisfies the following:

$$\text{Trace}_g \nabla(|df|^{p-2} df) = 0 \quad \text{on } M.$$

As in the case of a harmonic map, i.e. $p = 2$, any p -harmonic map $f : (M, g) \rightarrow (N, h)$ can be characterized as a critical point of the L^p energy functional $\int_M |df|^p$ if it is finite. In this note we generalize Theorem* to p -harmonic maps. However we cannot apply the above proposition to the case $p \neq 2$ because the subharmonicity of $|df|^2$ breaks down. Nevertheless we will show the following by modifying an idea developed in [T].

Theorem. *Let $f : (M, g) \rightarrow (N, h)$ and Σ be as in Theorem*. Suppose $f : (M \setminus \overline{\Sigma}, g) \rightarrow (N, h)$ is p -harmonic, i.e. $\text{Trace}_g \nabla(|df|^{p-2} df) = 0$ on $M \setminus \overline{\Sigma}$, and $\int_{M \setminus \overline{\Sigma}} |df|^q < +\infty$ for $q > p - 1$ with $p \geq 2$. Then the following holds:*

$$|df|(x) \leq \sup_{y \in \Sigma} |df|(y) \quad \text{for any } x \in M.$$

This result can be regarded as a maximum principle for L^q integrable solutions of a non-linear degenerate elliptic equation. In particular, we obtain the following (cf. [L-S], [N], [Sc-Y]).

Corollary. *Let (M, g) and (N, h) be a non-compact complete (connected) Riemannian manifold with non-negative Ricci curvature and a Riemannian manifold with non-positive Riemannian curvature respectively. Then any p -harmonic map $f : (M, g) \rightarrow (N, h)$ with L^q finite energy is constant if $q > p - 1$ with $p \geq 2$ ($q \geq 1$ when $p = 2$).*

2. PROOF OF THE THEOREM

By the Weitzenböck formula (cf. [E-L], (2.20) and (3.13)) we obtain the following:

$$\frac{1}{2} \Delta |df|^2 = |\nabla(df)|^2 - \langle df, (DD^* + D^*D)(df) \rangle + Q(df),$$

where D^* is the adjoint operator of

$$D : \Gamma\left(\bigwedge^s TM^* \otimes f^*TN\right) \rightarrow \Gamma\left(\bigwedge^{s+1} TM^* \otimes f^*TN\right)$$

and $Q(df)$ is given by the following:

$$Q(df) := \sum_{i=1}^m \langle \text{Ric}_M(df(e_i)), df(e_i) \rangle - \sum_{j,k=1}^m \langle \text{Riem}_N(df(e_j), df(e_k)) df(e_k), df(e_j) \rangle.$$

Since $Q(df) \geq 0$ on $M \setminus \overline{\Sigma}$ by the curvature condition and $D(df) = 0$, we obtain for any real number q

$$(1) \quad \begin{aligned} & \frac{1}{2} |df|^{q-2} \Delta |df|^2 + |df|^{q-2} \langle df, DD^*(df) \rangle \\ & \geq |df|^{q-2} |\nabla(df)|^2 \quad \text{on } M \setminus (\overline{\Sigma} \cup \{|df| = 0\}). \end{aligned}$$

If $c_0 := \sup_{y \in \Sigma} |df|(y) = +\infty$, then the assertion is trivial. We have only to consider the case $0 \leq c_0 < +\infty$. Let $M = M_* \cup M_+$, $M_* := \{|df| \leq c_0\}$, and $M_+ := \{|df| > c_0\}$. Assume $M_+ \neq \emptyset$. First we take a smooth non-negative function λ on a real line satisfying $\lambda \equiv 0$ on $(-\infty, c_0^2]$, $\lambda > 0$, $\lambda' \geq 0$ on $(c_0^2, +\infty)$ and $\lambda \equiv 1$ on $[c_1, +\infty)$, $c_1 > c_0^2$. By the completeness of g , for a fixed point $x_* \in M$ and any $r > 0$ there exists a non-negative Lipschitz continuous function ω on M such that $0 \leq \omega \leq 1$ on M , $\omega \equiv 1$ on $B_r(x_*)$, $\omega \equiv 0$ on $M \setminus B_{2r}(x_*)$, and $|d\omega| \leq C/r$ on M for $C > 0$ not depending on r , where $B_r(x_*)$ is the geodesic ball centered at $x_* \in M$ of radius $r > 0$. Setting $u := |df|$ it should be noted that $\overline{\Sigma} \cup \{u = 0\} \subset M_*$, u is smooth on M_+ , and $\lambda(u(x)^2) > 0$ if and only if $x \in M_+$. By (1) we have

$$(2) \quad \begin{aligned} \int \lambda(u^2) \omega^2 u^{q-2} |\nabla(df)|^2 & \leq \frac{1}{2} \int \lambda(u^2) \omega^2 u^{q-2} \Delta u^2 \\ & \quad + \int \lambda(u^2) \omega^2 u^{q-2} \langle df, DD^*(df) \rangle. \end{aligned}$$

Using integration by parts we have

$$(*) \quad \begin{aligned} \int \lambda(u^2) \omega^2 u^{q-2} \Delta u^2 & = - \int \langle \nabla(\lambda(u^2) \omega^2 u^{q-2}), \nabla u^2 \rangle \\ & = - \int \lambda'(u^2) \omega^2 u^{q-2} |\nabla u^2|^2 - 2 \int \lambda(u^2) \omega u^{q-2} \langle \nabla \omega, \nabla u^2 \rangle \\ & \quad - \int \lambda(u^2) \omega^2 \langle \nabla u^{q-2}, \nabla u^2 \rangle \\ & \leq \frac{2}{\varepsilon} \int \lambda(u^2) \omega^2 u^q |\nabla \omega|^2 - 2(q-2-\varepsilon) \int \lambda(u^2) \omega^2 u^{q-2} |\nabla u|^2 \end{aligned}$$

for any $q > 0$ and $\varepsilon > 0$. Here we remark the following:

$$D^*(vu^{p-2}df) = vD^*(u^{p-2}df) - u^{p-2}\mathbf{e}(dv)^*df$$

on $\{u > 0\}$ for any real number p and any smooth function v on M , where $\mathbf{e}(dv)^*$ is the adjoint operator of the left exterior multiplication by dv . In particular, by the p -harmonicity of f , i.e. $D^*(u^{p-2}df) = 0$, on M_+ we obtain

$$(3) \quad D^*(vu^{q-2}df) = -u^{p-2}\mathbf{e}(d(vu^{q-p}))^*df,$$

$$(4) \quad D^*(df) = (p-2)\mathbf{e}(d \log u)^*df$$

on M_+ for any $p > 1$ and $q > 0$. Using integration by parts, and the Cauchy-Schwarz inequalities $|\mathbf{e}(du)^*df| \leq |\nabla u|u$ and $|D^*(df)| \leq \sqrt{m}|\nabla(df)|$, we obtain

$$\begin{aligned}
 \int \lambda(u^2)\omega^2 u^{q-2} \langle df, DD^*(df) \rangle &= \int \langle D^*(\lambda(u^2)\omega^2 u^{q-2}df), D^*(df) \rangle \\
 &= - \int \left\{ 2\lambda'(u^2)u^{q-1} + (q-p)u^{q-3}\lambda(u^2) \right\} \omega^2 \langle \mathbf{e}(du)^*df, D^*(df) \rangle \\
 &\quad - 2 \int \lambda(u^2)\omega u^{q-2} \langle \mathbf{e}(d\omega)^*df, D^*(df) \rangle \quad \text{by (3)} \\
 (**) \quad &\leq -\min\{0, (q-p)(p-2)\} \int \lambda(u^2)u^{q-2}\omega^2 |\nabla u|^2 \\
 &\quad + 2 \int \lambda(u^2)\omega u^{q-1} |\nabla \omega| |D^*(df)| \quad \text{by (4)} \\
 &\leq -\min\{0, (q-p)(p-2)\} \int \lambda(u^2)u^{q-2}\omega^2 |\nabla u|^2 \\
 &\quad + \varepsilon \int \lambda(u^2)\omega^2 u^{q-2} |\nabla(df)|^2 + \frac{m}{\varepsilon} \int \lambda(u^2)u^q |\nabla \omega|^2
 \end{aligned}$$

for any $q > 0$ and $\varepsilon > 0$. Since $|\nabla u| \leq |\nabla(df)|$ on $\{u > 0\}$, by (2), (*), and (**), we obtain for any $q > 0$ and $0 < \varepsilon \leq 1$

$$\begin{aligned}
 &(q-1 + \min\{0, (q-p)(p-2)\} - 2\varepsilon) \int \lambda(u^2)\omega^2 u^{q-2} |\nabla u|^2 \\
 &\leq \frac{m+1}{\varepsilon} \int \lambda(u^2)u^q |\nabla \omega|^2.
 \end{aligned}$$

If $q > p-1$ with $p \geq 2$, then $\varepsilon := \min\{1, q-1 + \min\{0, (q-p)(p-2)\}\}/3 > 0$. Since $\lambda' \geq 0$ and $\sup \lambda' = 1$, we obtain

$$\int_{B_r(x_*)} \lambda(u^2)u^{q-2} |\nabla u|^2 \leq \frac{(m+1)C^2}{\varepsilon^2 r^2} \int_{M \setminus \overline{\Sigma}} u^q.$$

By the L^q integrability condition of u on $M \setminus \overline{\Sigma}$ and letting r tend to infinity we obtain

$$\int_M \lambda(u^2)u^{q-2} |\nabla u|^2 = 0.$$

Since $\lambda \geq 0$ and $\lambda(u(x)^2) > 0$ if and only if $x \in M_+$, the above equality implies $\nabla u \equiv 0$ on M_+ , i.e. $u = |df| \equiv \text{constant} > c_0$ on M_+ by the connectedness of M . If $\Sigma \neq \emptyset$, then by continuity there exists a point $x_* \in M_*$ such that $|df|(x_*) > c_0$, which is a contradiction and so M_+ is empty. If $\Sigma = \emptyset$, i.e. $c_0 = 0$, then $u \equiv c > 0$ on M . Hence by [Y-1], Theorem 7, $\int_M |df|^q = c^q \times \text{Vol}(M) = +\infty$, which is also a contradiction and so M_+ is empty, i.e. f is constant. This completes the proof of Theorem. In case of $p = 2$ and $q = 1$, the assertion of Corollary follows from [L-S], Theorem 2.2, (b).

REFERENCES

- [C] Cheng, S.Y., *Liouville theorem for harmonic maps*, Proceedings of Symposia in Pure Mathematics **36** (1980), 147-151. MR **81i**:58021
- [C-Y] Cheng, S.Y., Yau, S.T., *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math. **28** (1975), 333-354. MR **52**:6608
- [E-L] Eells, J., Lemaire, L., *A report on harmonic maps*, Bull. London Math. Soc. **10** (1978), 1-68. MR **82b**:58033

- [G-H] Goldberg, S.I., Har'el, Z., *A general Schwarz lemma for Riemannian manifolds*, Bull. Soc. Math. Grèce **18** (1977), 141-148. MR **80c**:53055
- [L-S] Li, P., Schoen, R., *L^p and mean value properties of subharmonic functions on Riemannian manifolds*, Acta Math. **153** (1984), 279-301. MR **86j**:58147
- [N] Nakauchi, N., *A Liouville type theorem for p -harmonic maps*, preprint.
- [Sc-Y] Schoen, R., Yau, S.T., *Harmonic maps and the topology of stable hypersurfaces and manifolds of nonnegative Ricci curvature*, Comment. Math. Helv. **51** (1976), 333-341. MR **55**:11302
- [Sh] Shen, C-L., *A generalization of the Schwarz-Ahlfors lemma to the theory of harmonic maps*, J. Reine Angew. Math. **348** (1984), 23-33. MR **85i**:58044
- [T] Takegoshi, K., *A volume estimate for strong subharmonicity and maximum principle on complete Riemannian manifolds*, to appear in Nagoya Mathematical Journal.
- [Y-1] Yau, S.T., *Some function theoretic properties of complete Riemannian manifolds and their applications to geometry*, Indiana Univ. Math. J. **25** (1976), 656-670. MR **54**:5502
- [Y-2] Yau, S.T., *A general Schwarz lemma for Kähler manifolds*, Amer. J. Math. **100** (1978), 197-203. MR **58**:6370

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, MACHIKANNEYAMA-CHO 1-16,
TOYONAKA-SHI OSAKA, 560 JAPAN

E-mail address: kensho@math.wani.osaka-u.ac.jp