ON AN ANALOGUE OF SELBERG'S EIGENVALUE CONJECTURE FOR $\mathrm{SL}_3(\mathbf{Z})$

SULTAN CATTO, JONATHAN HUNTLEY, JAY JORGENSEN, AND DAVID TEPPER

(Communicated by Dennis A. Hejhal)

ABSTRACT. Let $\mathcal{H}$ be the homogeneous space associated to the group $\mathrm{PGL}_3(\mathbf{R})$. Let $X = \Gamma \backslash \mathcal{H}$ where $\Gamma = \mathrm{SL}_3(\mathbf{Z})$ and consider the first nontrivial eigenvalue λ_1 of the Laplacian on $L^2(X)$. Using geometric considerations, we prove the inequality $\lambda_1 > 3\pi^2/10$. Since the continuous spectrum is represented by the band $[1, \infty)$, our bound on λ_1 can be viewed as an analogue of Selberg's eigenvalue conjecture for quotients of the hyperbolic half space.

§1. STATEMENT OF THE MAIN THEOREM

A fundamental question in the spectral theory of automorphic forms is whether small eigenvalues exist. More specifically, let G be a noncompact reductive group with finite center, Γ a nonuniform lattice, K a maximal compact subgroup of G , and set $X = \Gamma \backslash G/K$. It is well known from the theory of Eisenstein series that $L^2(X)$ has continuous spectrum for the ring of invariant differential operators, and in particular for the positive Laplacian, Δ . The continuous spectrum will be, in cases of interest such as $\mathrm{PGL}_n(\mathbf{R})$, an interval $[a, \infty)$ with $a > 0$. The question we referred to above is: Do nonconstant square integrable eigenforms exist with eigenvalue $\lambda < a$? This problem is important for various considerations in number theory. In the case $G = \mathrm{PGL}_2(\mathbf{R})$ and Γ is a congruence subgroup, Selberg conjectured that no such nontrivial small eigenvalues exist.

In this paper, we consider the case when

$$G = \mathrm{PGL}_3(\mathbf{R}) \quad \text{and} \quad \Gamma = \mathrm{SL}_3(\mathbf{Z}).$$

Our main result is the following.

Theorem. *Let λ_1 be the eigenvalue for the first nontrivial eigenform on $L^2(X)$. Then*

$$\lambda_1 > 3\pi^2/10 > 2.96088.$$

Received by the editors January 28, 1997.

1991 *Mathematics Subject Classification.* Primary 11F55; Secondary 22E40, 11F72.

The first named author acknowledges support from DOE grants DE-AC-0276-ER3074 and 3075 and PSC-CUNY Research Award No. 9203393.

The second named author acknowledges support from several PSC-CUNY grants. The third named author acknowledges support from NSF grant DMS-93-07023 and from the Sloan Foundation.

§2. NOTATION

Let $\mathcal{H} = G/K$ and set $X = \Gamma \backslash \mathcal{H}$. Explicit coordinates for $\tau \in \mathcal{H}$ via the Iwasawa decomposition are given by

$$\tau = \begin{bmatrix} 1 & x_2 & x_3 \\ 0 & 1 & x_1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_1 y_2 & 0 & 0 \\ 0 & y_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

with $y_1, y_2 > 0$, from which one can compute that the (positive) Laplacian Δ can be written as

$$\begin{aligned} -\Delta = & y_1^2 \frac{\partial^2}{\partial y_1^2} - y_1 y_2 \frac{\partial^2}{\partial y_1 \partial y_2} + y_2^2 \frac{\partial^2}{\partial y_2^2} + y_1^2 (x_2^2 + y_2^2) \frac{\partial^2}{\partial x_3^2} \\ & + y_1^2 \frac{\partial^2}{\partial x_1^2} + y_2^2 \frac{\partial^2}{\partial x_2^2} + 2y_1^2 x_2 \frac{\partial^2}{\partial x_1 \partial x_3} \end{aligned}$$

(see pages 17 and 33 of [Bu84]). The ring of invariant differential operators is spanned by the identity, the Laplacian Δ , and a third order operator Δ_3 (see [Bu84]). The invariant volume element is given by

$$dV = \frac{dx_1 dx_2 dx_3 dy_1 dy_2}{y_1^3 y_2^3}.$$

We shall not use the explicit formula for the invariant volume element; however, the above expression for the Laplacian will be necessary in our proof of the main theorem.

For our purposes, it is more convenient to work with functions on $\mathcal{H}$ that are $\mathrm{SL}_3(\mathbf{Z})$ invariant rather than considering functions on the quotient space X . To this end, we introduce a fundamental domain for $\Gamma \backslash \mathcal{H}$. Specifically, computations on page 56 of [Gr93] show that a fundamental domain $\mathcal{D}$ is described through the following set of inequalities:

$$\begin{aligned} v^{\frac{3}{2}} &< v^{\frac{3}{2}}(1 - x_2 + x_3)^2 + w(1 - x_1)^2 + w^{-1}; \\ v^{\frac{3}{2}} &< v^{\frac{3}{2}}(x_2 - x_3)^2 + w(1 - x_1)^2 + w^{-1}; \\ v^{\frac{3}{2}} &< v^{\frac{3}{2}}x_2^2 + w; \quad v^{\frac{3}{2}} < v^{\frac{3}{2}}x_3^2 + wx_1^2 + w^{-1}; \\ 1 &< w^{-2} + x_1^2; \quad 0 < x_1 < \frac{1}{2}; \quad 0 < x_2 < \frac{1}{2}; \quad -\frac{1}{2} < x_3 < \frac{1}{2}, \end{aligned}$$

where we have used the notation $w^{-1} = y_1$ and $v^{-\frac{3}{2}} = y_2^2 y_1$. Let S denote the Siegel set described via the inequalities

$$0 < x_1 < \frac{1}{2}, \quad 0 < x_2 < \frac{1}{2}, \quad -\frac{1}{2} < x_3 < \frac{1}{2}, \quad y_1 > \frac{\sqrt{3}}{2}, \quad y_2 > \frac{\sqrt{3}}{2}.$$

The set S contains the fundamental domain $\mathcal{D}$. Further, results from page 61 of [Gr93] show the existence of elements $\gamma_1, \dots, \gamma_{10} \in \mathrm{SL}_3(\mathbf{Z})$ such that $S \subset \bigcup_{i=1}^{10} \mathcal{D}\gamma_i$ (we have used the notation $\mathcal{D}\gamma$ to denote the image of the fundamental domain $\mathcal{D}$ under left multiplication by γ). The main aspects of the above points which we shall use are the assertions that for any $\tau \in S$ we have $y_1(\tau) > \sqrt{3}/2$ and that S is contained in ten translates of $\mathcal{D}$.

Recall that an automorphic form is a C^∞ function ϕ on $\mathcal{H}$ which satisfies the following properties:

- (i) $\phi(\gamma \circ \tau) = \phi(\tau)$ for $\gamma \in \mathrm{SL}_3(\mathbf{Z})$;
- (ii) $|\phi(\tau)| \ll y_1^{N_1} y_2^{N_2}$ for $\tau \in \mathcal{D}$ and integers N_1, N_2 ;
- (iii) ϕ is an eigenform for the ring of invariant differential operators.

An automorphic form is said to be a cusp form if it satisfies the additional property

$$(iv) \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \phi \left(\begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) d\xi_1 d\xi_3 = \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \phi \left(\begin{bmatrix} 1 & \xi_2 & \xi_3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) d\xi_2 d\xi_3 = 0.$$

Cusp forms are square integrable. Although we shall not need this fact, let us note that, from the theory of Eisenstein series, the only noncuspidal square integrable automorphic forms on X are constant.

§3. PROOF OF THE MAIN THEOREM

Our method of proof is a modification of that used by Roelcke to show that the small eigenvalue λ_1 for the quotient space $\mathrm{SL}_2(\mathbf{Z}) \backslash \mathrm{SL}_2(\mathbf{R}) / \mathrm{SO}_2(\mathbf{R})$ satisfies the bound $\lambda_1 > 3\pi^2/2$ (see page 511 of [He83]). We shall use the Fourier expansion of automorphic forms associated to $\mathrm{SL}_3(\mathbf{Z})$, as developed in Chapter IV of [Bu84].

Assume that ϕ is a nonconstant automorphic form, so then $\Delta\phi = \lambda\phi$ and $\phi\Delta\phi = \lambda\phi^2$. Through integration by parts, using the automorphic boundary conditions, and the fact that the Siegel domain S is contained in ten translates of the fundamental domain $\mathcal{D}$, we obtain the inequality

$$\frac{\int_S |\nabla\phi|^2 dV}{\int_S |\phi|^2 dV} < 10\lambda.$$

As on page 67 of [Bu84], let us expand ϕ in a Fourier expansion with respect to the abelian group

$$\left\{ \begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} : \xi_1, \xi_3 \in \mathbf{R} \right\}.$$

Specifically, we have $\phi(\tau) = \sum_{n_1, n_3} \phi_{n_1}^{n_3}(\tau)$ where

$$\phi_{n_1}^{n_3}(\tau) = \int_0^1 \int_0^1 \phi \left(\begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) e^{-2\pi i(n_1 \xi_1 + n_3 \xi_3)} d\xi_1 d\xi_3.$$

Observe that $\phi_0^0 = 0$ since ϕ is not constant and square integrable, hence cuspidal. Let

$$\Gamma_1^2 = \left\{ \begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & 1 \end{bmatrix} : \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbf{Z}) \right\},$$

and set Γ_∞^2 to be the subgroup of Γ_1^2 which stabilizes infinity. As on page 69 of [Bu84], we then can write $\phi_{n_1}^{n_3}$ as

$$\phi_{n_1}^{n_3}(\tau) = \sum_{\gamma \in \Gamma_\infty^2 \backslash \Gamma_1^2} \sum_{n_1=1}^{\infty} \phi_{n_1}^0(\gamma \circ \tau).$$

By a standard application of elliptic regularity, ϕ is necessarily C^∞ , hence we can interchange integration and summation and apply Parseval's theorem to obtain the inequality

$$\frac{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \nabla_\tau \phi_{n_1}^0(\gamma \circ \tau) \right|^2 dV}{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\gamma \circ \tau) \right|^2 dV} < 10\lambda.$$

Since ∇ is an invariant operator, we may differentiate the expressions in the numerator and then evaluate at $\gamma \circ \tau$, thus yielding

$$\frac{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \nabla_\tau \phi_{n_1}^0(\tau) \Big|_{\gamma \circ \tau} \right|^2 dV}{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\tau) \Big|_{\gamma \circ \tau} \right|^2 dV} < 10\lambda.$$

We now integrate by parts and consider the action of the Laplacian Δ on functions of the form $\phi_{n_1}^0$. Since each function $\phi_{n_1}^0$ is independent of x_3 , these terms in Δ annihilate $\phi_{n_1}^0$. Observe that all terms involving y_1, y_2 and x_2 are positive operators (compare with line (2.31) on page 32 of [Bu84]), so we obtain the bound

$$\Delta \phi_{n_1}^0 \geq -y_1^2 \cdot \frac{\partial^2}{\partial x_1^2} \phi_{n_1}^0 = y_1^2 \cdot 4\pi^2 n_1^2 \phi_{n_1}^0.$$

Since $y_1^2 > \frac{3}{4}$, we have $\Delta \phi_{n_1}^0 \geq \frac{3}{4} \cdot 4\pi^2 n_1^2 \phi_{n_1}^0 = 3\pi^2 n_1^2 \phi_{n_1}^0$. Combining this inequality with the above calculations and the cuspidality condition $\phi_0^0 = 0$, we obtain

$$10\lambda > \frac{\int_S \sum_{n_1=1}^{\infty} \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \Delta_\tau \phi_{n_1}^0(\tau) \Big|_{\gamma \circ \tau} \cdot \phi_{n_1}^0(\tau) \Big|_{\gamma \circ \tau} dV}{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\tau) \Big|_{\gamma \circ \tau} \right|^2 dV} \geq 3\pi^2,$$

hence $\lambda \geq 3\pi^2/10$. Since ϕ was any cusp form, we obtain the bound as asserted in the theorem.

§4. CONCLUDING REMARKS

As the continuous spectrum in this situation is $[1, \infty)$, our theorem implies an analogue of Selberg's eigenvalue conjecture. Note that our bound is stronger than the result for $\mathrm{SL}_3(\mathbf{Z})$ from [Mi96] where it was proved that $\lambda_1 \geq 1$. In general, our method applies to $G = \mathrm{SL}_n(\mathbf{R})$ with $\Gamma = \mathrm{SL}_n(\mathbf{Z})$ to give the bound $\lambda_1 > 3\pi^2/M$ where M is the number of fundamental domains which intersect a Siegel set containing the fundamental domain constructed in [Gr93]; however, for $n \geq 4$, this bound is rather weak. Finally, let us remark that our theorem is indeed a consequence of the Ramanujan conjecture, which asserts that all nontrivial automorphic representations come from tempered representations.

REFERENCES

- [Bu84] BUMP, D.: *Automorphic forms on $GL(3, \mathbf{R})$* . Lecture Notes in Mathematics **1083** New York: Springer-Verlag (1984). MR **86g**:11028
- [Gr93] GREINER, D.: On the shape of fundamental domains in $GL(n, R)$. *Pacific Journal Math.* **160** (1993) 53-65.
- [He83] HEJHAL, D. A.: *The Selberg Trace Formula for $PSL(2, \mathbf{R})$, vol. 2*. Lecture Notes in Mathematics **1001** New York: Springer-Verlag (1983). MR **86e**:11040
- [Mi96] MILLER, S.: Spectral and cohomological applications of the Rankin-Selberg method. *Inter. Math. Res. Notices* **1** (1996) 15-26. MR **97a**:11079

(S. Catto) THE GRADUATE SCHOOL AND BARUCH COLLEGE, THE CITY UNIVERSITY OF NEW YORK, NEW YORK, NEW YORK 10010 AND DEPARTMENT OF PHYSICS, THE ROCKEFELLER UNIVERSITY, 1230 YORK AVENUE, NEW YORK, NEW YORK 10021-6339

(J. Huntley and D. Tepper) DEPARTMENT OF MATHEMATICS, BARUCH COLLEGE CUNY, 17 LEXINGTON AVENUE, NEW YORK, NEW YORK 10010

(J. Jorgenson) SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY 08540

Current address: Department of Mathematics, Oklahoma State University, Stillwater, Oklahoma 74078

E-mail address: `jjorgen@littlewood.math.okstate.edu`