A CONVOLUTION ESTIMATE FOR A MEASURE ON A CURVE IN $\mathbb{R}^4$. II

DANIEL M. OBERLIN

(Communicated by Christopher D. Sogge)

ABSTRACT. This paper contains almost-sharp $L^p - L^q$ convolution estimates for measures on the curve (t, t^2, t^3, t^4) in $\mathbb{R}^4$.

Let γ be the curve in $\mathbb{R}^4$ defined by $\gamma(t) = (t, t^2, t^3, t^4)$, $t \in \mathbb{R}$. For an interval $I \subseteq \mathbb{R}$ we consider the convolution operator T on $\mathbb{R}^4$ defined by

$$Tf(x) = \int_I f(x - \gamma(t)) dt, \quad x \in \mathbb{R}^4.$$

Our interest is in determining the type-set $\mathcal{T}$ of T —the set of all $(\frac{1}{p}, \frac{1}{q}) \in [0, 1] \times [0, 1]$ such that $T(L^p(\mathbb{R}^4)) \subseteq L^q(\mathbb{R}^4)$. Let $A = (0, 0)$, $B = (\frac{4}{10}, \frac{3}{10})$, $C = (\frac{5}{10}, \frac{4}{10})$, $D = (\frac{6}{10}, \frac{5}{10})$, $E = (\frac{7}{10}, \frac{6}{10})$, and $F = (1, 1)$. Since T is a convolution operator, $\mathcal{T}$ lies on or below the closed segment $[A, F]$. Estimating Tf when f is the characteristic function of a small ball shows that $\mathcal{T}$ lies on or above the line through A and B and so, by duality, on or below the line through F and E . Estimating Tf when f is the characteristic function of

$$[0, \varepsilon] \times [0, \varepsilon^2] \times [0, \varepsilon^3] \times [0, \varepsilon^4]$$

shows that $\mathcal{T}$ lies on or above the line through B and E . Thus $\mathcal{T}$ lies inside the closed quadrilateral $ABEF$ (the closed convex hull of A , B , E , and F). If the interval of integration I is unbounded, then homogeneity considerations show that $\mathcal{T} \subseteq [B, E]$. It is conjectured that these necessary conditions for the boundedness of T are also sufficient. By duality and interpolation, it would suffice to show that $B \in \mathcal{T}$ if $I = (0, \infty)$. The best previous result [O1] is that $[C, D] \subseteq \mathcal{T}$ for any I . Here we establish an almost sharp result when I is bounded.

Theorem. *If I is bounded, then, excepting possibly the segments $(A, B]$, $[B, C)$, $(D, E]$, and $[E, F)$, the closed quadrilateral $ABEF$ lies in $\mathcal{T}$.*

Proof. Let $G = (\frac{1}{2}, \frac{1}{2})$. By duality, interpolation, and the result of [O1], it is enough to show that $[G, B] \subseteq \mathcal{T}$. And a homogeneity argument shows that it is enough to do this for $I = [0, 1]$. To this end we will study an analytic family T_z of convolution operators given formally by

$$T_z f(x) = \frac{1}{\Gamma(\frac{z+1}{2})} \int_0^1 \int_{-1}^1 f(x - \gamma(t) - u\gamma''(t)) |u|^z du dt.$$

Received by the editors May 12, 1997.

1991 *Mathematics Subject Classification.* Primary 42B15.

The author was partially supported by a grant from the National Science Foundation.

We will establish, with acceptable estimates on the growth of the operator norms, that

$$(1) \quad T_z \text{ is bounded on } L^2(\mathbb{R}^4) \text{ when } \operatorname{Re}(z) = -\frac{3}{2}$$

and, if $H = (\frac{1}{3}, \frac{1}{6})$, that

$$(2) \quad T_z \text{ maps } L^p(\mathbb{R}^4) \text{ into } L^q(\mathbb{R}^4) \text{ if } \left(\frac{1}{p}, \frac{1}{q}\right) \in [G, H) \text{ and } \operatorname{Re} z = -\frac{2}{3}.$$

Analytic interpolation and the fact that T_{-1} is a multiple of T then show that $[G, B) \subseteq \mathcal{T}$ as claimed.

To establish (1) we need to estimate the Fourier transform $\widehat{T}_{-\frac{3}{2}+is}$ given formally by

$$\widehat{T}_{-\frac{3}{2}+is}(y) = \frac{1}{\Gamma((-\frac{1}{2}+is)/2)} \int_0^1 \int_{-1}^1 e^{-iy \cdot (\gamma(t) + u\gamma''(t))} |u|^{-\frac{3}{2}+is} du dt.$$

Because the integral

$$\int_{-1}^1 |u|^{-\frac{2}{3}} du$$

converges, it is enough to estimate the integral

$$(3) \quad \int_0^1 \int_{-\infty}^{\infty} e^{-iy \cdot (\gamma(t) + u\gamma''(t))} |u|^{-\frac{3}{2}+is} du dt.$$

If $y = (y_1, y_2, y_3, y_4)$, write $p(t) = \sum_{j=1}^4 y_j t^j$. Since the Fourier transform of the one-dimensional distribution $|u|^{-\frac{3}{2}+is} du$ is a multiple of $|x|^{\frac{1}{2}-is} dx$, (3) is essentially

$$\int_0^1 e^{-ip(t)} |p''(t)|^{\frac{1}{2}-is} dt.$$

A result of [O3] bounds this integral by $C(1+|s|)^{\frac{1}{2}}$ independently of y . Thus

$$\|\widehat{T}_{-\frac{3}{2}+is}\|_{\infty} \leq C(1+|s|)^{\frac{1}{2}},$$

yielding (1).

To prove (2) we begin by defining

$$\begin{aligned} S_1 f(x) &= \int_0^{\infty} \int_{\frac{1}{2}}^1 f(x - \gamma(t) - u\gamma''(t)) du dt \doteq \mu_1 * f(x), \\ S_2 f(x) &= \int_0^{\infty} \int_{-1}^{-\frac{1}{2}} f(x - \gamma(t) - u\gamma''(t)) du dt \doteq \mu_2 * f(x). \end{aligned}$$

With $S = S_1 + S_2$, our immediate goal is to establish that

$$(4) \quad S \text{ maps } L^3(\mathbb{R}^4) \text{ into } L^6(\mathbb{R}^4).$$

We will do this by adapting an argument from [O2]. Now (4) follows from the two multilinear estimates

$$(5) \quad \left| \int_{\mathbb{R}^4} \prod_{j=1}^6 \mu_{\ell} * f_j(x) dx \right| \leq C \prod_{j=1}^6 \|f_j\|_3, \quad \ell = 1, 2.$$

Utilizing Christ's remarkable observation about multilinear interpolation ([C1], pp. 227–228), (5) will follow from the estimates

$$(6) \quad \left| \int_{\mathbb{R}^4} \prod_{j=1}^6 \mu_\ell * f_j(x) dx \right| \leq C \|f_1\|_1 \prod_{j=2}^6 \|f_j\|_{5,1}, \quad \ell = 1, 2,$$

where the symbol $\|\cdot\|_{p,r}$ stands for a Lorentz space norm on $\mathbb{R}^4$. It is enough to prove (6) with f_1 replaced by the point mass at the origin in $\mathbb{R}^4$ and so (6) reduces to the estimate

$$\|S_\ell f\|_{L^5(\mu)} \leq C \|f\|_{5,1}.$$

We will actually establish the dual of this estimate,

$$(7) \quad \|S_\ell^* g\|_{L^{\frac{5}{4},\infty}} \leq C \|g\|_{L^{\frac{5}{4}}(\mu_\ell)},$$

for functions g on the two-dimensional manifold that is the support of μ_ℓ . Let ϕ be the map of $\mathbb{R}^4$ into itself defined by

$$\phi(t, u, s, v) = \gamma(t) + u\gamma''(t) - \gamma(s) - v\gamma''(s).$$

Computations show that

$$(8) \quad \phi \text{ is at most three to one on the set } \{t \neq s\}$$

and that the absolute value J of the Jacobian of ϕ is given by

$$(9) \quad J(t, u, s, v) = 24(s-t)^2|(s-t)^2 - 6(u+v)|.$$

Let m denote Lebesgue measure on $\mathbb{R}^4$. From (8) it follows that if $U \subseteq \mathbb{R}^4$ is open and if the measure λ_U is defined on $\phi(U)$ by

$$\int_{\phi(U)} h d\lambda_U = \int_U h \circ \phi(t, u, s, v) J(t, u, s, v) d(t, u, s, v),$$

then

$$(10) \quad m(E) \leq \lambda_U(E) \leq 3m(E), \quad E \subseteq U.$$

Let $Q_1 = \{(s, v) : 0 < s < \infty, \frac{1}{2} < v < 1\}$, $Q_2 = \{(s, v) : 0 < s < \infty, -1 < v < -\frac{1}{2}\}$. If f is a function on $\mathbb{R}^4$ and g is defined on the support of μ_ℓ , then we have, writing $g(s, v)$ for $g(\gamma(s) + v\gamma''(s))$,

$$\begin{aligned} \int_{\mathbb{R}^4} f(x) S_\ell^* g(x) dx &= \int_{\mathbb{R}^4} S_\ell f(y) g(y) d\mu_\ell(y) \\ &= \int_{Q_\ell \times Q_\ell} f(\phi(t, u, s, v)) \frac{g(s, v)}{J(t, u, s, v)} J(t, u, s, v) d(t, u, s, v). \end{aligned}$$

The last integral is bounded by the product of the two Lorentz norms

$$\|f \circ \phi\|_{L^{5,1}(Jdm)}, \quad \|g/J\|_{L^{\frac{5}{4},\infty}(Jdm)}.$$

By (10) the first of these is bounded by $\|f\|_{5,1}$. We will establish (7) by observing that there is a constant C such that, for any $\alpha > 0$,

$$\int_{(Q_\ell \times Q_\ell) \cap \{|g(x,v)| > \alpha J(t,u,s,v)\}} J(t, u, s, v) d(t, u, s, v) \leq \frac{C}{\alpha^{\frac{5}{4}}} \int_{Q_\ell} |g(s, v)|^{\frac{5}{4}} d(s, v).$$

This will follow from the inequality, uniform in $A > 0$ and $(s, v) \in Q_\ell$,

$$\int_{Q_\ell \cap \{J(t, u, s, v) < A\}} J(t, u, s, v) d(t, u) \leq CA^{\frac{5}{4}}.$$

Taking account of (9) it is enough to observe that there is C such that for $A > 0$ and $6 < |k| < 12$ we have

$$(11) \quad \int_{\{t^2|t^2-k|<A\}} t^2|t^2-k| dt \leq CA^{\frac{5}{4}}.$$

But this inequality is easily checked by considering separately the cases of large and small A . The first case follows from the fact that if $|t|$ is large, then $t^2|t^2-k| \sim t^4$, leading to (11). In the case of small A , $t^2|t^2-k| < A$ implies either $t^2|t^2-k| \sim Ct^2$ or $t^2|t^2-k| \sim C|t \pm \sqrt{k}|$, leading to estimates of $CA^{\frac{3}{2}}$ or CA^2 respectively for the left-hand side of (11). These complete the proof of (4).

Now define operators V_k by

$$V_k f(x) = \int_0^1 \int_{2^{-k} < |u| < 2^{-k+1}} f(x - \gamma(t) - u\gamma''(t)) du dt.$$

Let $\|V_k\|_{pq}$ denote the norm of V_k from $L^p(\mathbb{R}^4)$ to $L^q(\mathbb{R}^4)$. Then, since

$$T_{-\frac{3}{2}} \sim \sum_{k=1}^{\infty} 2^{\frac{2k}{3}} V_k,$$

(2) will follow from the estimate

$$(12) \quad \|V_k\|_{pq} \leq C2^{-k(1-2\alpha)}$$

for $(\frac{1}{p}, \frac{1}{q})$ on $[G, H]$ with $\frac{1}{p} - \frac{1}{q} = \alpha$ along with the fact that $\alpha < \frac{1}{6}$. Inequality (12) comes from interpolating the cases $\alpha = 0$ (corresponding to $G = (\frac{1}{2}, \frac{1}{2})$) and $\alpha = \frac{1}{6}$ (corresponding to $H = (\frac{1}{3}, \frac{1}{6})$). The first of these two estimates is clear. The second is a consequence of (4) and a homogeneity argument:

For $x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4$ and $\lambda \in \mathbb{R}$, let $[\lambda]x$ be the nonisotropic dilate of x given by $(\lambda x_1, \lambda^2 x_2, \lambda^3 x_3, \lambda^4 x_4)$. For $\lambda > 0$, let $V_{[\lambda]}$ be the dilate of V given by

$$V_{[\lambda]} f(x) = \int_0^\infty \int_{\frac{1}{2} < |u| < 1} f(x - [\lambda](\gamma(t) + u\gamma''(t))) du dt.$$

Then it is easy to see that

$$\|V_{[\lambda]}\|_{pq} = \lambda^{-(\frac{10}{p} - \frac{10}{q})} \|S\|_{pq}$$

and that

$$V_{[\lambda]} f(x) = \lambda^{-3} \int_0^\infty \int_{\frac{\lambda^2}{2} < |u| < \lambda^2} f(x - \gamma(t) - u\gamma''(t)) du dt.$$

Taking $\lambda = 2^{-\frac{k}{2}}$ and combining these two observations with (4) gives (12) for $\alpha = \frac{1}{6}$ (and $(\frac{1}{p}, \frac{1}{q}) = H = (\frac{1}{3}, \frac{1}{6})$).

ADDED IN PROOF

The recent paper [GSW] uses a modification of the method of this paper to show that T is of restricted weak type at the points B and E . And the more recent paper [C2] uses an entirely different (geometric) method to prove the n -dimensional analogue of this last result.

REFERENCES

- [C1] M. Christ, *On the restriction of the Fourier transform to curves: endpoint results and the degenerate case*, Trans. Amer. Math. Soc. **287** (1985), 223–238. MR **87b**:42018
- [C2] M. Christ, *Convolution, Curvature and Combinatorics, a Case Study*, preprint.
- [GSW] A. Greenleaf, A. Seeger, and S. Wainger, *On x-ray transforms for rigid line complexes and integrals over curves in $\mathbb{R}^4$* , preprint.
- [O1] D. Oberlin, *A convolution estimate for a measure on a curve in $\mathbb{R}^4$* , Proc. Amer. Math. Soc. **125** (1997), 1355–1361. MR **97g**:42009
- [O2] D. Oberlin, *Multilinear proofs for two theorems on circular averages*, Colloq. Math. **63** (1992), 187–190. MR **93m**:42005
- [O3] D. Oberlin, *Oscillatory integrals with polynomial phase*, Math. Scand. **69** (1991), 45–56.

DEPARTMENT OF MATHEMATICS, FLORIDA STATE UNIVERSITY, TALLAHASSEE, FLORIDA 32306-4510