THE TRANSVERSAL HOMOCLINIC POINTS ARE DENSE IN THE CODIMENSION-1 HÉNON-LIKE STRANGE ATTRACTORS

YONGLUO CAO

(Communicated by Mary Rees)

ABSTRACT. We consider the codimension-1 Hénon-like strange attractors Λ . We prove that the transversal homoclinic points are dense in Λ , and that hyperbolic periodic points are dense in Λ . Moreover the hyperbolic periodic points that are heteroclinically related to the primary fixed point (transversal intersection of stable and unstable manifolds) are dense in Λ .

1. INTRODUCTION

In 1976 [3] Hénon performed a numerical study of the family of diffeomorphism of the plane $h_{a,b}(x, y) = (1 - ax^2 + y, bx)$ and detected for parameter values $a = 1.4, b = 0.3$, what seemed to be nontrivial attractor with a highly intricate geometric structure. This family has since then been the subject of intense research. But its dynamics are still far from being completely understood.

In a remarkable paper [1], Benedicks and Carleson show that if $b > 0$ is small enough then for a positive measure set of a -values near $a = 2$, the corresponding diffeomorphism $h_{a,b}$ exhibits a strange attractor.

Subsequently, this result has been much extended by Mora and Viana: The Hénon-like attractors appear whenever one generically unfolds a quadratic homoclinic tangency through one-parameter families of locally dissipative surface diffeomorphism [4].

Recently Viana [6] overcame many difficulties that arose when extending the BC method to a higher dimensional setting and showed the existence of codimension-1 Hénon-like strange attractors for unfoldings of codimension-1 homoclinic tangencies in higher dimensions.

In general in a manifold M of dimension 2 or more, the codimension-1 Hénon-like strange attractors Λ of diffeomorphism φ is defined as follows:

1. There is a saddle fixed point $p \in \Lambda$ whose unstable manifold has dimension 1, and Λ is the closure of the unstable manifold of the saddle p .
2. The basin of Λ contains an open set.

Received by the editors January 9, 1997 and, in revised form, September 30, 1997.

1991 *Mathematics Subject Classification.* Primary 58F11, 58F12.

Key words and phrases. Hénon-like strange attractor, transversal homoclinic point.

The author was partially supported by IMPA, TWAS and NSF of China and Jiangsu and “Nonlinear Science” Basic Project.

3. There is a point z_1 on the unstable manifold $w^u(p)$ such that its forward orbit is dense in Λ and $\|(d\varphi^n)_{z_1}(\nu)\| \geq e^{cn} \|\nu\|$ for all $n \geq 0$, some constant $c > 0$ and some nonzero vector ν .

A construction of Sinai-Bowen-Ruelle (SRB) invariant measure for these attractors was given by Benedicks and Young [2].

There exists a direction in the tangent space at z_1 which is exponentially contracted by both positive and negative iterates of $D\varphi$, clearly the presence of such point is an obstruction to (uniform) hyperbolicity of the attractors. It implies the nonhyperbolicity of Λ .

It is well known that for an Axiom-A attractor, by the definition of hyperbolicity, it follows that the transversal homoclinic points are dense in it. Moreover the hyperbolic periodic points are dense in it, and any two periodic points are heteroclinically related (transversal intersection of stable and unstable manifolds).

We know that the codimension-1 Hénon-like strange attractors are nonhyperbolic. In this paper, we prove the following results.

Theorem. *The transversal homoclinic points are dense in the codimension-1 Hénon-like strange attractor.*

Corollary. *The hyperbolic periodic points are dense in the codimension-1 Hénon-like strange attractor. Moreover the hyperbolic periodic points that are heteroclinically related to the primary fixed point also are dense in it.*

This paper is arranged as follows. In Section 2 we give some preliminaries. Section 3 is devoted to the proof of the Theorem.

2. PRELIMINARIES

In this section, we first give some definitions and facts.

First, we define the concept of a Hénon-like map, let $f_a(x) = 1 - ax^2$ and $\hat{f}_a(x, Y) = (1 - ax^2, 0^{m-1})$, where $a \in [0, 3]$, $(x, Y) \in [-2, 2] \times [-1, 1]^{m-1}$.

Given $b > 0$, a C^k ($k \geq 3$) family of diffeomorphism φ_a is a nonsingular b -perturbation of $\hat{f}_a$, if $\|\varphi(a, x, Y) - \hat{f}_a(a, x, Y)\| \leq b$, where $\varphi(a, x, Y) = \varphi_a(x, Y)$, $\hat{f}_a(a, x, Y) = \hat{f}_a(x, Y) = (1 - ax^2, 0^{m-1})$ and $\|\cdot\|$ is the C^k norm.

It follows from the main Theorem in [1], Theorem B in [4] and the proof of Theorem A in [6] that if b is small, then there exists a positive measure set Δ in the parameter space near 2, the corresponding diffeomorphism φ_a possesses a codimension-1 Hénon-like strange attractor Λ . Moreover $\Lambda = \overline{w^u(p_a)}$, where p_a is the primary fixed point of φ_a , which is the continuation of the fixed point $(\frac{1}{2}, 0^{m-1})$ of $\hat{f}_2$.

Now we recall some notations, concepts and definitions which will be used throughout the paper.

Fix $a \in \Delta$, and denote $\varphi = \varphi_a$, $p = p_a$. Let W denote the global unstable manifold of p .

2.1. The critical set. We use the following notation. For $A \in GL(m, R)$, if $v \mapsto \frac{|Av|}{|v|}$ is not constant, let $f(A)$ denote the unit vector that is maximally expanding by A , and $e(A) = \{f(A)\}^\perp$ denote the contractive hyperplane of A . Write $e_n(z) = e(D\varphi_z^n)$ if it makes sense. For $z \in W$, let $\tau \in T_z R^m$ denote a unit vector, tangent to W .

A certain subset $\mathcal{C}$ of W called the critical set is singled out in sects.5 and 6 of [1] and section 5 of [4] [6] to play the role of 0 in 1-dimension. Each $z_0 \in \mathcal{C}$ has the property that tangent direction at $\varphi(z_0)$ is (exponentially) contracted by all forward iterates of $D\varphi$. This can be thought of as the moral equivalent to $f'_a(0) = 0$.

2.2. Controlled orbits. For $(v, V) \in R \times R^{m-1}$ we define $|slope|(v, V) = \|V/v\|$.

Definition. A curve in R^m is called a $C^2(b)$ curve if $|slope|(\gamma') \leq 10b$ and its curvature $\kappa \leq 10b$.

- (i) Let $z_0 = (x_0, Y_0)$ be a critical point lying on a piece of $C^2(b)$ curve $\gamma \subset W$ and $\zeta_0 = (\xi_0, \mathcal{Y}_0)$ be an arbitrary point in $(-\delta, \delta) \times R^{m-1}$. We say that ζ_0 is in tangential position with respect to z_0 if there is $\widehat{z}_0 = (\widehat{x}_0, \widehat{Y}_0) \in \gamma$ with $\widehat{x}_0 = \xi_0$ and $|\widehat{Y}_0 - \mathcal{Y}_0| < |\widehat{x}_0 - x_0|^4$.
- (ii) Let $\zeta_0 \in W$, and let $\zeta_n = \varphi^n \zeta_0$. We say that the orbit of ζ_0 is controlled on the time interval $[0, N)$, $N \leq \infty$, if the following holds. If $\zeta_0 \notin \mathcal{C}$, let $n_1 < n_2 < \dots$ be the times when ζ_n is in $(-\delta, \delta) \times R^{m-1}$. If $\zeta_0 \in \mathcal{C}$, take $n_1 > 0$ to be the first time ζ_n return to $(-\delta, \delta) \times R^{m-1}$. Then for every $n_i \in [0, N)$, ζ_{n_i} is attached to a critical point with respect to which it is in tangential position. This critical point is denoted by $z(\zeta_{n_i})$ and is called the binding point of ζ_{n_i} .

2.3. Bound periods. Let $z_0 \in \mathcal{C}$, and let ζ_0 be in tangential position with respect to z_0 . As a first approximation we define the bound period $\tilde{p}(\zeta_0, z_0)$ to be the largest k such that

$$|\varphi^j \zeta_0 - \varphi^j z_0| < e^{-\beta j} \quad j < k.$$

(Recalled that $\alpha = 10^{-6}$, $\beta = 14\alpha$.)

Now consider ζ_0 whose orbit is controlled on $[0, \infty)$, and let $\tilde{p}_i = \tilde{p}(\zeta_{n_i}, z(\zeta_{n_i}))$. Note that it is entirely possible for ζ_n with $n_i < n < n_i + \tilde{p}_i$ to return to $(-\delta, \delta) \times R^{m-1}$, so that bound periods can be initiated in the middle of bound periods. It is shown in section 6.2 of [1], section 8 of [4] and section 6 of [6] that by modifying slightly the definition of $\tilde{p}$, bound periods can be mode nested. More precisely, one can choose $\{p_i\}_{i=1}^\infty$ in such a way that for all i ,

- (i) $p_i \leq \tilde{p}_i$ and $|\varphi^{p_i} \zeta_{n_i} - \varphi^{p_i} z(\zeta_{n_i})| \geq e^{-\beta^* p_i}$ for some $\beta^* \approx \beta$,
- (ii) if n_j is such that $n_i < n_j < n_i + p_i$, then $n_j + p_j \leq n_i + p_i$.

The bound period between ζ_{n_i} and $z(\zeta_{n_i})$ is then defined to be p_i . It is further required that if the bound relation between z_{n_i} and $z(z_{n_i})$ is still in effect at time $n_j > n_i$, then we must have $z(z_{n_j}) = z(\varphi^{n_j - n_i} z(z_{n_i}))$.

2.4. Bound of free states. Conceptually, it is convenient to divide a controlled orbit into free and bound states. Let ζ_0 be as above.

- For $n \leq n_1$, ζ_n is free.
- For $n_1 < n < n_1 + p_1$, ζ_n is said to be in bound state.
- Let n_i be the first return to $(-\delta, \delta) \times R^{m-1}$ after $n_1 + p_1$. Then ζ_n is free for $n_1 + p_1 \leq n \leq n_i$, and ζ_{n_i} is called a free return.
- ζ_n is in bound state again during the period $n_i < n < n_i + p_i$, and so on.

Let $\mathcal{P} = \{I_{\mu_j}\}$ be the partition of $(-\delta, \delta)$, given by $(-\delta, \delta) = \bigcup_{|\mu| \geq \mu_0} I_\mu$, where $I_\mu = (e^{-(\mu+1)}, e^{-\mu})$ for $\mu > 0$, $I_\mu = -I_{-\mu}$, for $\mu < 0$.

Lemma 2.1. *Let $\xi_0 \in W$ and assume that $\xi_i \notin \mathcal{C}$ for all i , then ξ_i is in a free state infinitely often, and the following holds for $\tau_i = D\varphi_{\xi_0}^i \tau$, where τ denotes a unit vector, tangent to W at point ξ_0 .*

(I) *Expansion outside of $(-\delta, \delta) \times R^{m-1}$.*

There are $c_0 > 0$ and $M_0 \in \mathbb{Z}^+$ such that

(1) *If ξ_i is free and $\xi_i, \dots, \xi_{i+M_0} \notin (-\delta, \delta) \times R^{m-1}$, then*

$$\frac{|\tau_{i+M_0}|}{|\tau_i|} \geq e^{c_0 M_0};$$

(2) *If $\xi_i \notin (-\delta, \delta) \times R^{m-1}$ is free, and $k > i$ is the first time return, namely $\xi_k \in (-\delta, \delta) \times R^{m-1}$, then*

$$\frac{|\tau_k|}{|\tau_i|} \geq e^{c_0(k-i)}.$$

(II) *Estimate of bound period.*

There is $c_1 \approx \log 2$, such that following holds: If $\xi_i \in (-\delta, \delta) \times R^{m-1}$ is free and becomes bound at this time $z(\xi_i) \in \mathcal{C}$ with bound period p , then

(a) *If $e^{-\mu-1} \leq |\xi_i - z(\xi_i)| \leq e^{-\mu}$, then $\frac{1}{2}\mu \leq p \leq 5\mu$;*

(b) *$\frac{|\tau_{i+j}|}{|\tau_i|} \geq 3|\xi_i - z(\xi_i)| \cdot \left| D\varphi_{z(\xi_i)}^j \begin{pmatrix} 0 \\ 1^{m-1} \end{pmatrix} \right| \geq |\xi_i - z(\xi_i)| e^{c_1 j}$, $0 \leq j \leq p$;*

(c) *$\frac{|\tau_{i+p}|}{|\tau_i|} \geq e^{\frac{1}{3}c_1 p}$.*

Proof. See Corollary 1 in section 2.1.1 of [2]. □

For x_0 with $|x_0| \ll \delta$, let $\mathcal{P}_{[x_0]}$ denote a copy of $\mathcal{P}$ with 0 moved to x_0 , more precisely, let $h : (-\delta, \delta) \rightarrow (-\delta, \delta)$ be a piecewise linear homeomorphism taking the points $-\delta, -e^{-(\mu_0+1)}, x_0, e^{-(\mu_0+1)}, \delta$, to $-\delta, -e^{-(\mu_0+1)}, 0, e^{-(\mu_0+1)}, \delta$ respectively and let $\mathcal{P}_{[x_0]} = h^{-1}\mathcal{P}$.

Furthermore, if γ is a horizontal curve and $z_0 = (x_0, Y_0)$ is such that $|x_0| \ll \delta$, then $\mathcal{P}_{[z_0]} = \mathcal{P}_{[x_0]}$ is a partition on $\gamma \cap ((-\delta, \delta) \times R^{m-1})$. Once γ, z_0 are fixed, for the sake of simplicity, we will consider I_μ as though it is a subsegment of γ . Let $J_\mu^+ = I_{\mu-1} \cup I_\mu \cup I_{\mu+1}$, in particular, $J_{\mu_0}^+ = I_{\mu_0-1} \cup I_{\mu_0} \cup I_{\mu_0+1}$.

Remark 1. For the point in J_μ^+ , we have defined bound period as in section 2.3, for a segment $I \subset J_\mu^+$, we define its bound period $p = \max_{x \in J_\mu^+} p(x)$, from Lemma 2.1, it follows that $\frac{|\tau_p|}{|\tau|} \geq e^{\frac{1}{6}c_1 p}$.

If γ_0 is a curve segment, let $\gamma_j = \varphi^j \gamma_0$, when we say that a segment $\gamma \subset W$ is free, it will assumed implicitly that $\forall z \in \gamma, z$ is controlled during the time interval $(-\infty, 0)$.

Define the projection map $\pi : R \times R^{m-1} \rightarrow R$ by $\pi(x, Y) = x$.

Lemma 2.2. *Let γ be a free segment of W , then*

(1) $\forall z \in \gamma, |\text{slope}|(\tau(z)) < \frac{2b}{\delta}$;

(2) $\forall z \in \gamma \cap (-\delta, \delta) \times R^{m-1}, |\text{slope}|(\tau(z)) < 10b$.

Proof. See Lemma 1 in section 2.2.1 of [2]. □

3. THE MAIN THEOREM

In this section, we will give the proof of the Theorem.

Lemma 3.1. *For one-dimensional map $f_2(x) = 1 - 2x^2$, $x \in [-1, 1]$, $\{f_2^{-n}\hat{x}_2, n \geq 0\}$ is dense on $[-1, 1]$, where $\hat{x}_2$ is the fixed point of f_2 .*

Proof. The map f_2 has an absolutely continuous invariant measure with positive entropy. Therefore f_2 is exact on $[-1, 1]$, that is to say, given an interval $I \subset [-1, 1]$, there exists $N(I)$ such that $f_2^{N(I)}I = [-1, 1]$, from which it follows that $\{f_2^{-n}\hat{x}_2, n \geq 0\}$ is dense on $[-1, 1]$. $\square$

Lemma 3.2. *There exist a_0, a_1 , sufficiently close to 2 with $a_0 < a_1 < 2$, such that for the map f_a , $a \in [a_0, a_1]$, $\exists$ a neighbourhood V of $\hat{x}_2$ and intervals $B^\pm \subset I_{\pm\mu_0}$, $A^\pm \subset I_{\pm(\mu_0-1)}$ and two integers $N, M \in \mathbb{Z}^+$ such that $f_a^N B^\pm \supset V$, $f_a^M A^\pm \supset V$ and $\hat{x}_a \in \text{Int}V$, where $\hat{x}_a$ is a fixed point of f_a corresponding to $\hat{x}_2$.*

Proof. From Lemma 3.1, it follows that there exist intervals $B^\pm$ and $A^\pm$, a neighbourhood V of $\hat{x}_2$ and $N, M \in \mathbb{Z}^+$ such that $f_2^N B^\pm \supset V$ and $f_2^M A^\pm \supset V$. By openness, this implies there are a_0, a_1 sufficiently near 2, such that $\hat{x}_a \in V$ and $f_a^N B^\pm \supset V$, $f_a^M A^\pm \supset V$ for $a \in [a_0, a_1]$. $\square$

Lemma 3.3. *If $a \in [a_0, a_1]$, then both $\hat{f}_a^N(B^\pm \times \{0^{m-1}\})$ and $\hat{f}_a^M(A^\pm \times \{0^{m-1}\})$ intersect $w^s(\hat{x}_a, 0^{m-1})$ transversally.*

Proof. For $\hat{f}_a, a \in [a_0, a_1]$, the stable set of $(\hat{x}_a, 0^{m-1})$ contains a $(m-1)$ -dimension hyperplane through the point $(\hat{x}_a, 0^{m-1})$, which is perpendicular to the x -axis. From Lemma 3.2, it follows that $\hat{f}_a^N(B^\pm \times \{0^{m-1}\})$ and $\hat{f}_a^M(A^\pm \times \{0^{m-1}\})$ intersects $w^s(\hat{x}_a, 0^{m-1})$ transversally. $\square$

Lemma 3.4. *There exists b_0 small enough such that if $b < b_0$ then both $\varphi_a^N(\tilde{B}^\pm)$ and $\varphi_a^M(\tilde{A}^\pm)$ intersect $w^s(p_a)$, for $a \in [a_0, a_1]$, where $\tilde{B}^\pm \subset B^\pm \times [-1, 1]^{m-1}$, $\tilde{A}^\pm \subset A^\pm \times [-1, 1]^{m-1}$, and $\pi(\tilde{B}^\pm) = B^\pm \times \{0^{m-1}\}$, $\pi(\tilde{A}^\pm) = A^\pm \times \{0^{m-1}\}$, and $\tilde{B}^\pm$ (respectively $\tilde{A}^\pm$) are close to $B^\pm \times \{0^{m-1}\}$ (respectively $A^\pm \times \{0^{m-1}\}$) in the sense of the Hausdorff metric.*

Proof. According to the differentiable dependence of stable manifolds in a compact set and Lemma 3.3, it follows that there exists b_0 small enough that provided $b < b_0$, then for $a \in [a_0, a_1]$, both $\varphi_a^N(B^\pm \times \{0^{m-1}\})$ and $\varphi_a^M(A^\pm \times \{0^{m-1}\})$ intersect $w^s(p_a)$ transversally. Therefore if $\tilde{B}^\pm \subset B^\pm \times [-1, 1]^{m-1}$ and $\pi(\tilde{B}^\pm) = B^\pm \times \{0^{m-1}\}$, $\tilde{B}^\pm$ is close to $B^\pm \times \{0^{m-1}\}$ in the sense of the Hausdorff metric. Then because of the differentiable dependence on the map, $\varphi_a^N(\tilde{B}^\pm)$ intersects $w^s(p_a)$ transversally. Analogously, if $\tilde{A}^\pm \subset A^\pm \times [-1, 1]^{m-1}$ and $\pi(\tilde{A}^\pm) = A^\pm \times \{0^{m-1}\}$, $\tilde{A}^\pm$ is close to $A^\pm \times \{0^{m-1}\}$ in the sense of the Hausdorff metric, then $\varphi_a^M(\tilde{A}^\pm)$ intersects $w^s(p_a)$ transversally. $\square$

Theorem. *There exists $b_0 > 0$ small enough such that if $b < b_0$ then for $a \in \Delta \cap [a_0, a_1]$ the transversal homoclinic points are dense in Λ .*

Proof. Since W is dense in Λ , it is sufficient to prove that the transversal homoclinic points are dense in H_1 , where H_1 denotes the primary segment through the fixed point p in W .

Given $I \subset H_1$, let z_0 be the critical point in H_1 , $\mathcal{P}_{[z_0]}$ denotes the partition of $H_1 \cap (-\delta, \delta)$ relative to z_0 , and $I_{\mu_0}, I_{-\mu_0}$ are outmost interval of $\mathcal{P}_{[z_0]}$. Without loss of generality, we assume $I \not\supset I_{\pm\mu_0}, I \not\supset I_{\pm(\mu_0-1)}$. Otherwise, by Lemma 3.3 and 3.4, we have that I contains the transversal homoclinic points.

Now we consider two cases.

(I) $I \cap (-\delta, \delta) \times R^{m-1} = \emptyset$.

First, we prove that there exist n_1 and $\mu_1 > 0$ such that $\varphi^{n_1}I \supset I_{\mu_1}$ or $\varphi^{n_1}I \supset I_{-\mu_1}$. If $\mu_1 = \mu_0$ or $\mu_1 = \mu_0 - 1$, then by Lemma 3.3 and 3.4, we have that I contains the transversal homoclinic point.

From Lemma 2.1 and 2.2, it follows that there exists n such that $\varphi^n I$ intersects $(-\delta, \delta) \times R^{m-1}$ and $\varphi^n I \cap (-\delta, \delta) \times R^{m-1}$ bound to some point $z_1 \in \mathcal{C}$, moreover $\varphi^n I$ is almost horizontal.

If $\varphi^n I \cap (-\delta, \delta) \times R^{m-1}$ contains some I_μ , then let $n_1 = n$ and μ_1 denote the minimal absolute value of μ .

Otherwise we have two subcases:

- (1) $\varphi^n I \cap (-\delta, \delta) \times R^{m-1} \subset I_{\mu_0} \times R^{m-1}$, moreover $\varphi^n I \cap [(-\delta, \delta) \times R^{m-1}]^c \neq \emptyset$.

If case (1) occurs, moreover the segment $\varphi^n I \cap [(-\delta, \delta) \times R^{m-1}]^c$ is contained in $I_{(\mu_0-1)}$ or $I_{-(\mu_0-1)}$. Then keep iterating $\varphi^n I$. From Lemmas 2.1 and 2.2, there must be some n' such that $\varphi^{n'+n} I$ intersects $(-\delta, \delta) \times R^{m-1}$ again. Moreover $|\varphi^{n'+n} I| > 2|\varphi^n I|$ and $\varphi^{n'+n} I$ is almost horizontal.

If case (1) occurs, but the segment $\varphi^n I \cap [(-\delta, \delta) \times R^{m-1}]^c$ contains $I_{(\mu_0-1)}$ or $I_{-(\mu_0-1)}$, then let $n_1 = n$ and $\mu_1 = \mu_0 - 1$.

- (2) $\varphi^n I \subset J_\mu^+$.

If this case occurs, then keep iterating $\varphi^n I$. From Lemma 2.1 (II) and Remark 1, it follows that $|\varphi^{n+p} I| \geq e^{\frac{1}{6}c_1 p} |\varphi^n I| > 2|\varphi^n I|$. Keep iterating $\varphi^{n+p} I$ until there is some n'' such that $\varphi^{n+p+n''} I \cap (-\delta, \delta) \times R^{m-1}$ is nonempty. By Lemma 2.1, we have $|\varphi^{n+p+n''} I| > |\varphi^{n+p} I| > 2|\varphi^n I|$, and by Lemma 2.2, we have $\varphi^{n+p+n''} I$ is almost horizontal.

By repeating the above arguments, we conclude that there must be n_1 such that $\varphi^{n_1} I$ contain some I_μ and $\varphi^{n_1} I$ is almost horizontal, denote the minimal absolute value of μ by μ_1 .

Next, we iterate I_{μ_1} or $I_{-\mu_1}$ (in term of $\varphi^{n_1} I \supset I_{\mu_1}$ or $I_{-\mu_1}$). By Lemma 2.1 (II), it follows that

$$|\varphi^{p(\pm\mu_1)} I_{\pm\mu_1}| \geq e^{\frac{1}{3}c_1 p(\pm\mu_1)} |I_{\pm\mu_1}|.$$

There exists $l(\pm\mu_1) \geq 0$ such that $\varphi^{l(\pm\mu_1)+p(\pm\mu_1)} I_{\pm\mu_1}$ intersects $(-\delta, \delta) \times R^{m-1}$. If $\varphi^{l(\pm\mu_1)+p(\pm\mu_1)} I_{\pm\mu_1}$ contains I_μ of some $\mathcal{P}_{[z]}$, then denote the minimal absolute value of μ by μ_2 . Otherwise repeat the arguments as above, keep iterating; there must be n_2 such that $\varphi^{n_2} I_{\pm\mu_1}$ contains some I_μ and $\varphi^{n_2} I_{\pm\mu_1}$ is almost horizontal, then denote the minimal absolute value of μ by μ_2 .

From Lemma 2.1, it follows that $|\varphi^{n_2} I_{\mu_1}| \geq e^{\frac{1}{3}c_1 p} |I_{\mu_1}|$, since $p \geq \frac{1}{2}\mu_1$, it has $|\varphi^{n_2} I_{\mu_1}| \geq 2e^{-(\mu_1-1)}$. Thus we get that $0 < \mu_2 < \mu_1$.

We then consider I_{μ_2} or $I_{-\mu_2}$ as above, repeat the arguments as above, and we have $0 < \mu_k < \mu_{k-1} < \dots < \mu_2 < \mu_1$. Therefore there must be $\tilde{N}$, such that $\varphi^{\tilde{N}} I$ contains an almost horizontal segment J which contains some I_i , for $i = \pm\mu_0, \pm(\mu_0 - 1)$.

From Lemma 3.4, for b_0 small enough, if $b < b_0$, then $\varphi^N \varphi^{\tilde{N}} I$ or $\varphi^M \varphi^{\tilde{N}} I$ intersects $w^s(p)$ transversally. Hence I contains the transversal homoclinic points.

- (II) $I \cap (-\delta, \delta) \times R^{m-1} \neq \emptyset$.

If I contains some I_μ , let μ_1 denote the minimal absolute value of μ . Keep iterating, analogous to (I). If not, then I must be contained in some J_μ^+ for $\mu \geq \mu_0$, keep iterating as above. We also have that I contains the transversal homoclinic points.

Summarizing (I) and (II), it follows that the transversal homoclinic points are dense in H_1 . Therefore the transversal homoclinic points are dense in W . $\square$

Corollary. *The hyperbolic periodic points are dense in the codimension-1 Hénon-like strange attractor. Moreover the hyperbolic periodic points that are heteroclinically related to the primary fixed point are also dense in it.*

Proof. From the proof of the Theorem, it follows the transversal homoclinic points of $w^s(p)$ are dense in Λ . By the Smale-Birkhoff Theorem, we have that the hyperbolic points that are heteroclinically related to the primary fixed point p are dense and the hyperbolic periodic points are dense in Λ . $\square$

ACKNOWLEDGMENTS

Part of the work was done while the author visited IMPA. The author thanks TWAS, IMPA for their support and hospitality, Professor M. Viana and Zhang Zhifen for their guidance and encouragement, Professor L. S. Young and J. Kennedy for their discussions, and the referees for their suggestions.

REFERENCES

1. M.Benedicks & L.Carleson, *The dynamics of the Hénon map*, Ann. Math. **133** (1991), 73-169. MR **92d**:58116
2. M.Benedicks & L.S.Young, *Sinai-Bowen-Ruelle measures for certain Hénon maps*, Inventiones Math. **112** (1993), 541-576. MR **94e**:58074
3. M.Hénon, *A two-dimensional mapping with a strange attractor*, Commun. Math. Phys **50** (1976), 69-77. MR **54**:10917
4. L.Mora & M.Viana, *Abundance of strange attractors*, Acta .Math. **171** (1993), 1-71. MR **94k**:58089
5. J.Palis & F.Takens, *Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations*, Cambridge Press, 1993. MR **94h**:58129
6. M.Viana, *Strange attractors in higher dimensions*, Bull.Braz.Math.Soc. **24** (1993), 13-62. MR **94k**:58093

DEPARTMENT OF MATHEMATICS, SUZHOU UNIVERSITY, SUZHOU 215006, JIANGSU, PEOPLE'S REPUBLIC OF CHINA

E-mail address: ylcao@nsad.suda.edu.cn