A NOTE ON HOLOMORPHIC MAPS WITH UNIPOTENT JACOBIAN MATRICES

YU QING CHEN

(Communicated by Steven R. Bell)

ABSTRACT. We prove that a holomorphic map $H : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is invertible if its Jacobian matrix JH is unipotent.

1. INTRODUCTION

Let $\mathbb{C}$ be the complex number field. Given a polynomial map $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ with $F = (f_1, f_2, \dots, f_n)$, where $f_i \in \mathbb{C}[z_1, z_2, \dots, z_n]$, a simple algebraic argument tells us that

$$J_F = \det\left[\frac{\partial f_i}{\partial z_j}\right] \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$$

whenever F is invertible. The Jacobian Conjecture asserts that the converse is true also.

The Jacobian Conjecture is false for holomorphic maps. An easy example is

$$\begin{aligned} F : \mathbb{C}^2 &\rightarrow \mathbb{C}^2, \\ f_1 &= e^{z_1}, \\ f_2 &= z_2 e^{-z_1}. \end{aligned}$$

In [BCW] the Jacobian Conjecture has been reduced to the Unipotent Jacobian Conjecture, which states

The Unipotent Jacobian Conjecture. *If the Jacobian matrix JF of F is a unipotent matrix, then F is invertible.*

We suspect that this conjecture could also be true for holomorphic maps. In this note, we give a proof of the Unipotent Jacobian Conjecture for holomorphic maps with $n = 2$.

2. MAIN RESULTS

In this section, we first prove a theorem concerning holomorphic maps $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ with $JF^2 = 0$. The idea of the proof is obtained from [CSW]. We then apply the theorem to the case $n = 2$ and $F(z) = H(z) - z$, where $H(z)$ is an arbitrary holomorphic map $H : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ with unipotent Jacobian matrix. This yields that H is invertible, i.e. Corollary 2.3.

Received by the editors September 26, 1997.

1991 *Mathematics Subject Classification.* Primary 32H99.

Theorem 2.1. *Let $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a holomorphic map. The following statements are equivalent:*

- (1) $JF(z)^2 = 0$ for all $z \in \mathbb{C}^n$,
- (2) $F(z + JF(z)z') = F(z)$ for all $z, z' \in \mathbb{C}^n$, and
- (3) $JF(z + JF(z)z')JF(z) = 0$ for all $z, z' \in \mathbb{C}^n$.

Proof. (1) $\Rightarrow$ (2). Using Taylor expansion

$$f(z + y) = f(z) + \sum_{k=1}^{\infty} \frac{1}{k!} \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \prod_{s=1}^k y_{i_s},$$

one has

$$\begin{aligned} f_i(z + JF(z)z') &= \\ f_i(z) + \sum_{k=1}^{\infty} \frac{1}{k!} \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \prod_{s=1}^k \sum_{j_s=1}^n \frac{\partial f_{i_s}(z)}{\partial z_{j_s}} z'_{j_s} &= \\ f_i(z) + \sum_{k=1}^{\infty} \frac{1}{k!} \sum_{j_1, j_2, \dots, j_k=1}^n \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \prod_{s=1}^k \frac{\partial f_{i_s}(z)}{\partial z_{j_s}} \prod_{s=1}^k z'_{j_s}. \end{aligned}$$

Define

$$D_{j_1, j_2, \dots, j_k}^{[i]} = \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \prod_{s=1}^k \frac{\partial f_{i_s}}{\partial z_{j_s}}.$$

We now show that $D_{j_1, j_2, \dots, j_k}^{[i]} = 0$ for all $k \geq 1$ and all $1 \leq j_1, j_2, \dots, j_k, i \leq n$ by induction on k . When $k = 1$, the $D_{j_1}^{[i]}$'s are the entries of $JF(z)^2$, and therefore are equal to 0. Suppose $D_{j_1, j_2, \dots, j_k}^{[i]} = 0$ for all $1 \leq j_1, j_2, \dots, j_k, i \leq n$. Then

$$\begin{aligned} D_{j_1, j_2, \dots, j_{k+1}}^{[i]} &= \sum_{i_1, i_2, \dots, i_{k+1}=1}^n \frac{\partial^{k+1} f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_{k+1}}} \prod_{s=1}^{k+1} \frac{\partial f_{i_s}(z)}{\partial z_{j_s}} \\ &= \sum_{i_{k+1}=1}^n \frac{\partial D_{j_1, j_2, \dots, j_k}^{[i_{k+1}]}}{\partial z_{i_{k+1}}} \frac{\partial f_{i_{k+1}}}{\partial z_{j_{k+1}}} \\ &\quad - \sum_{i_{k+1}=1}^n \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \sum_{s=1}^k \frac{\partial^2 f_{i_s}(z)}{\partial z_{j_s} \partial z_{i_{k+1}}} \frac{\partial f_{i_{k+1}}(z)}{\partial z_{j_{k+1}}} \prod_{t=1, t \neq s}^k \frac{\partial f_{i_t}(z)}{\partial z_{j_t}}. \end{aligned}$$

Since

$$\sum_{i_{k+1}=1}^n \frac{\partial f_{i_s}(z)}{\partial z_{i_{k+1}}} \frac{\partial f_{i_{k+1}}(z)}{\partial z_{j_{k+1}}} = 0,$$

we have

$$\sum_{i_{k+1}=1}^n \frac{\partial^2 f_{i_s}(z)}{\partial z_{j_s} \partial z_{i_{k+1}}} \frac{\partial f_{i_{k+1}}(z)}{\partial z_{j_{k+1}}} = - \sum_{i_{k+1}=1}^n \frac{\partial f_{i_s}(z)}{\partial z_{i_{k+1}}} \frac{\partial^2 f_{i_{k+1}}(z)}{\partial z_{j_s} \partial z_{j_{k+1}}}.$$

Thus

$$\begin{aligned}
 & D_{j_1, j_2, \dots, j_{k+1}}^{[i]} \\
 &= \sum_{s=1}^k \sum_{i_1, i_2, \dots, i_k=1}^n \frac{\partial^k f_i(z)}{\partial z_{i_1} \partial z_{i_2} \cdots \partial z_{i_k}} \sum_{i_{k+1}=1}^n \frac{\partial f_{i_s}(z)}{\partial z_{i_{k+1}}} \frac{\partial^2 f_{i_{k+1}}(z)}{\partial z_{j_s} \partial z_{j_{k+1}}} \prod_{t=1, t \neq s}^k \frac{\partial f_{i_t}(z)}{\partial z_{j_t}} \\
 &= \sum_{s=1}^k \sum_{i_{k+1}=1}^n D_{j_1, j_2, \dots, j_{s-1}, i_{k+1}, j_{s+1}, \dots, j_{k-1}, j_k}^{[i]} \frac{\partial^2 f_{i_{k+1}}}{\partial z_{j_s} \partial z_{j_{k+1}}} \\
 &= 0.
 \end{aligned}$$

(2) $\Rightarrow$ (3). We fix z and consider z' as variable and compute the Jacobian matrix with respect to z' in the left-hand side of (2). (3) $\Rightarrow$ (1). Set $z' = 0$ in (3). $\square$

Theorem 2.2. *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a holomorphic map. Then $JF(z)^2 = 0$ if and only if there exist an entire function $f : \mathbb{C} \rightarrow \mathbb{C}$ and constants a, b, c_1 , and c_2 in $\mathbb{C}$, such that $F = (bf(az_1 + bz_2) + c_1, -af(az_1 + bz_2) + c_2)$.*

Proof. If $JF(z) = 0$ for all $z \in \mathbb{C}^2$, then F is a constant map and the assertion is trivial. If $JF(z) \neq 0$, we define

$$\Omega = \{z \in \mathbb{C}^2 \mid JF(z) \neq 0\}.$$

Note that Ω is an open subset of $\mathbb{C}^2$. For every $z \in \Omega$, define

$$L_z = \{z + JF(z)z' \mid z' \in \mathbb{C}^2\}.$$

Since $JF(z) \neq 0$ and $JF(z)^2 = 0$, L_z is a complex line in $\mathbb{C}^2$ passing through z . From Theorem 2.1(2), $F|_{L_z}$ is a constant map. We claim that all these lines are parallel to each other. Suppose to the contrary. Then there are two lines L_{z_1} and L_{z_2} , $z_1, z_2 \in \Omega$, meeting at a point $z \in \mathbb{C}^2$. Thus, F is constant on $L_{z_1} \cup L_{z_2}$. Also for every $z \in \Omega$, L_z meets at least one of L_{z_1} and L_{z_2} . Therefore, F is a constant map on

$$\Omega \subset \bigcup_{z \in \Omega} L_z.$$

Since Ω is open, F is a constant map on $\mathbb{C}^2$, which contradicts $JF(z) \neq 0$. Hence there exist two complex numbers a and b , $|a| + |b| \neq 0$, such that

$$(a, b)JF(z)z' = 0$$

for all $z, z' \in \mathbb{C}^2$. Therefore

$$(a, b)JF(z) = 0$$

for all $z \in \mathbb{C}^2$. Let

$$\begin{aligned}
 G : \mathbb{C}^2 &\rightarrow \mathbb{C}^2, \\
 g_1 &= a'z_1 + b'z_2, \\
 g_2 &= az_1 + bz_2,
 \end{aligned}$$

be a linear map such that G is invertible. Set $H = GFG^{-1}$. The Jacobian matrix

$$JH(z) = JG(z)JF(G^{-1}(z))JG(z)^{-1}$$

satisfies $JH(z)^2 = 0$ and its second row equals 0. This implies

$$\begin{aligned} H : \mathbb{C}^2 &\rightarrow \mathbb{C}^2, \\ h_1 &= h(z_2), \\ h_2 &= \gamma, \end{aligned}$$

where h is an entire function and γ is a constant. If we denote $\det JG(z)^{-1}$ by d , dh by f , $-db'\gamma$ by c_1 and $da'\gamma$ by c_2 , then

$$\begin{aligned} F : \mathbb{C}^2 &\rightarrow \mathbb{C}^2, \\ f_1 &= bf(az_1 + bz_2) + c_1, \\ f_2 &= -af(az_1 + bz_2) + c_2. \end{aligned}$$

The converse is obvious and the proof is completed. $\square$

Corollary 2.3. *If the Jacobian matrix JH of a holomorphic map $H : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is a unipotent matrix, then H is invertible.*

Proof. Let $H(z) = z + F(z)$, where F is a holomorphic map with nilpotent Jacobian matrix. By Theorem 2.2, it is easy to check that $G(z) = z - F(z - F(z))$ is the inverse of H . $\square$

Remark 1. If F is a polynomial map, Theorem 2.2 can be derived from a result of [BCW].

Remark 2. Given a holomorphic function $f : \mathbb{C}^2 \rightarrow \mathbb{C}$, there is a holomorphic function $g : \mathbb{C}^2 \rightarrow \mathbb{C}$ such that the holomorphic map $F = (f, g)$ or $(g, f) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ has nilpotent Jacobian matrix JF if and only if f satisfies the partial differential equation

$$(*) \quad \left(\frac{\partial f}{\partial z_1}\right)^2 \frac{\partial^2 f}{\partial z_2^2} + \left(\frac{\partial f}{\partial z_2}\right)^2 \frac{\partial^2 f}{\partial z_1^2} = 2 \frac{\partial f}{\partial z_1} \frac{\partial f}{\partial z_2} \frac{\partial^2 f}{\partial z_1 \partial z_2}.$$

This follows since $\det(JF) = 0$ and $\text{Tr}(JF) = 0$ and we can then eliminate the function g by using the mixed second derivative of g . Thus Theorem 2.2 is equivalent to the following, which can also be proved in the same way as the combination of proofs of Theorem 2.1 and Theorem 2.2:

Theorem 2.4. *Let f be a holomorphic function on $\mathbb{C}^2$. Then f satisfies the differential equation (*) if and only if $f = h(az_1 + bz_2)$, where h is an analytic function on $\mathbb{C}$ and a and b are constants.*

ACKNOWLEDGMENT

The author would like to thank Professor Henry H. Glover for his guidance and encouragement.

REFERENCES

- [BCW] H. Bass, E. H. Connell and D. Wright, *The Jacobian Conjecture: Reduction of Degree and Formal Expansion of the Inverse*, Bull. AMS **7** (1982), 287–330. MR **83k**:14028
- [CSW] C. C. Cheng, T. Sakalis and S. S. Wang, *A Case of the Jacobian Conjecture*, J. Pure Appl. Algebra **96** (1994), 15–18. MR **95i**:14018a

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OHIO 43210
E-mail address: yuqchen@math.ohio-state.edu