ON THE RANGE AND THE KERNEL OF THE OPERATOR $X \mapsto AXB - X$

A. MAZOUZ

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let $L(H)$ denote the algebra of (bounded linear) operators on the separable complex Hilbert space H , and let $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ denote a norm ideal in $L(H)$. For $A, B \in L(H)$, let the derivation $\delta_{A,B}: L(H) \rightarrow L(H)$ be defined by $\delta_{A,B}(X) = AX - XB$, and let $\Delta_{A,B}: L(H) \rightarrow L(H)$ be defined by $\Delta_{A,B}(X) = AXB - X$. The main result of this paper is to show that if A, B are contractions, then for every operator $T \in \mathfrak{J}$ such that $ATB = T$, then $\|AXB - X + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}}$ for all $X \in \mathfrak{J}$.

1. INTRODUCTION

Recently Du Hong Ke ([2]) proved that if A, B are contractions, then for every operator S such that $ASB = S$, $A^*SB^* = S$, then

$$\|AXB - X + S\| \geq \|S\| \quad \text{for all operators } X \in L(H).$$

Duggal ([4]) proved that if A, B are contractions, then $S \in C_2$ and $ASB = S$ imply

$$\|AXB - X + S\|_2^2 = \|AXB - X\|_2^2 + \|S\|_2^2,$$

for all $X \in L(H)$, where C_2 denotes the (Hilbert) space of Hilbert-Schmidt operators on H . In this note, we shall prove the following theorem.

Theorem 1. *If A, B are contractions and $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a norm ideal in $L(H)$ and $T \in \mathfrak{J}$ is such that $ATB = T$, then*

$$\|AXB - X + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}} \quad \text{for all operators } X \in L(H).$$

2. SOME PRELIMINARIES

Definition 2.1 ([5]). A proper two-sided ideal $\mathfrak{J}$ in $L(H)$ is said to be a norm ideal if there is a norm on $\mathfrak{J}$ satisfying the following properties:

- i) $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a Banach space;
- ii) $\|AXB\|_{\mathfrak{J}} \leq \|A\| \|X\|_{\mathfrak{J}} \|B\|$ for all $A, B \in L(H)$ and for all $X \in \mathfrak{J}$;
- iii) $\|X\|_{\mathfrak{J}} = \|X\|$ for X a rank one operator.

Remark. If $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a norm ideal, then the norm $\|\cdot\|_{\mathfrak{J}}$ is unitarily invariant, in the sense that $\|UAV\|_{\mathfrak{J}} = \|A\|_{\mathfrak{J}}$ for all A in $\mathfrak{J}$ and unitary U, V in $L(H)$.

Received by the editors December 2, 1996 and, in revised form, October 16, 1997.
 1991 *Mathematics Subject Classification.* Primary 47B47, 47D50.

Example. Each proper ideal of $L(H)$ is contained in the ideal of compact operators. For any compact operator A , denote by $s_1(A) \geq s_2(A) \geq \dots$ the singular values of A , i.e., the eigenvalues of $(A^*A)^{\frac{1}{2}}$.

Two special families of unitarily invariant norms satisfying conditions i), ii), and iii) of Definition 2.1 are the Schatten p -norms defined as

$$\|A\|_p = \left(\sum_{j=1}^{\infty} s_j(A)^p \right)^{\frac{1}{p}}, \quad 1 \leq p \leq \infty,$$

where by convention

$$\|A\|_{\infty} = \max s_j(A) = s_1(A) = \|A\|,$$

and the Ky Fan norms defined as $\|A\|_k = \sum_{j=1}^k s_j(A)$, $k \geq 1$.

Theorem 2.2 ([3]). a) Let $A, B^* \in L(H)$ be contractions with C_0 completely non-unitary parts. If $\Delta_{A,B}(X) = 0$, then $\overline{\text{ran } X}$ reduces A , $\ker^{\perp} X$ reduces B^* , and $A|_{\overline{\text{ran } X}}$ and $B^*|_{\ker^{\perp} X}$ are unitarily equivalent operators.

b) Let A and B be contractions such that $\Delta_{A,B}(X) = 0$ and X a compact operator. Then the conclusions of part a) hold.

3. MAIN RESULTS

Theorem 3.1. If U is a unitary operator, $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a norm ideal in $L(H)$ and $T \in \mathfrak{J}$ is such that $TU = UT$, then

$$\|UX - XU + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}} \quad \text{for all operators } X \in \mathfrak{J}.$$

Proof. The proof is similar to J. Anderson's ([1, Theorem 1.4, p. 136]).

Corollary 3.2. If U, V are unitary operators, $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a norm ideal in $L(H)$ and $T \in \mathfrak{J}$ is such that $UT = TV$, then

$$\|\delta_{U,V}(X) + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}} \quad \text{for all operators } X \in \mathfrak{J}.$$

Proof. On $H \oplus H$, let

$$W = \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix}, \quad \tilde{T} = \begin{pmatrix} 0 & T \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \tilde{X} = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}.$$

Then $\tilde{T} \in \mathfrak{J} \oplus \mathfrak{J}$ and $W\tilde{X} - \tilde{X}W + \tilde{T} = \begin{pmatrix} 0 & UX - XV + T \\ 0 & 0 \end{pmatrix}$. Since $UT = TV$, it follows that $W\tilde{T} = \tilde{T}W$. By Theorem 3.1 we have $\|W\tilde{X} - \tilde{X}W + \tilde{T}\|_{\mathfrak{J} \oplus \mathfrak{J}} \geq \|\tilde{T}\|_{\mathfrak{J}}$ and so $\|UX - XV + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}}$.

Corollary 3.3. If U is a unitary operator, $(\mathfrak{J}; \|\cdot\|_{\mathfrak{J}})$ is a norm ideal in $L(H)$, and $T \in \mathfrak{J}$ is such that $UTU = T$. Then

$$\|UXU - X + T\|_{\mathfrak{J}} \geq \|T\|_{\mathfrak{J}} \quad \text{for all operators } X \in \mathfrak{J}.$$

Proof. Let T be such that $UTU = T$. Then $TU = U^*T$ and so

$$\begin{aligned} \|UXU - X + T\|_{\mathfrak{J}} &= \|U(XU - U^*X + U^*T)\|_{\mathfrak{J}} \\ &= \|XU - U^*X + U^*T\|_{\mathfrak{J}} \\ &\geq \|U^*T\|_{\mathfrak{J}} \quad (\text{Corollary 3.2}) \\ &\geq \|T\|_{\mathfrak{J}}. \end{aligned}$$

Corollary 3.4. *If U, V are unitary operators, $(\mathfrak{I}; \|\cdot\|_{\mathfrak{I}})$ is a norm ideal in $L(H)$, and $T \in \mathfrak{I}$ is such that $UTV = T$, then*

$$\|UXV - X + T\|_{\mathfrak{I}} \geq \|T\|_{\mathfrak{I}} \quad \text{for all operators } X \in \mathfrak{I}.$$

Proof. On $H \oplus H$, let

$$W = \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix}, \quad \tilde{T} = \begin{pmatrix} 0 & T \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \tilde{X} = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}.$$

Then $\tilde{T} \in \mathfrak{I} \oplus \mathfrak{I}$ and $W\tilde{X}W - \tilde{X} + \tilde{T} = \begin{pmatrix} 0 & UXV - X + T \\ 0 & 0 \end{pmatrix}$. Since $UTV = T$, it follows that $W\tilde{T}W = \tilde{T}$. By Corollary 3.3 we have $\|W\tilde{X}W - \tilde{X} + \tilde{T}\|_{\mathfrak{I} \oplus \mathfrak{I}} \geq \|\tilde{T}\|_{\mathfrak{I}}$ and so

$$\|UXV - X + T\|_{\mathfrak{I}} \geq \|T\|_{\mathfrak{I}}.$$

Proof of Theorem 1. Let T be an operator such that $\Delta(T) = 0$ and $T \in \mathfrak{I}$. By Theorem 2.2b) $\overline{\text{ran } X}$ reduces A , $\ker^{\perp} X$ reduces B^* and $A|_{\overline{\text{ran } X}}$ and $B^*|_{\ker^{\perp} X}$ are unitary operators. Put $H_1 = H = \overline{\text{ran } T} \oplus \overline{\text{ran } T}^{\perp}$, $H_2 = H = \ker(T)^{\perp} \oplus \ker(T)$ so that we get decompositions of operators respectively:

$$W = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix}, \quad B^* = \begin{pmatrix} B_1^* & 0 \\ 0 & B_2^* \end{pmatrix}.$$

For linear operators X, T from H_2 into H_1 we have:

$$X = \begin{pmatrix} X_1 & X_2 \\ X_3 & X_4 \end{pmatrix}, \quad T = \begin{pmatrix} T_1 & 0 \\ 0 & 0 \end{pmatrix}.$$

So

$$\begin{aligned} \|\Delta_{A,B}(X) + T\|_{\mathfrak{I}} &= \left\| \begin{pmatrix} T_1 + \Delta_{A_1,B_1}(X_1) & * \\ * & * \end{pmatrix} \right\|_{\mathfrak{I}} \\ &\geq \left\| \begin{pmatrix} T_1 + \Delta_{A_1,B_1}(X_1) & 0 \\ 0 & 0 \end{pmatrix} \right\|_{\mathfrak{I}} \\ &\geq \|T_1 + \Delta_{A_1,B_1}(X_1)\|_{\mathfrak{I}}. \end{aligned}$$

Since A_1, B_1 are unitary operators, then Corollary 3.4 implies that

$$\|\Delta_{A,B}(X) + T\|_{\mathfrak{I}} \geq \|T_1\|_{\mathfrak{I}} = \|T\|_{\mathfrak{I}}.$$

REFERENCES

1. J. Anderson, *On normal derivations*, Proc. Amer. Math. Soc. **38** (1973), 135–140. MR **47**:875
2. Du Hong Ke, *Another generalization of Anderson's theorem*, Proc. Amer. Math. Soc. **123** (1995), 2709–2714. MR **95k**:47032
3. B. P. Duggal, *On intertwining operators*, Monatsh. Math. **106** (1988), 139–148. MR **89k**:47031
4. B. P. Duggal, *A remark on normal derivations of Hilbert-Schmidt type*, Monatsh. Math. **112** (1991), 265–270. MR **92m**:47067
5. I. C. Gohberg and M. G. Krein, *Introduction to the theory of linear nonself-adjoint operators*, Transl. Math. Monographs, Vol. 18, Amer. Math. Soc., Providence, RI, 1969. MR **39**:7447

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ MONTPELLIER II, PLACE E.-BATAILLON, 34060 MONTPELLIER CEDEX, FRANCE