IDENTITIES OF INCOMPLETE KLOOSTERMAN SUMS

YE YANGBO

(Communicated by Dennis A. Hejhal)

ABSTRACT. Identities between incomplete Kloosterman sums and incomplete hyper-Kloosterman sums are established.

1. INTRODUCTION

In recent years various identities of Kloosterman sums and other exponential sums have been proved and their roles in relative trace formulas and Langlands' functoriality conjecture have been studied (Jacquet and Ye [3], Mao and Rallis [6], Ye [7], [8], [9], [10], and Zagier [11]). Some of these identities imply the fundamental lemmas of relative trace formulas and hence provide us new information about the corresponding functoriality of group representations.

More specifically, an identity is proved in Ye [10] between a complete hyper-Kloosterman sum and an exponential sum over a quadratic number field. The fundamental lemma of a relative trace formula for GL_n would follow from this identity and other conjectured identities. This relative trace formula would in turn characterize quadratic base change for GL_n .

In Ye [9] an identity is established between an exponential sum of Kloosterman type and an exponential sum over a cyclic algebraic number field of a prime degree. Although we do not know whether this identity is related to a relative trace formula, it provides us a new interpretation of Davenport-Hasse relations of Gauss sums (Davenport and Hasse [2]).

On the other hand Kloosterman sums and related exponential sums have important applications in number theory, based on estimation of individual sums and their weighted sums. We are especially interested in estimation of incomplete Kloosterman sums and incomplete hyper-Kloosterman sums. In Bump, Duke, Hoffstein, and Iwaniec [1] estimation of hyper-Kloosterman sums was used in improving bounds of Fourier coefficients of Maass cusp forms. Moreover, Luo, Rudnick, and Sarnak in [5] established a new lower bound of the first eigenvalue for Maass cusp forms based in part on estimates of hyper-Kloosterman sums. Recently Luo in [4] established a new bound for incomplete hyper-Kloosterman sums.

In this article we will establish identities between incomplete hyper-Kloosterman sums and incomplete classical Kloosterman sums in the case of prime power moduli. We will see that these incomplete sums are essentially the same and hence estimation of one sum implies estimation of the other sum.

Received by the editors November 18, 1997.

1991 *Mathematics Subject Classification*. Primary 11L05; Secondary 11F70.

The author was supported in part by NSF Grant #DMS 97-01225.

From the viewpoint of base change this identity is closely related to the identity in Ye [9]. Indeed, we may regard a local hyper-Kloosterman sum as the lifting of a local Kloosterman sum to the direct sum of n copies of local field $\mathbb{Q}_p$. Then the exponential sum in [9] is the lifting of an exponential sum of Kloosterman type to an unramified extension field of $\mathbb{Q}_p$ of degree n .

Our first result is an identity between incomplete exponential sums. We denote $e(x) = e^{2\pi i x}$ and set $\varepsilon_p = 1$ if $p \equiv 1 \pmod{4}$ and $\varepsilon_p = i$ if $p \equiv 3 \pmod{4}$. We will use Legendre symbols.

Theorem 1. *Let $c > 1$ be an odd integer such that $p^2 | c$ for any prime factor p of c . Let e be a nonzero integer and $m > 1$ an integer satisfying $m \equiv 1 \pmod{\phi(c)}$, where ϕ is the Euler function. For any z with $(z, c) = 1$ and any x with $1 \leq x \leq c$ we have*

$$\sum_{\substack{1 \leq b_1 \leq x, \\ b_2, \dots, b_m \pmod{c}, \\ (b_1, c) = \dots = (b_m, c) = 1}} e\left(\frac{1}{c}(b_1 + \dots + b_m + zb_1^e \bar{b}_2 \dots \bar{b}_m)\right) \\ = c^{(m-1)/2} \left(\prod_{\substack{p|c \text{ to an} \\ \text{odd power}}} \left(\frac{c_2 \dots c_r}{p}\right) \varepsilon_p^r \right) \sum_{\substack{1 \leq b \leq x, \\ (b, c) = 1}} e\left(\frac{1}{c}(b + zb^e \overline{(2-m)})\right)$$

where $y\bar{y} \equiv 1 \pmod{c}$ for any y with $(y, c) = 1$. Here for each prime p which divides c to an odd power the even constant r with $1 < r \leq m-3$ and the nonzero integers $c_2, \dots, c_r$ can be solely determined by p and m .

For instance, when $p = 3$, we have $r = (m-1)/3$ and $c_2 = \dots = c_r = 1$. For the general case see the proof of Lemma 1. If we choose $e = \phi(c) - 1$ in Theorem 1, we can get an identity between incomplete Kloosterman sums and incomplete hyper-Kloosterman sums.

Theorem 2. *Let $c > 1$ be an odd integer such that $p^2 | c$ for any prime factor p of c . Let $m > 1$ be an integer satisfying $m \equiv 1 \pmod{\phi(c)}$. For any z with $(z, c) = 1$ and any x with $1 \leq x \leq c$ we have*

$$\sum_{\substack{1 \leq b_1 \leq x, \\ b_2, \dots, b_m \pmod{c}, \\ (b_1, c) = \dots = (b_m, c) = 1}} e\left(\frac{1}{c}(b_1 + \dots + b_m + z\bar{b}_1 \dots \bar{b}_m)\right) \\ = c^{(m-1)/2} \left(\prod_{\substack{p|c \text{ to an} \\ \text{odd power}}} \left(\frac{c_2 \dots c_r}{p}\right) \varepsilon_p^r \right) \sum_{\substack{1 \leq b \leq x, \\ (b, c) = 1}} e\left(\frac{1}{c}(b + z\overline{(2-m)}\bar{b})\right).$$

2. MELLIN TRANSFORMS

2.1. Mellin transforms of hyper-Kloosterman sums. Following the methods used in Ye [9] we can deduce an identity between hyper-Kloosterman sum and the classical Kloosterman sum. Moreover, this identity can be written in terms of incomplete exponential sums. Let p be an odd prime and $q = p^a$ with $a \geq 1$. Fix a nonzero integer e and an additive character ψ_p of order zero of the local field $\mathbb{Q}_p$. Denote by R_p the ring of integers in $\mathbb{Q}_p$ and by $R_p^\times$ the group of invertible elements

of R_p . Let S be a subset of $R_p^\times$ such that $S = S(1 + qR_p)$. For $m > 1$ we define an incomplete exponential sum

$$Kl_m(z; e; S) = \sum_{\substack{b_1 \in S/(1+qR_p), \\ b_2, \dots, b_m \in R_p^\times/(1+qR_p)}} \psi_p \left(\frac{1}{q} \left(b_1 + \dots + b_m + \frac{zb_1^e}{b_2 \dots b_m} \right) \right)$$

where $z \in R_p^\times$. If we take $e = -1$, then $Kl_m(z; e; S)$ becomes an incomplete hyper-Kloosterman sum. Let χ be any multiplicative character of $\mathbb{Q}_p$. Then the Mellin transform of this incomplete exponential sum is

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ &= q^m \int_{\substack{z \in R_p^\times, \\ b_1 \in S, \\ b_2, \dots, b_m \in R_p^\times}} \chi^{-1}(z) \psi_p \left(\frac{1}{q} \left(b_1 + \dots + b_m + \frac{zb_1^e}{b_2 \dots b_m} \right) \right) dz db_1 \dots db_m. \end{aligned}$$

Changing variables from z to $x = zb_1^e/(b_2 \dots b_m q)$ and from b_i to $y_i = b_i/q$ for $i = 1, \dots, m$, we get

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ (1) \quad &= q^{-1} \chi^{-1}(q^{m-e}) \\ & \cdot \int_{\substack{y_1 \in q^{-1}S, \\ x, y_2, \dots, y_m \in q^{-1}R_p^\times}} \chi^{-1}(xy_1^{-e}y_2 \dots y_m) \psi_p \left(x + y_1 + \dots + y_m \right) dx dy_1 \dots dy_m. \end{aligned}$$

In this section we will assume that $a > 1$ and consider three cases: (i) χ is ramified with its conductor exponent $a(\chi) = a$, (ii) χ is ramified with $a(\chi) \neq a$, and (iii) χ is unramified.

2.2. The case of $a(\chi) = a > 1$. In this case the integral $\int_{q^{-1}R_p^\times} \chi^{-1}(x) \psi_p(x) dx$ equals the local ε -factor $\varepsilon(\chi, \psi_p)$. Therefore

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ &= q^{-1} \chi^{-1}(q^{m-e}) \left(\varepsilon(\chi, \psi_p) \right)^m \int_{q^{-1}S} \chi^{-1}(y_1^{-e}) \psi_p(y_1) dy_1. \end{aligned}$$

We need a lemma to evaluate $\varepsilon(\chi, \psi_p)^m$:

Lemma 1. *Let p be an odd prime satisfying $(p, m) = (p-1, m) = 1$ and $m \equiv 1 \pmod{\phi(q)}$. Let χ be a ramified character with the conductor exponent $a(\chi) =$*

$a > 1$. Set $q = p^a$. Then

$$\begin{aligned} & \left(\varepsilon(\chi, \psi_p; dx) \right)^{m-1} \\ &= q^{(m-1)/2} \chi \left(\frac{q^{m-1}}{2-m} \right) && \text{if } a \text{ is even;} \\ &= q^{(m+r-1)/2} \chi \left(\frac{q^{m-1}}{2-m} \right) \\ &\quad \cdot \int_{z_1, \dots, z_r \in p^{(a-1)/2} R_p} \chi \left(1 + z_1^2 + c_2 z_2^2 + \dots + c_r z_r^2 \right) dz_1 \cdots dz_r \\ & && \text{if } a \text{ is odd} \end{aligned}$$

where $1 < r \leq m-3$ and $(c_2, p) = \dots = (c_r, p) = 1$. Here $r \equiv 0 \pmod{2}$, and $c_2, \dots, c_r \in \mathbf{Z}$ are solely determined by p and m .

Proof. We start from computation of $\varepsilon(\chi, \psi_p; dx)^m$:

$$\begin{aligned} \left(\varepsilon(\chi, \psi_p; dx) \right)^m &= \int_{(q^{-1} R_p^\times)^m} \chi^{-1}(x_1 \cdots x_m) \psi_p(x_1 + \dots + x_m) dx_1 \cdots dx_m \\ &= q^m \chi(q^m) \int_{(R_p^\times)^m} \chi^{-1}(y_1^m y_2 \cdots y_m) \\ &\quad \cdot \psi_p \left(\frac{y_1(1 + y_2 + \dots + y_m)}{q} \right) dy_1 \cdots dy_m \end{aligned}$$

where we changed variables from $x_1, \dots, x_m$ to $y_1 = qx_1$, $y_2 = qx_2/y_1$, $\dots$, $y_m = qx_m/y_1$. Since $m \equiv 1 \pmod{\phi(q)}$ and the conductor exponent of χ is a , we have $\chi(y_1^m) = \chi(y_1)$. Consequently

$$\begin{aligned} \left(\varepsilon(\chi, \psi_p; dx) \right)^m &= q^{m-1} \chi(q^{m-1}) \int_{\substack{x \in q^{-1} R_p^\times, \\ y_2, \dots, y_m \in R_p^\times}} \chi^{-1}(xy_2 \cdots y_m) \\ &\quad \cdot \psi_p \left(x(1 + y_2 + \dots + y_m) \right) dx dy_2 \cdots dy_m \end{aligned}$$

where $x = y_1/q$. From $a(\chi) = a > 1$ we conclude that the integral with respect to x is non-zero only when $1 + y_2 + \dots + y_m \in R_p^\times$. Therefore

$$\begin{aligned} \left(\varepsilon(\chi, \psi_p; dx) \right)^m &= q^{m-1} \chi(q^{m-1}) \varepsilon(\chi, \psi_p; dx) \\ (2) \quad &\cdot \int_{\substack{y_2, \dots, y_m \in R_p^\times, \\ 1 + y_2 + \dots + y_m \in R_p^\times}} \chi \left(\frac{1 + y_2 + \dots + y_m}{y_2 \cdots y_m} \right) dy_2 \cdots dy_m. \end{aligned}$$

To compute the integral with respect to $y_2, \dots, y_m$ we set $y_2 = y(1+u)$ with $u \in p^{[(a+1)/2]} R_p$ and $y \in R_p^\times / (1 + p^{[(a+1)/2]} R_p)$ with $y \notin -(1 + y_3 + \dots + y_m) + pR_p$. Then the integral with respect to y_2 becomes a sum with respect to y of integrals

with respect to u :

$$\begin{aligned} \sum_y \int_{p^{[(a+1)/2]} R_p} \chi\left(\frac{1+y(1+u)+y_3+\cdots+y_m}{yy_3\cdots y_m(1+u)}\right) du \\ = \sum_y \chi\left(\frac{1+y+y_3+\cdots+y_m}{yy_3\cdots y_m}\right) \\ \cdot \int_{p^{[(a+1)/2]} R_p} \chi\left(1 - \frac{1+y_3+\cdots+y_m}{1+y+y_3+\cdots+y_m}u\right) du \end{aligned}$$

because any higher powers of u are contained in the conductor of χ . We observe that the integrand on the right side is an additive character of u ; hence the integral with respect to u is non-zero only if $1+y_3+\cdots+y_m \in p^{[a/2]}R_p$. Note here $[a/2] \geq 1$ because $a > 1$. Therefore the integral on the right side of (2) is actually taken over $y_2, \dots, y_m \in R_p^\times$ with $1+y_3+\cdots+y_m \in p^{[a/2]}R_p$. By similar arguments we conclude that for this integral the variables $y_2, \dots, y_m \in R_p^\times$ satisfy conditions

$$\begin{aligned} 1+y_3+\cdots+y_m &\in p^{[a/2]}R_p, \\ 1+y_2+y_4+\cdots+y_m &\in p^{[a/2]}R_p, \\ &\dots\dots\dots \\ 1+y_2+\cdots+y_{m-1} &\in p^{[a/2]}R_p. \end{aligned}$$

From these conditions we can see that $y_3, \dots, y_m \in y_2 + p^{[a/2]}R_p$. Then the condition $1+y_3+\cdots+y_m \in p^{[a/2]}R_p$ further implies that $1+(m-2)y_2 \in p^{[a/2]}R_p$, i.e., $y_2, \dots, y_m \in -1/(m-2) + p^{[a/2]}R_p$. Setting $y_i = -(1+x_i)/(m-2)$ with $x_i \in p^{[a/2]}R_p$ for $i = 2, \dots, m$, we can write the integral on the right side of (2) as

$$\begin{aligned} \int_{(p^{[a/2]}R_p)^{m-1}} \chi\left((2-m)^{m-2} \frac{1+x_2+\cdots+x_m}{(1+x_2)\cdots(1+x_m)}\right) dx_2 \cdots dx_m \\ (3) \quad = \chi\left((2-m)^{m-2}\right) \int_{(p^{[a/2]}R_p)^{m-1}} \chi\left(1 - \sum_{2 \leq i < j \leq m} x_i x_j\right) dx_2 \cdots dx_m. \end{aligned}$$

Since $m \equiv 1 \pmod{\phi(q)}$, we can replace $\chi\left((2-m)^{m-2}\right)$ by $\chi^{-1}(2-m)$. When $a > 1$ is even, this last integral equals

$$\begin{aligned} \int_{\substack{y_2, \dots, y_m \in R_p^\times, \\ 1+y_2+\cdots+y_m \in R_p^\times}} \chi\left(\frac{1+y_2+\cdots+y_m}{y_2 \cdots y_m}\right) dy_2 \cdots dy_m \\ (4) \quad = q^{(1-m)/2} \chi^{-1}(2-m). \end{aligned}$$

When $a > 1$ is odd, the integrand on the right side of equation (3) becomes $\chi(1-x_2(x_3+\cdots+x_m)-\sum_{2 < i < j \leq m} x_i x_j)$, and hence the integral with respect to x_2 is nonzero only if $x_3+\cdots+x_m \in p^{(a+1)/2}R_p$. When this is the case the integral with respect to x_2 equals $p^{-(a-1)/2}$. We then can set $x_3 = -(x_4+\cdots+x_m)+p^{(a+1)/2}R_p$.

By integrating with respect to x_3 we get

$$\begin{aligned} & \int_{\substack{y_2, \dots, y_m \in R_p^\times, \\ 1+y_2+\dots+y_m \in R_p^\times}} \chi^{-1} \left(\frac{1+y_2+\dots+y_m}{y_2 \cdots y_m} \right) dy_2 \cdots dy_m \\ &= p^{-a} \chi^{-1}(2-m) \\ & \cdot \int_{x_4, \dots, x_m \in p^{(a-1)/2} R_p} \chi \left(1 + (x_4 + \dots + x_m)^2 - \sum_{3 < i < j \leq m} x_i x_j \right) dx_4 \cdots dx_m. \end{aligned}$$

To complete the square for x_4 we set $z_1 = x_4 + (x_5 + \dots + x_m)/2 \in p^{(a-1)/2} R_p$; the integrand then becomes

$$\chi \left(1 + z_1^2 + \frac{3}{4}(x_5 + \dots + x_m)^2 - \sum_{4 < i < j \leq m} x_i x_j \right).$$

If $p \neq 3$ we can complete the square for x_5 . If $p = 3$, then the integrand simplifies to

$$\chi \left(1 + z_1^2 - \sum_{4 < i < j \leq m} x_i x_j \right)$$

and we can repeat the above computation to integrals with respect to $x_5, \dots, x_m$. By an induction argument we conclude that

$$\begin{aligned} & \int_{\substack{y_2, \dots, y_m \in R_p^\times, \\ 1+y_2+\dots+y_m \in R_p^\times}} \chi^{-1} \left(\frac{1+y_2+\dots+y_m}{y_2 \cdots y_m} \right) dy_2 \cdots dy_m \\ (5) \quad &= q^{(r-m+1)/2} \chi^{-1}(2-m) \\ & \cdot \int_{z_1, \dots, z_r \in p^{(a-1)/2} R_p} \chi \left(1 + z_1^2 + c_2 z_2^2 + \dots + c_r z_r^2 \right) dz_1 \cdots dz_r \end{aligned}$$

when $a > 1$ is odd, where $1 < r \leq m-3$, $r \equiv m-1 \pmod{2}$, and $c_2, \dots, c_r \in R_p^\times$. Since $m \equiv 1 \pmod{\phi(q)}$, we have $m \equiv 1 \pmod{2}$ and $r \equiv 0 \pmod{2}$. By changing variables we can further assume that $c_2, \dots, c_r$ are nonzero integers relatively prime to p . The results from (2), (4), and (5) prove the Lemma. Q.E.D.

When $a > 1$ is even, Lemma 1 implies that

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) K_l(z; e; S) dz \\ &= q^{(m-3)/2} \chi^{-1} \left(q^{1-e} (2-m) \right) \varepsilon(\chi, \psi_p) \int_{q^{-1}S} \chi^{-1}(y^{-e}) \psi_p(y) dy. \end{aligned}$$

Since $a(\chi) = a$ the above can be written as

$$\begin{aligned} & q^{(m-3)/2} \chi^{-1} \left(q^{1-e} (2-m) \right) \int_{q^{-1}R_p^\times} \chi^{-1}(x) \psi_p(x) dx \int_{q^{-1}S} \chi^{-1}(y^{-e}) \psi_p(y) dy \\ &= q^{(m-1)/2} \chi^{-1}(2-m) \int_{R_p^\times} \chi^{-1}(z) dz \sum_{y \in S/(1+qR_p)} \psi_p \left(\frac{y + zy^e}{q} \right) \end{aligned}$$

where we set $z = q^{1-e}xy^{-e}$ and then changed variables from y to y/q . If we change variables from z to $x = (2-m)z$ we finally get

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ (6) \quad &= q^{(m-1)/2} \int_{R_p^\times} \chi^{-1}(x) dx \sum_{y \in S/(1+qR_p)} \psi_p \left(\frac{1}{q} \left(y + \frac{xy^e}{2-m} \right) \right). \end{aligned}$$

When $a(\chi) = a > 1$ is odd we get that

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ &= q^{(m+r-3)/2} \chi \left(\frac{q^{e-1}}{2-m} \right) \\ & \cdot \int_{(p^{(a-1)/2}R_p)^r} \chi \left(1 + z_1^2 + c_2 z_2^2 + \cdots + c_r z_r^2 \right) dz_1 \cdots dz_r \\ & \cdot \int_{\substack{x \in q^{-1}R_p^\times, \\ y \in q^{-1}S}} \chi^{-1}(xy^{-e}) \psi_p(x+y) dx dy. \end{aligned}$$

Changing variables from y to y/q and from x to z by

$$x = \frac{zy^e}{(2-m)q} \left(1 + z_1^2 + c_2 z_2^2 + \cdots + c_r z_r^2 \right)$$

we get

$$\begin{aligned} & q^{(m+r+1)/2} \int_{R_p^\times} \chi^{-1}(z) dz \int_S \psi_p \left(\frac{1}{q} \left(y + \frac{zy^e}{2-m} \right) \right) dy \\ & \cdot \int_{(p^{(a-1)/2}R_p)^r} \psi_p \left(\frac{zy^e}{(2-m)q} \left(z_1^2 + c_2 z_2^2 + \cdots + c_r z_r^2 \right) \right) dz_1 \cdots dz_r. \end{aligned}$$

Changing variables from $z_1, \dots, z_r$ to $x_i = z_i p^{(1-a)/2} \in R_p$ for $i = 1, \dots, r$, the innermost integral becomes

$$\begin{aligned} & p^{r(1-a)/2} \int_{(R_p)^r} \psi_p \left(\frac{zy^e}{p} \left(z_1^2 + c_2 z_2^2 + \cdots + c_r z_r^2 \right) \right) dz_1 \cdots dz_r \\ &= q^{-r/2} \gamma \left(\frac{2zy^e}{p}, \psi_p \right) \gamma \left(\frac{2c_2 zy^e}{p}, \psi_p \right) \cdots \gamma \left(\frac{2c_r zy^e}{p}, \psi_p \right). \end{aligned}$$

Here we used the formula

$$\int_{R_p} \psi_p\left(\frac{bx^2}{2}\right) dx = |b|_p^{-1/2} \gamma(b, \psi_p)$$

for $|b|_p > 1$, where $\gamma(b, \psi_p)$ is the Weil constant.

Now we want to choose a special character ψ_p . Let $\varphi = \varphi_{\mathbb{R}} \prod_p \varphi_p$ be a character of the adèle ring $\mathbf{A}$ of $\mathbb{Q}$ which is trivial on $\mathbb{Q}$ such that $\varphi_{\mathbb{R}}(x) = e^{2\pi i x} = e(x)$ and the order of φ_p is zero for every prime p . Let $c = p_1^{a_1} \cdots p_s^{a_s}$ be an odd positive integer where $p_1, \dots, p_s$ are distinct primes and every $a_j > 1$ for $j = 1, \dots, s$. Now we define a character ψ_{p_j} for each $j = 1, \dots, s$ by $\psi_{p_j}(x) = \bar{\varphi}_{p_j}(xp_j^{a_j}/c)$ for any $x \in \mathbb{Q}_p$. Then the order of each ψ_{p_j} is still zero. When $p = p_j$ we have

$$\begin{aligned} & q^{-r/2} \gamma\left(\frac{2zy^e}{p}, \psi_p\right) \gamma\left(\frac{2c_2zy^e}{p}, \psi_p\right) \cdots \gamma\left(\frac{2c_rzy^e}{p}, \psi_p\right) \\ &= q^{-r/2} \gamma\left(\frac{2zy^e p_j^{a_j}}{pc}, \bar{\varphi}_p\right) \gamma\left(\frac{2c_2zy^e p_j^{a_j}}{pc}, \bar{\varphi}_p\right) \cdots \gamma\left(\frac{2c_rzy^e p_j^{a_j}}{pc}, \bar{\varphi}_p\right). \end{aligned}$$

Because of $p > 2$ and our specific choice of character φ , we have

$$\gamma\left(\frac{2c_izy^e p_j^{a_j}}{pc}, \bar{\varphi}_p\right) = \left(\frac{2c_izy^e (\overline{cp_j^{-a_j}})}{p}\right) \varepsilon_p$$

for $i = 1, \dots, r$, where $x\bar{x} \equiv 1 \pmod{p}$. Here we use the Legendre symbol, and set $\varepsilon_p = 1$ if $p \equiv 1 \pmod{4}$ and $\varepsilon_p = i$ if $p \equiv 3 \pmod{4}$. Consequently

$$\begin{aligned} & \int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz \\ &= q^{(m+1)/2} \int_{R_p^\times} \chi^{-1}(z) dz \int_S \psi_p\left(\frac{1}{q}\left(y + \frac{zy^e}{2-m}\right)\right) dy \\ & \quad \cdot \left(\frac{2zy^e (\overline{cp_j^{-a_j}})}{p}\right) \left(\frac{2c_2zy^e (\overline{cp_j^{-a_j}})}{p}\right) \cdots \left(\frac{2c_rzy^e (\overline{cp_j^{-a_j}})}{p}\right) \varepsilon_p^r \\ &= q^{(m-1)/2} \int_{R_p^\times} \chi^{-1}(z) dz \sum_{y \in S/(1+qR_p)} \psi_p\left(\frac{1}{q}\left(y + \frac{zy^e}{2-m}\right)\right) \\ & \quad \cdot \left(\frac{c_2 \cdots c_r}{p}\right) \varepsilon_p^r \end{aligned} \tag{7}$$

because r is even, when $a > 1$ is odd and $p = p_j$ for $j = 1, \dots, s$.

2.3. The case of $a(\chi) \neq a > 1$. In this case we have

$$\int_{R_p^\times} \chi^{-1}(z) Kl_m(z; e; S) dz = 0 \tag{8}$$

because the integral with respect to x on the right side of (1) vanishes. On the other hand the expressions on the right side of (6) and (7) are equal to zero because the integrals there with respect to x vanish.

2.4. The case of unramified χ . Now we assume that χ is unramified. Then the right side of (1) vanishes, because $q = p^a$ with $a > 1$. Therefore (8) holds in this case. By the same reason, the right side of (6) and (7) are zero because the integrals with respect to z vanish.

3. IDENTITIES OF EXPONENTIAL SUMS

3.1. Local identities. By the equations in (6) and (7) and results in Subsections 2.3 and 2.4, we prove the following identities of incomplete exponential sums in the case of $a > 1$.

Theorem 3. *Let S be a subset of $R_p^\times$ satisfying $S(1 + qR_p) = S$ where $q = p^a$ with $a > 1$ and p being an odd prime. Let e be a nonzero integer and $m > 1$ an integer congruent to 1 modulo $\phi(q)$, where ϕ is the Euler function. Then for c , p , and ψ_p chosen above we have*

$$\begin{aligned}
 & \sum_{\substack{b_1 \in S/(1+qR_p), \\ b_2, \dots, b_m \in R_p^\times/(1+qR_p)}} \psi_p \left(\frac{1}{q} \left(b_1 + \dots + b_m + \frac{zb_1^e}{b_2 \dots b_m} \right) \right) \\
 (9) \quad &= q^{(m-1)/2} \sum_{y \in S/(1+qR_p)} \psi_p \left(\frac{1}{q} \left(y + \frac{zy^e}{2-m} \right) \right) \\
 & \quad \text{when } a > 1 \text{ is even} \\
 &= q^{(m-1)/2} \left(\frac{c_2 \dots c_r}{p} \right) \varepsilon_p^r \sum_{y \in S/(1+qR_p)} \psi_p \left(\frac{1}{q} \left(y + \frac{zy^e}{2-m} \right) \right) \\
 & \quad \text{when } a > 1 \text{ is odd,}
 \end{aligned}$$

for any $z \in R_p^\times$, where r with $1 < r \leq m$ and $c_2, \dots, c_r$ with $(c_2, p) = \dots = (c_r, p) = 1$ are solely determined by p and m .

3.2. The proof of Theorem 1. These local identities imply Theorem 1. To see this we use the same odd $c = p_1^{a_1} \dots p_s^{a_s}$. For each j let S_j be a subset of $R_{p_j}^\times$ such that $S_j(1 + p_j^{a_j} R_{p_j}) = S_j$. Let $m > 1$ be an integer which satisfies $m \equiv 1 \pmod{\phi(c)}$, where ϕ is the Euler function. Fix a nonzero integer e . For any integer z relatively prime to c Theorem 3 implies that

$$\begin{aligned}
 & \prod_{1 \leq j \leq s} \sum_{\substack{b_{1j} \in S_j/(1+p_j^{a_j} R_{p_j}), \\ b_{2j}, \dots, b_{mj} \in R_{p_j}^\times/(1+p_j^{a_j} R_{p_j})}} \psi_{p_j} \left(\frac{1}{p_j^{a_j}} \left(b_{1j} + \dots + b_{mj} + \frac{zb_{1j}^e}{b_{2j} \dots b_{mj}} \right) \right) \\
 &= \prod_{1 \leq j \leq s} p_j^{a_j(m-1)/2} \sum_{y_j \in S_j/(1+p_j^{a_j} R_{p_j})} \psi_{p_j} \left(\frac{1}{p_j^{a_j}} \left(y_j + \frac{zy_j^e}{2-m} \right) \right) \\
 & \cdot \prod_{\substack{1 \leq j \leq s, \\ \text{with } a_j \text{ being odd}}} \left(\frac{c_{2j} \dots c_{r_j j}}{p_j} \right) \varepsilon_{p_j}^{r_j}
 \end{aligned}$$

where r_j and $c_{2j}, \dots, c_{r_j j}$ are the constants in Theorem 3 for $p = p_j$, $j = 1, \dots, s$. Let S be the subset of a reduced residue system modulo c which consists of numbers b_1 satisfying $b_1 \equiv b_{1j} \pmod{p_j^{a_j}}$ for some $b_{1j} \in S_j/(1 + p_j^{a_j} R_{p_j})$, for all $j = 1, \dots, s$, according to the Chinese remainder theorem. Then the above equality can be

written as

$$\begin{aligned} & \sum_{\substack{b_1 \in S, \\ b_2, \dots, b_m \bmod c, \\ (b_2, c) = \dots = (b_m, c) = 1}} \prod_{1 \leq j \leq s} \psi_{p_j} \left(\frac{1}{p_j^{a_j}} (b_1 + \dots + b_m + z b_1^e \bar{b}_2 \dots \bar{b}_m) \right) \\ &= c^{(m-1)/2} \left(\prod_{\substack{1 \leq j \leq s, \\ \text{with } a_j \text{ odd}}} \left(\frac{c_{2j} \dots c_{r_j j}}{p_j} \right) \varepsilon_{p_j}^{r_j} \right) \sum_{y \in S} \prod_{1 \leq j \leq s} \psi_{p_j} \left(\frac{y + z \left(\overline{2-m} \right) y^e}{p_j^{a_j}} \right). \end{aligned}$$

We rewrite the above in terms of $\bar{\varphi}_{p_j}$. Note that $\bar{\varphi}_p(x/c) = 1$ for any integer x , if $p \neq p_1, \dots, p_s$. Since the global character φ is trivial on $\mathbb{Q}$, we can replace the product of $\bar{\varphi}_{p_j}$ by $\varphi_{\mathbb{R}}$ and the equation above becomes

$$\begin{aligned} & \sum_{\substack{b_1 \in S, \\ b_2, \dots, b_m \bmod c, \\ (b_2, c) = \dots = (b_m, c) = 1}} e \left(\frac{1}{c} (b_1 + \dots + b_m + z b_1^e \bar{b}_2 \dots \bar{b}_m) \right) \\ &= c^{(m-1)/2} \left(\prod_{\substack{1 \leq j \leq s, \\ \text{with } a_j \text{ odd}}} \left(\frac{c_{2j} \dots c_{r_j j}}{p_j} \right) \varepsilon_{p_j}^{r_j} \right) \sum_{y \in S} e \left(\frac{1}{c} \left(y + z \left(\overline{2-m} \right) y^e \right) \right). \end{aligned}$$

This completes the proof of Theorem 1.

Q.E.D.

ACKNOWLEDGEMENT

The author would like to express thanks to Wenzhi Luo for helpful discussions.

REFERENCES

- [1] D. Bump, W. Duke, J. Hoffstein, and H. Iwaniec, An estimate for the Hecke eigenvalues of Maass forms, *Inter. Math. Res. Notices*, **4** (1992), 75-81. MR **93d**:11047
- [2] H. Davenport and H. Hasse, Die Nullstellen der Kongruenzzetafunktionen in gewissen zyklischen Fällen, *J. reine angew. Math.*, **172** (1935), 151-182.
- [3] H. Jacquet and Y. Ye, Relative Kloosterman integrals for $GL(3)$, *Bulletin de la Société mathématique de France*, **120** (1992), 263-295. MR **94c**:11047
- [4] W. Luo, Bounds for incomplete hyper-Kloosterman sums, preprint.
- [5] W. Luo, Z. Rudnick, and P. Sarnak, On Selberg's eigenvalue conjecture, *Geom. Funct. Anal.*, **5** (1995), 387-401. MR **96h**:11045
- [6] Z. Mao and S. Rallis, A trace formula for dual pairs, *Duke Math. J.*, **87** (1997), 321-341. CMP 97:11
- [7] Y. Ye, Kloosterman integrals and base change for $GL(2)$, *J. reine angew. Math.* **400** (1989), 57-121. MR **90i**:11134
- [8] Y. Ye, The lifting of Kloosterman sums, *J. Number Theory*, **51** (1995), 275-287. MR **97a**:11126
- [9] Y. Ye, The lifting of an exponential sum to a cyclic algebraic number field of a prime degree, *Transactions Amer. Math. Soc.* **350** (1998), 5003-5015. CMP 97:08
- [10] Y. Ye, Exponential sums for $GL(n)$ and their applications to base change, *J. Number Theory* **68** (1998), 112-130. CMP 98:07
- [11] D. Zagier, Modular forms associated to real quadratic fields, *Invent. Math.* **30** (1975), 1-46. MR **52**:3062

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242-1419

E-mail address: yey@math.uiowa.edu

Current address: Department of Mathematics, The University of Hong Kong, Hong Kong