A SEPARABLE SPACE WITH NO SCHAUDER DECOMPOSITION

G. ALLEXANDROV, DENKA KUTZAROVA, AND A. PLICHKO

(Communicated by Dale Alspach)

ABSTRACT. We combine some known results to remark that there exists a separable Banach space which fails to have a Schauder decomposition. It can be chosen as a subspace of Gowers-Maurey space without any unconditional basic sequence.

The following problem was raised in [Si] (Problem 15.1, p. 494): Does every separable Banach space have a Schauder decomposition? This question goes back to J. R. Retherford [R].

Recall that a sequence $\{X_n\}_{n=1}^\infty$ of closed subspaces of a Banach space X is said to be a Schauder decomposition of X if every $x \in X$ has a unique representation of the form $x = \sum_{n=1}^\infty x_n$, with $x_n \in X_n$ for every n .

Let GM be Gowers-Maurey space which does not contain any unconditional basic sequence [GM]. As was observed by W. B. Johnson, GM has in fact a stronger property, namely it is hereditarily indecomposable (*H.I.*); i.e., no infinite-dimensional closed subspace can be written as a direct sum $Y \oplus Z$, where Y and Z are infinite-dimensional closed subspaces. It is known that every block subspace of GM contains uniform copies of ℓ_1^n . This follows from the lower f -estimate and Krivine's theorem as in [S]. Then, by Szankowski's refinement of Enflo's criterion (see [LT2, p. 111, Remark 1]), we immediately obtain the following.

Proposition. *There exists a subspace X of GM which does not have the compact approximation property (*C.A.P.*).*

Remark 1. For the same purpose we can as well use other *H.I.* spaces constructed after the breakthrough of W. T. Gowers and B. Maurey. For example, there are subspaces without the *C.A.P.* of the super-reflexive *H.I.* spaces in [F] in the case when they contain uniform copies of ℓ_p^n for $p \neq 2$. One can also use the asymptotic ℓ_1 hereditarily indecomposable spaces constructed in [AD] and [ADKM]. The existence of uniform copies of ℓ_1^n in these spaces follows directly from the definition and one does not need to apply Krivine's theorem. Therefore, they also have subspaces without the *C.A.P.*

Corollary. *The space X is an example of a separable Banach space with no Schauder decomposition.*

Received by the editors November 9, 1998.

1991 *Mathematics Subject Classification.* Primary 46B15.

The authors were partially supported by the Bulgarian Ministry of Education and Science under contract MM 506/95.

Proof. Assume the contrary, i.e. X has a Schauder decomposition $\{X_n\}_{n=1}^\infty$.

Case 1. $\{X_n\}_{n=1}^\infty$ is a finite-dimensional decomposition. This is impossible since the existence of an *F.D.D.* implies *B.A.P.* which in turn implies *C.A.P.* (see [LT1]) and this contradicts the above Proposition.

Case 2. There exists m such that X_m is infinite-dimensional. Denote $Y = [X_n : n \neq m]$. Then $X = X_m \oplus Y$ which is also impossible because X_m and Y are closed infinite-dimensional subspaces of X , X is a closed subspace of GM , and GM is *H.I.* $\square$

Remark 2. Clearly, the result is true hereditarily in all the above mentioned *H.I.* spaces, e.g. we have that every subspace of GM has a further subspace which has no Schauder decomposition.

REFERENCES

- [AD] S. Argyros and I. Deliyanni, *Examples of asymptotic ℓ_1 spaces*, Trans. Amer. Math. Soc. **349** (1997), 973–995. MR **97f**:46021
- [ADKM] S. Argyros, I. Deliyanni, D. Kutzarova and A. Manousakis, *Modified mixed Tsirelson spaces*, J. Funct. Anal. **159** (1998), 43–109. CMP 99:04
- [GM] W. T. Gowers and B. Maurey, *The unconditional basic sequence problem*, J. Amer. Math. Soc. **6** (1993), 851–874. MR **94k**:46021
- [F] V. Ferenczi, *A uniformly convex hereditarily indecomposable Banach space*, Israel J. Math. **102** (1997), 199–225. MR **98m**:46013
- [LT1] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I, Sequence Spaces*, Springer-Verlag, 1977. MR **58**:17766
- [LT2] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II, Function Spaces*, Springer-Verlag, 1979. MR **81c**:46001
- [R] J. R. Retherford, *Some remarks on Schauder bases II*, Rev. Rouman. Math. Pure et Appl. **13** (1968), 521–527. MR **38**:4969
- [S] Th. Schlumprecht, *An arbitrarily distortable Banach space*, Israel J. Math. **76** (1991), 81–95. MR **93h**:46023
- [Si] I. Singer, *Bases in Banach spaces II*, Springer-Verlag, 1981. MR **82k**:46024

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF SOFIA, BULGARIA

INSTITUTE OF MATHEMATICS, BULGARIAN ACADEMY OF SCIENCES, BULGARIA

Current address: Department of Mathematics, University of South Carolina, Columbia, South Carolina 29208

DEPARTMENT OF MATHEMATICS, PEDAGOGICAL UNIVERSITY, KIROVOGRAD, UKRAINE