A GEOMETRIC PROOF OF A THEOREM ABOUT NON-DUAL RENORMINGS

LIBOR VESELÝ

(Communicated by Dale Alspach)

ABSTRACT. We give a simple geometric proof of a result by Davis and Johnson that every nonreflexive Banach space X admits an equivalent norm in which X is not isometric to a dual space. Moreover, our renorming keeps unchanged the original norm on a given finite-codimensional subspace and makes this subspace norm-one complemented.

Let X be a real Banach space. We shall say that a norm $\|\cdot\|$ on X (or the space $(X, \|\cdot\|)$) is *non-dual* if $(X, \|\cdot\|)$ is not isometric to a dual space.

W. J. Davis and W. B. Johnson proved in [DJ] that every nonreflexive Banach space admits a non-dual renorming. This result was strengthened by D. van Dulst and I. Singer ([vDS]), who produced a renorming of any nonreflexive space such that the renormed space is not norm-one complemented in its bidual. Finally, S. V. Konyagin gave a quite simple proof of the following yet stronger result [Ko]: *every nonreflexive Banach space has an equivalent norm in which a three-point set fails to have Chebyshev centers*. (See, e.g., [Ho] for the definition of Chebyshev centers and for the fact that each bounded subset of X admits a Chebyshev center whenever X is norm-one complemented in its bidual.)

We present here a simple short geometric proof of a stronger form of the result by Davis and Johnson (Corollary 2). Here “simple” means that it uses only classical tools (James’ theorem, Krein-Šmul’yan theorem) and not results from renorming theory.

For a Banach space X , we denote by B_X its closed unit ball and by X^* its dual space.

Theorem. *Let $(X, \|\cdot\|)$ be a nonreflexive Banach space. Then there exists a norm $\|\cdot\|$ on $X \oplus \mathbb{R}$ such that*

- (a) $\|(x, 0)\| = \|x\|$;
- (b) *the natural projection $P: X \oplus \mathbb{R} \rightarrow X$, $P(x, t) = x$, has norm one*;
- (c) $(X \oplus \mathbb{R}, \|\cdot\|)$ *is non-dual*.

Proof. Let $f \in X^*$ be any functional that does not attain its norm (cf. [Ja]). Let us denote

$$L = B_X \cap f^{-1}(0), \quad C = \overline{\text{conv}} \left[(B_X \times \{0\}) \cup (L \times \{1\}) \cup (L \times \{-1\}) \right].$$

Received by the editors September 22, 1998 and, in revised form, November 30, 1998.
 1991 *Mathematics Subject Classification*. Primary 46B03; Secondary 46B10.

(The reader is invited to use an easily made diagram.) Then $B_{X \oplus_1 \mathbb{R}} \subset C \subset B_{X \oplus_\infty \mathbb{R}}$, and C is symmetric. (The symbols $\oplus_1$ and $\oplus_\infty$ denote respectively the ℓ_1 -sum and the ℓ_∞ -sum.) Thus C is the unit ball of an equivalent norm $\|\cdot\|$ on $X \oplus \mathbb{R}$, and moreover,

$$B_X = \{x \in X : (x, 0) \in C\} \subset P(C) \subset P(B_{X \oplus_\infty \mathbb{R}}) = B_X.$$

Thus (a), (b) hold.

To prove (c), suppose that $(X \oplus \mathbb{R}, \|\cdot\|)$ is isometric to Z^* . If we denote by w^* the weak-star topology $\sigma((X \oplus \mathbb{R}, \|\cdot\|), Z)$, then C is w^* -compact, and hence also the set

$$L_0 := L \times \{0\} = ((0, 1) + C) \cap ((0, -1) + C)$$

is w^* -compact. This implies that also

$$(f^{-1}(0) \times \mathbb{R}) \cap C = L \times [-1, 1] = \text{conv} \left[((0, 1) + L_0) \cup ((0, -1) + L_0) \right]$$

is w^* -compact. By the Krein-Šmul'yan theorem (cf. [D-S] or [Sch]), $f^{-1}(0) \times \mathbb{R}$ is w^* -closed since it is bw^* -closed. But $f^{-1}(0) \times \mathbb{R}$ is the kernel of the functional $F = (f, 0) \in X^* \oplus \mathbb{R} = (X \oplus \mathbb{R})^*$. Hence F can be identified with an element of Z , and F attains its norm on C . In other words, $F^{-1}(\|F\|)$ intersects C . But this is in contradiction with the fact that f does not attain its norm on B_X . Indeed, $f^{-1}(\|f\|)$ does not intersect B_X , and (by (a), (b)) $\|f\| = \|F\|$; thus $F^{-1}(\|F\|) = f^{-1}(\|f\|) \times \mathbb{R}$ does not intersect $B_X \times \mathbb{R}$ and the latter set contains C . $\square$

Corollary 1. *For every positive integer n , each nonreflexive Banach space is isometric to an n -codimensional norm-one complemented subspace of a non-dual Banach space.*

Proof. Apply the Theorem n times. $\square$

From Corollary 1, we obtain the following strengthening of the theorem by Davis and Johnson.

Corollary 2. *Let E be a nonreflexive Banach space and let $X \subset E$ be a proper closed subspace of finite codimension. Then E has an equivalent non-dual norm which coincides with the original norm on X and makes X norm-one complemented.*

Proof. Observe that E is isomorphic with $X \oplus \mathbb{R}^n$ for $n = \text{codim } X$ (where the norm on X is the one inherited from E), and X is nonreflexive. Apply Corollary 1. $\square$

REFERENCES

- [DJ] W. J. DAVIS and W. B. JOHNSON, *A renorming of non-reflexive Banach spaces*, Proc. Amer. Math. Soc. **37** (1973), 486–488. MR **46**:9693
- [vDS] D. van DULST and I. SINGER, *On Kadec-Klee norms on Banach spaces*, Studia Math. **54** (1976), 205–211. MR **52**:14937
- [D-S] N. DUNFORD and J. T. SCHWARTZ, *Linear Operators I*, New York 1958. MR **22**:8302
- [Ho] R. B. HOLMES, *A course in Optimization and Best Approximation*, Lecture Notes in Mathematics 257, Springer-Verlag, 1972. MR **54**:8381
- [Ja] R. C. JAMES, *Reflexivity and the supremum of linear functionals*, Ann. Math. **66** (1957), 159–169. MR **19**:755g

- [Ko] S. V. KONYAGIN, *A remark on renormings of nonreflexive spaces and the existence of a Chebyshev center*, Moscow Univ. Math. Bull. **43**, no. 2 (1988), 55–56. MR **89g**:46033
- [Sch] H. H. SCHAEFER, *Topological Vector Spaces*, Graduate Texts in Mathematics, Vol. 3, Springer-Verlag, Berlin 1971. MR **49**:7722

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA C. SALDINI 50,
20133 MILANO, ITALY

E-mail address: `libor@vmimat.mat.unimi.it`