UNIFORM BOUNDS FOR STABLY INTEGRAL POINTS ON ELLIPTIC CURVES

PATRICIA L. PACELLI

(Communicated by Ron Donagi)

ABSTRACT. We show that a conjecture by Lang and Vojta regarding integral points on varieties of logarithmic general type implies the existence of a uniform bound on the number of stably S -integral points on an elliptic curve over a degree- d number field.

1. INTRODUCTION

Many recent results have shown how various conjectures in arithmetic geometry imply uniform boundedness results for points on curves. S. Lang [L] has conjectured that the rational points on a variety of general type are not Zariski dense, a conjecture which can be viewed as a generalization to higher dimensions of the Mordell Conjecture. L. Caporaso, J. Harris, and B. Mazur [CHM] use this conjecture to prove a fascinating result; they show that Lang's conjecture implies a uniform bound on the number of rational points on a curve of genus bigger than 1 over a fixed number field. In [P], the author shows that this bound depends solely on the genus and the degree of the number field, extending a result by D. Abramovich ([A1]).

Abramovich has also shown that a conjecture made by Lang and P. Vojta [V] regarding the non-denseness of integral points on varieties of logarithmic general type implies a result similar to the [CHM] result, but for stably integral points on elliptic curves. More precisely, let X be a variety of logarithmic general type (see Definition 1.5) defined over a number field K . Let S be a finite set of places of K and let $\mathcal{O}_{K,S}$ be the ring of S -integers. If $\mathcal{X}$ is a model of X over $\text{Spec}(\mathcal{O}_{K,S})$, Lang and Vojta conjecture that the set of S -integral points of $\mathcal{X}$ is not Zariski dense, a generalization of both Mordell's conjecture and Siegel's theorem. Now, let E be an elliptic curve defined over a number field K , with S as above. A point $P \in E(K)$ is said to be stably S -integral, i.e. $P \in E(K, S)$, if P is S -integral on the complement of the origin after semistable reduction (see Definition 1.7 for a more precise statement). In [A2], Abramovich shows that a consequence of the Lang-Vojta conjecture is the existence of a uniform bound, depending on K and S , on the number of stably S -integral points of E . In this paper we show that,

Received by the editors September 2, 1997.

1991 *Mathematics Subject Classification.* Primary 11Gxx, 14Gxx.

It is a pleasure to thank Dan Abramovich and Glenn Stevens for encouraging me to pursue this work. Dan Abramovich was also kind enough to read earlier versions of this paper and provide extremely helpful comments.

once again, this bound is independent of the actual number field and, in fact, only depends on S and the degree of K . Our main theorem is the following:

Theorem 1.1. *Assume the Lang-Vojta conjecture. Fix an integer $d > 0$ and fix a finite set of places S_0 in $\mathbf{Q}$ containing 3. Then there exists a bound $N(d, S_0)$ depending on d and S_0 , such that for any number field K of degree d , for any finite set S of places of K lying over S_0 , and for any elliptic curve E defined over K , it follows that $|E(K, S)| \leq N(d, S_0)$, where $E(K, S)$ denotes the set of stably S -integral points of E .*

If the elliptic curve E with which we start is actually semistable, then stably S -integral points are simply S -integral points. This leads immediately to the following corollary, just as in [A2]:

Corollary 1.2. *Assume the Lang-Vojta conjecture. Fix an integer $d > 0$ and fix a finite set of places S_0 in $\mathbf{Q}$ containing 2 and 3. Then there exists a bound $N(d, S_0)$, depending on d and S_0 , such that for any number field K of degree d , for any finite set S of places of K lying over S_0 , and for any semistable elliptic curve E defined over K , it follows that $|E(\mathcal{O}_{K,S})| \leq N(d, S_0)$.*

We have another corollary, which is that the Lang-Vojta conjecture combined with Theorem 1.1 implies L. Merel's theorem on uniform boundedness for torsion points on elliptic curves. Of course, Merel's theorem is unconditional, but it is perhaps interesting to see here the connections with Lang-Vojta and stably integral points. More specifically, we have:

Corollary 1.3. *Assume the Lang-Vojta conjecture. Fix a positive integer d , and let E/K be an elliptic curve over any number field K of degree d . Then there exists a bound $N(d)$ on the order of the group of torsion points of $E(K)$.*

Proof. By [KM], it is enough to bound the number of torsion points of prime order on any elliptic curve E over any number field K of degree d . We can add level 3 structure to E , producing a field extension L of K of bounded degree; say $[L : K] \leq B$, where E/L has semistable reduction over all primes $p > 3$. Define a set S as follows:

$$S = \{v \in M_K^0 \mid \text{ord}_v(p) \geq p - 1 \text{ for some primes } p \text{ such that } p - 1 \leq B \cdot d\},$$

where T is the set of places of L lying over S and over the primes 2 and 3. To see this, first, suppose that $p - 1 > B \cdot d$. Then if P is not integral in $E(L)$ for some place v , it follows from Theorem VIII.7.1 in [S] that $\text{ord}_v(p) \geq p - 1 > B \cdot d$. This is a contradiction, however, as $\text{ord}_v(p)$ is at most the degree of the field. Now suppose that $p - 1 \leq B \cdot d$. If P is not integral for some place v , it follows from above that $v \in T$.

The above claim says that torsion points of prime order in $E(K)$ are stably S -integral, where the set S depends on the degree d . But Theorem 1.1 implies that there is a bound, $N(d)$, on the number of such points, and so the corollary is proven. $\square$

The methods used in this paper to prove Theorem 1.1 are very close to those of [P], and much of the definitions and notation will be similar, as we show that the ideas of that paper apply in this new setting.

Recall the following definitions:

Definition 1.4. A line bundle L on a variety X is *big* if for high values of k , global sections of $H^0(X, L^{\otimes k})$ determine a birational map of X into projective space.

Definition 1.5. Let X be a quasi-projective variety over $\mathbf{C}$. Let $f : Y \rightarrow X$ be a resolution of singularities, so that f is a proper, birational morphism, and Y is a smooth variety. Let $Y \subset \overline{Y}$ be a projective compactification such that $\overline{Y}$ is smooth, and the reduced variety $D = \overline{Y} \setminus Y$ is a divisor of normal crossings. Let $\omega_{\overline{Y}}$ be the dualizing sheaf of $\overline{Y}$. Then X is said to be of *logarithmic general type* if $\omega_{\overline{Y}}(D)$ is big.

Let X be a quasi-projective variety of logarithmic general type defined over a number field K . Let S be a finite set of places of K , and let R be the ring of S integers. Choose a model $\mathcal{X}$ of X over R . The following is a conjecture made by Lang and Vojta.

Conjecture 1.6 (Lang-Vojta). *The set of integral points $\mathcal{X}(R)$ is not Zariski dense in $\mathcal{X}$.*

We now give a precise definition of stably S -integral points:

Definition 1.7. Let E/K be an elliptic curve, S a finite set of places of K , and $P \in E(K)$. Assume E admits a semistable model $\mathcal{E}$ over $\mathcal{O}_{L,T}$, where L is a finite extension of K and T is the set of places of L above S . Let $\mathcal{E}_0 = \mathcal{E} \setminus 0$. We say P is *stably S -integral*, denoted $P \in E(K, S)$, if $P \in \mathcal{E}_0(\mathcal{O}_{L,T})$, i.e., if P is T -integral on the complement of the origin of the semistable model $\mathcal{E}$.

Remark 1.8. Suppose that S contains all places of K dividing 2 and 3, and that $v \notin S$ is a place at which E has additive reduction. Then $P \in E(K)$ is stably S -integral if P does not go through the zero component of the Néron model of E at v . (See [A2], section 4, for a discussion of stably S -integral points in terms of stably minimal models of elliptic curves.)

Remark 1.9. After adding level 3 structure, an elliptic curve admits semistable reduction away from the prime 3. Thus, a point $P \in E(K)$ is stably S -integral if it is S -integral on the Néron model after adding level 3 structure to E . Note also that stably S -integral points remain stably S -integral after field extensions.

Example 1.10. Let $E/\mathbf{Q}$ be the curve given by

$$y^2 = x^3 + 5.$$

Let $S = \{2, 3\}$, and let $P = (-1, 2)$. The curve E has additive reduction at the prime 5, but making the substitution

$$x = \sqrt[3]{5}x', \quad y = \sqrt{5}y',$$

we obtain a semistable model of E over $L = \mathbf{Q}(\sqrt[6]{5})$ given by

$$\mathcal{E} : y'^2 = x'^3 + 1.$$

Under this change of variables, the point P is sent to the point $\left(\frac{-1}{\sqrt[3]{5}}, \frac{2}{\sqrt{5}}\right)$. Therefore, if T is the set of places of L lying over S , we see that P is not T -integral, and so P is not a stably S -integral point of E .

Example 1.11. Let $E/\mathbf{Q}$ be the curve given by

$$E : y^2 = x^3 + 5x,$$

and let $S = \{2, 3\}$ and $P = (20, 90)$. Once again, E has additive reduction at 5, but if we take $a = \sqrt{-5}$ and apply the change of coordinates:

$$x = ax', \quad y = a^{\frac{3}{2}}y',$$

we obtain a semistable model for E over the field $L = \mathbf{Q}(\sqrt{a})$:

$$\mathcal{E} : y'^2 = x'^3 - x'.$$

Under the above change of variables, the point P is sent to the point $(-4a, 18\sqrt{a})$, which is integral. Thus, P is stably S -integral on E .

In [RS], K. Rubin and A. Silverberg study the universal family $E_1 \rightarrow \mathbf{P}^1$ of semistable elliptic curves over $\mathbf{C}$ with full level 3 structure. The surface E_1 can be identified with the total space of the elliptic pencil:

$$\lambda(X^3 + Y^3 + Z^3) - 3\mu XYZ = 0,$$

which maps to $\mathbf{P}^1$ via $[\lambda : \mu]$. We choose the section of the pencil over the point $[X : Y : Z] = [1 : -1 : 0]$ as the origin of the elliptic surface, and denote this origin by Θ .

Remark 1.12. Given any elliptic curve E/K as in Theorem 1.1, it is possible to add level 3 structure by replacing K with an extension of bounded degree. This ensures that E now appears as a fiber of the family E_1 , and, moreover, does not affect our result, since the bound in Theorem 1.1 depends on the degree of the number field. (A more detailed argument of a similar situation in the case of d -gonal curves is given in section 5 of [P].)

Fix a positive integer d and a finite set of places S_0 of $\mathbf{Q}$ as in the statement of Theorem 1.1.

We fix an integer n , and let $E_n \rightarrow \mathbf{P}^1$ be the n -th fibered product of E_1 over $\mathbf{P}^1$: $E_n = (E_1)_{\mathbf{P}^1}^n$. We will be working with subvarieties of E_n^d , where $E_n^d \rightarrow (\mathbf{P}^1)^d$. Denote by Y_n the symmetric product $\text{Sym}^d(E_n)$.

For $i = 1, \dots, d$ and for $j = 1, \dots, n$, we have projections

$$f_i : E_n^d \rightarrow E_n$$

$$h_j : E_n \rightarrow E_1.$$

Let Θ_n be the divisor on E_n^d which comes from the pullbacks of the origin Θ :

$$\Theta_n = \sum_{i=1}^d \sum_{j=1}^n f_i^* h_j^*(\Theta).$$

Similarly, for any positive integer k we will denote by Θ_{n+k} the corresponding divisor on E_{n+k}^d . We will let Θ_{Y_n} denote the image of Θ_n in the quotient Y_n .

Suppose K is any number field of degree d , S a finite set of places of K lying over S_0 , and E an elliptic curve defined over K . Let $\sigma_1, \dots, \sigma_d$ denote the d embeddings of K into $\bar{K}$. Suppose that $P_1, \dots, P_n$ are in $E(K, S)$. Recall that this implies the existence of a field extension L of K , and a set of places T lying over S , along with a semistable model $\mathcal{E}$ of E over $\mathcal{O}_{L,T}$, such that $P_i \in (\mathcal{E} \setminus 0)(\mathcal{O}_{L,T})$ for each i . By Remark 1.12 we may assume that E/L has level 3 structure and appears as a fiber of the surface E_1 .

It follows that the n -tuple $(P_1, \dots, P_n)$ of points of $E(K, S)$ gives rise to a T -integral point on $E_n \setminus \left(\sum_{j=1}^n h_j^* \Theta\right)$, and hence also yields a T -integral point $y_{P_1, \dots, P_n} \in Y_n \setminus \Theta_{Y_n}$:

$$y_{P_1, \dots, P_n} = \{(P_1, \dots, P_n)^{\sigma_1}, \dots, (P_1, \dots, P_n)^{\sigma_d}\}.$$

Definition 1.13. A point $y_{P_1, \dots, P_n} \in Y_n \setminus \Theta_{Y_n}$ is said to be *off-diagonal* if the points $P_i \in E(K, S)$ are pairwise distinct.

Definition 1.14. Choose $m \geq n$. An off-diagonal integral point $y_{(P_1, \dots, P_n)} \in Y_n$ is *m-prolongable* if :

1. There exists an off-diagonal point $y_{(P_1, \dots, P_m)} \in Y_m \setminus \Theta_{Y_m}$ such that the restriction of $y_{(P_1, \dots, P_m)}$ to the first n coordinates yields $y_{(P_1, \dots, P_n)}$.
2. Each P_i , $i = 1, \dots, m$, is defined over the same number field K of degree d , but is not defined over any smaller field.

We will call $E_n^{(m)}$ the set of m -prolongable points in $Y_n \setminus \Theta_{Y_n}$, and will denote by $F_n^{(m)}$ the Zariski closure $\overline{E_n^{(m)}}$. Let $F_n = \bigcap_{m > n} F_n^{(m)}$, and let Θ_{F_n} be the restriction of Θ_{Y_n} to F_n . We will later denote by $\Theta_{F_{n+k}}$ the restriction of $\Theta_{Y_{n+k}}$ to F_{n+k} . Note that by the Noetherian property of the Zariski topology, we know that $F_n = F_n^{(m_0)}$ for some m_0 .

I claim that our goal is to show that, for some n , F_n is empty. A short argument shows that this suffices to prove Theorem 1.1: whenever a fiber of E_1 contains a large number of integral points, say m , then for any $n < m$, it follows that $E_n^{(m)}$ is non-empty. If Theorem 1.1 is not true, then m grows without bound as one varies the fibers of E_1 , implying that for arbitrarily large n , the F_n 's will be non-empty. Moreover, this implies that for all smaller n 's the F_n 's will also be non-empty, due to the surjectivity of the natural projection maps $F_{n+1} \rightarrow F_n$.

Just as in [P], Lemmas 2.6 and 2.7, the projection $F_{n+1} \rightarrow F_n$ is surjective, and each component of F_n has a component of F_{n+1} above it of positive relative dimension. We will perform an induction argument on the relative dimension of F_{n+1} over F_n . We know that this dimension can't be greater than d , as this is the relative dimension of Y_{n+1} over Y_n , and we also know that it is at least 1. What we will do is show that, for any l , if the relative dimension of F_{n+1} over F_n is at least l , then we obtain a contradiction, so it must be at least $l + 1$. Eventually we will get to $l = d$ where the argument must stop, and we will have to conclude that each F_n is empty.

Assume that for all n and for all $y \in F_n$, the dimension of the fiber above y in F_{n+1} is greater than or equal to l . Suppose there exists an element $y \in F_n$ with the dimension of the fiber above y in F_{n+1} exactly l . By the semicontinuity of the fiber dimension of projective maps, there is an irreducible component of F_n , call it M_n , with the properties that M_n contains y , and the general fibers in F_{n+1} over M_n have dimension equal to l . It also follows by induction that for any integer k the dimensions of the general fibers in F_{n+k} above M_n are equal to kl .

The next section of the paper will be devoted to proving the following proposition:

Proposition 1.15. *For large values of k and n , suppose that F is a component of $F_{n+k} \setminus \Theta_{F_{n+k}}$ such that F has relative dimension kl over F_n and the image of F in F_{n+1} has relative dimension l over F_n . Then F is of logarithmic general type.*

Remember, however, that by definition the set of integral points in $F_{n+k} \setminus \Theta_{F_{n+k}}$ is dense; the Lang-Vojta conjecture combined with the work above implies a contradiction. Thus we have discovered that, for all n and for all $y \in F_n$, the dimension of the fiber above y in F_{n+1} must be at least $l + 1$.

Continuing by induction on this relative dimension, we will be forced to conclude that the conjecture implies a contradiction, unless F_{n+k} is empty for large values of k and n , hence proving Theorem 1.1.

2. PROOF OF PROPOSITION 1.15

By definition each F_n is contained in $Y_n \setminus \Theta_{Y_n}$. Define G_n to be the subvariety of E_n^d which is the inverse image of F_n under the quotient map given by the action of the symmetric group. Fix n . As in the proof of Lemma 4.1 in [P], we may replace each G_{n+k} , for $k \geq 0$, with an irreducible component of relative dimension kl over G_n .

We have projection maps $\pi : G_{n+1} \rightarrow G_n$ such that $\pi(G_{n+1}) = G_n$. Notice also that $G_{n+k} \subseteq (G_{n+1})_{G_n}^k$ (see the proof of Lemma 4.1 in [P]).

Let ϕ be the projection of G_n to $(\mathbf{P}^1)^d$, and for $i = 1, \dots, d$, let f_i be the d projections from $(\mathbf{P}^1)^d$ to $\mathbf{P}^1$. Let $B = \phi(G_n)$, and set $B_i = f_i \circ \phi(G_n)$, for $i = 1, \dots, d$. Notice that B is irreducible since it is the image of the irreducible G_n , and $B \subset B_1 \times \dots \times B_d$, where B surjects to each B_i . Finally, let $\tilde{B}$ be a desingularization of B with the property that its discriminant locus is a divisor with normal crossings. In other words, we consider the pullback of $E_1^d \rightarrow (\mathbf{P}^1)^d$ to $\tilde{B}$, and require that the discriminant locus be a divisor of normal crossings, where the discriminant locus is the set of points in $\tilde{B}$ which have singular fibers lying above them. Let $g_i : \tilde{B} \rightarrow \mathbf{P}^1$ be the composition of the maps and inclusions in the diagram below:

$$\tilde{B} \rightarrow B \subset B_1 \times \dots \times B_d \subseteq (\mathbf{P}^1)^d \xrightarrow{f_i} \mathbf{P}^1.$$

Define families of curves:

$$C_i = g_i^*(E_1 \rightarrow \mathbf{P}^1) = E_1 \times_{\mathbf{P}^1} \tilde{B}.$$

By an abuse of notation, we shall pull back all G_{n+k} 's ($k \geq 0$) to $\tilde{B}$ and pull back the C_i 's to G_n , but keep the notation the same. Lemma 4.1 in [P] shows that this new G_{n+1} is contained inside the product of d families of curves, $C_1 \times_{G_n} \dots \times_{G_n} C_d$, and it surjects onto each family.

From [P], §4.2, we also know that there exist d surjective generically finite maps, σ_i , from G_{n+1} to a product of l of the C_i 's fibered over G_n , where l is the relative dimension of G_{n+1} over G_n . In particular, for each $i = 1, \dots, d$, we have

$$\sigma_i : G_{n+1} \rightarrow C_{j_1(i)} \times_{G_n} \dots \times_{G_n} C_{j_l(i)},$$

where one of the indices $j_k(i)$ is equal to i .

There is a projection from G_n down to $\tilde{B}$, and we can now obtain maps from G_n to fibered products of families of curves over $\tilde{B}$:

$$G_n \rightarrow C_1^{m_1} \times_{\tilde{B}} \dots \times_{\tilde{B}} C_d^{m_d}.$$

Hence we can express the σ_i 's as:

$$\sigma_i : G_{n+1} \rightarrow C_{j_1(i)} \times_{\tilde{B}} \dots \times_{\tilde{B}} C_{j_l(i)} \times_{\tilde{B}} C_1^{m_1} \times_{\tilde{B}} \dots \times_{\tilde{B}} C_d^{m_d}.$$

We proceed as in [P], §4.2; using the fact that $G_{n+k} \subseteq (G_{n+1})_{G_n}^k$, we apply the σ_i 's to the various copies of G_{n+1} in the above product, obtaining a generically finite map from G_{n+k} to a product of powers of the C_i 's:

$$G_{n+k} \rightarrow C_1^{k_1} \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_d^{k_d}.$$

We shall denote by V_{n+k} (where $k \geq 0$) the product on the right above:

$$V_{n+k} = C_1^{k_1} \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_d^{k_d}.$$

If we increase k , we can increase the above exponents k_i . For instance, in going from G_{n+k} to G_{n+k+1} , we obtain one more copy of G_{n+1} , allowing us to apply one more of the σ_i 's.

We now turn our attention to the divisors on these products of families of curves. This is where our work here differs from that done in [P]. Notice that each C_i contains the divisor $g_i^*(\Theta)$, the pullback of the origin of the surface E_1 . For $i = 1, \dots, d$, we may choose a projection

$$p_i : C_1 \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_d \rightarrow C_i.$$

Let Z denote this product

$$Z = C_1 \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_d,$$

and define a divisor D_Z on Z by

$$D_Z = \sum_{i=1}^d p_i^* g_i^*(\Theta).$$

Denote by Z_i the product

$$Z_i = C_{j_1(i)} \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_{j_l(i)},$$

where, as above, for each i one of the subscripts $j_k(i)$ is equal to i , and define the divisor D_{Z_i} on Z_i by

$$D_{Z_i} = \sum_{k=1}^l p_{j_k(i)}^* g_{j_k(i)}^*(\Theta).$$

Let τ be the map from G_{n+1} to Z induced by the inclusion $G_{n+1} \subset G_n \times_{\tilde{B}} Z$, and let W be the image of G_{n+1} in Z , where $e : W \hookrightarrow Z$ is the inclusion map.

Lemma 2.1. *We have that*

$$e \circ \tau(G_{n+1} \setminus \Theta_{n+1}) \subset Z \setminus D_Z.$$

Moreover, for each $i = 1, \dots, d$, we have dominant morphisms

$$\sigma_i : G_{n+1} \setminus \Theta_{n+1} \rightarrow Z_i \setminus D_{Z_i}.$$

Proof. The first statement follows automatically, since $(e \circ \tau)^*(D_Z) \subset \Theta_{n+1}$, and hence the complement maps to the complement as desired. For the second statement, consider the surjective maps $\sigma_i : G_{n+1} \rightarrow Z_i$. When restricted to $G_{n+1} \setminus \Theta_{n+1}$, their image is dense in $Z_i \setminus D_{Z_i}$. $\square$

For large n , let V_n be previously defined product

$$V_n = C_1^{k_1} \times_{\tilde{B}} \cdots \times_{\tilde{B}} C_d^{k_d}.$$

Define the canonical projections

$$r_i : V_n \rightarrow C_i^{k_i}$$

for $i = 1, \dots, d$, and, for $j_i = 1, \dots, k_i$,

$$q_{j_i} : C_i^{k_i} \rightarrow C_i.$$

Then V_n contains the divisor

$$D_{V_n} = \sum_{i=1}^d \sum_{j_i=1}^{k_i} r_i^* q_{j_i}^* g_i^* \Theta.$$

It follows that $G_n \backslash \Theta_n$ maps generically finitely to $V_n \backslash D_{V_n}$, and for positive k , $G_{n+k} \backslash \Theta_{n+k}$ maps generically finitely to $V_{n+k} \backslash D_{V_{n+k}}$.

From [A2], Lemma 3, we know that the relative dualizing sheaf $\omega_{E_1/\mathbf{P}^1}(\Theta)$ is big. This is the essential ingredient in proving the following lemma.

Lemma 2.2. *For large n , both $\omega_{V_n/\tilde{B}}(D_{V_n})$ and $\omega_{V_n}(D_{V_n})$ are big.*

Proof. The proof follows exactly as in the proof of Lemma 4.5 in [P]. □

Remark 2.3. In order to apply Lemma 4.5 of [P], we need to know that Lemmas 3.5 and 4.4 of [P] still hold when line bundles of the form $L_1 \otimes L_2$ are replaced by invertible sheaves of the form $L_1(D_1) \otimes L_2(D_2)$ where the D_i 's are divisors. For these lemmas to apply to logarithmic canonical differential forms it is important that the sum of the divisors D_i is a reduced divisor, as such forms must have simple poles along the divisor (see [I], Chapter 11). This does hold, however, as the divisors D_{V_n} which we work with here are constructed as sums of pullbacks of *horizontal* divisors via *distinct* maps, and hence are reduced.

Let $\mathcal{G}_k$ be the quotient of $G_{n+k} \backslash \Theta_{n+k}$ modulo the action of the symmetric group. To prove Proposition 1.15 we need to show that $\mathcal{G}_k$ is of log-general type. We know from [A2], Lemma 2, that for all $n > 2$, $V_n \backslash D_{V_n}$ has log-canonical singularities. We also know that V_n has canonical singularities, as it is a family of nodal curves (see, for instance, Lemma 3.3 of [CHM]). These two facts are important, as they show that the singularities do not prohibit the extension of sections to a desingularization. Thus we will be able to use our projections to products of families of curves to pull back log-differential forms to $G_{n+k} \backslash \Theta_{n+k}$, and to pull them back even further to a desingularization.

By Lemma 4 in [A1], what we must do is show that not only is $\omega_{G_{n+k}}(\Theta_{G_{n+k}})$ big, but also that its sections vanish to sufficiently high order on the fixed points of the symmetric group action on $G_{n+k} \backslash \Theta_{n+k}$. This will let us conclude that any desingularization of $\mathcal{G}_k$ is of log-general type, and hence will prove Proposition 1.15.

By Hironaka's theorem we may choose an equivariant resolution of singularities, $\widetilde{G_{n+k}}$, of G_{n+k} , such that the pullback of Θ_{n+k} , call it $\widetilde{\Theta_{n+k}}$, is a divisor of normal crossings. Denote by r the desingularization map

$$r : \widetilde{G_{n+k}} \backslash \widetilde{\Theta_{n+k}} \rightarrow G_{n+k} \backslash \Theta_{n+k}.$$

Let π_0 denote the map from $G_{n+k} \backslash \Theta_{n+k}$ to $G_n \backslash \Theta_n$, and let σ_0 be the map from $G_n \backslash \Theta_n$ to $V_n \backslash D_{V_n}$. For $j = 1, \dots, k$, let π_j denote the map from $G_{n+k} \backslash \Theta_{n+k}$ to $G_{n+1} \backslash \Theta_{n+1}$. Finally, recall that for each i we have the map σ_i from $G_{n+1} \backslash \Theta_{n+1}$

to $Z_i \setminus D_{Z_i}$. Thus, we have

$$\begin{aligned} \pi_0 : G_{n+k} \setminus \Theta_{n+k} &\rightarrow G_n \setminus \Theta_n, \\ \sigma_0 : G_n \setminus \Theta_n &\rightarrow V_n \setminus D_{V_n}, \\ \pi_j : G_{n+k} \setminus \Theta_{n+k} &\rightarrow G_{n+1} \setminus \Theta_{n+1}, \\ \sigma_i : G_{n+1} \setminus \Theta_{n+1} &\rightarrow Z_i \setminus D_{Z_i}. \end{aligned}$$

The remaining constructions and calculations are almost exactly as in [P], the only difference being the divisors. Define

$$V_{1,i} = V_n \times_{\tilde{B}} Z_i,$$

and

$$V_{k,i} = V_n \times_{\tilde{B}} (Z_i)^k_{\tilde{B}}.$$

Let $D_{V_{1,i}}$ and $D_{V_{k,i}}$ be the usual divisors on $V_{1,i}$ and $V_{k,i}$, respectively.

Recall the fact that

$$G_{n+k} \setminus \Theta_{n+k} \subseteq (G_{n+1} \setminus \Theta_{n+1})^k_{G_n \setminus \Theta_n}.$$

We use this to define a map $(\sigma_0, \sigma_i, \dots, \sigma_i)$ from $G_{n+k} \setminus \Theta_{n+k}$ to $V_{k,i} \setminus D_{V_{k,i}}$ in the following manner: apply σ_0 to G_n , and apply the σ_i 's to the copies of G_{n+1} . In other words, if

$$P = (P_1, \dots, P_n, P_{n+1}, \dots, P_{n+k}) \in G_{n+k} \setminus \Theta_{n+k},$$

then

$$(\sigma_0, \sigma_i, \dots, \sigma_i)(P) = (\sigma_0(P_1, \dots, P_n), \sigma_i(P_{n+1}), \dots, \sigma_i(P_{n+k})).$$

Define projections $\rho_i : Z \rightarrow Z_i$

$$\rho_i(C_1 \times_{\tilde{B}} \dots \times_{\tilde{B}} C_d) = C_{j_1(i)} \times_{\tilde{B}} \dots \times_{\tilde{B}} C_{j_l(i)},$$

where we also have

$$\rho_i : Z \setminus D_Z \rightarrow Z_i \setminus D_{Z_i}.$$

Let Φ_{n+1} and Φ_{n+k} be the locus of points fixed by the group action in $G_{n+1} \setminus \Theta_{n+1}$ and $G_{n+k} \setminus \Theta_{n+k}$, respectively. It is important to note that the image in W of the fixed points Φ_{n+1} is not all of W . This follows by the same argument as in Lemma 4.2 of [P].

From Lemma 2.2 we know that $\omega_{V_n}(D_{V_n})$ is big; it also follows that $\omega_{V_{k,i}}(D_{V_{k,i}})$ is big, since the families of curves appear in $V_{k,i}$ with even higher exponents than in V_n .

To finish the proof of Proposition 1.15, we use the fact that $\omega_{V_{k,i}}(D_{V_{k,i}})$ is big to pull back sections to $G_{n+k} \setminus \Theta_{n+k}$ and to $\widetilde{G_{n+k} \setminus \Theta_{n+k}}$, and then show that these sections vanish to sufficiently high order on the locus of fixed points. Unfortunately, we can't simply pull back sections along any one projection; to guarantee that the sections we pull back will vanish along the fixed points, we must pull back sections via many different projections and tensor them together.

If we pull back sections of $\omega_{V_{k,i}}(D_{V_{k,i}})$, we see that

$$r^* \left((\sigma_0, \sigma_i, \dots, \sigma_i)^* \omega_{V_{k,i}}(D_{V_{k,i}}) \right) \hookrightarrow \omega_{\widetilde{G_{n+k} \setminus \Theta_{n+k}}}(\widetilde{\Theta_{n+k}}).$$

Moreover, if we tensor over all $i = 1, \dots, d$, we have that

$$r^* \left(\bigotimes_i (\sigma_0, \sigma_i, \dots, \sigma_i)^* \omega_{V_{k,i}}(D_{V_{k,i}}) \right) \hookrightarrow \omega_{G_{n+k}}^{\otimes d}(\widetilde{d\Theta_{n+k}}).$$

Let's rewrite the left-hand side above. We can break it down into the portion which comes from V_n and the portion which comes from the powers of Z_i :

$$\begin{aligned} & r^* \left(\bigotimes_i (\sigma_0, \sigma_i, \dots, \sigma_i)^* \omega_{V_{k,i}}(D_{V_{k,i}}) \right) \\ &= r^* \pi_0^* \sigma_0^* \omega_{V_n}^{\otimes d}(dD_{V_n}) \otimes r^* \pi_1^* \left(\bigotimes_i \sigma_i^* \omega_{Z_i/\tilde{B}}(D_{Z_i}) \right) \otimes \dots \otimes r^* \pi_k^* \left(\bigotimes_i \sigma_i^* \omega_{Z_i/\tilde{B}}(D_{Z_i}) \right). \end{aligned}$$

Let's isolate and expand one of the last k terms in the above product. We shall use the fact that $\sigma_i = \rho_i \circ e \circ \tau$; hence $\sigma_i^* = \tau^* e^* \rho_i^*$. Letting $1 \leq j \leq k$, we see that:

$$\begin{aligned} & r^* \pi_j^* \left(\bigotimes_i \sigma_i^* \omega_{Z_i/\tilde{B}}(D_{Z_i}) \right) = r^* \pi_j^* \left(\bigotimes_i \tau^* e^* \rho_i^* \omega_{Z_i/\tilde{B}}(D_{Z_i}) \right) \\ &= r^* \pi_j^* \left(\bigotimes_i \left(\tau^* e^* \left(\omega_{C_{j_1(i)/\tilde{B}}}(D_{C_{j_1(i)}}) \otimes \dots \otimes \omega_{C_{j_l(i)/\tilde{B}}}(D_{C_{j_l(i)}}) \right) \right) \right) \\ &= r^* \pi_j^* \tau^* e^* \left(\bigotimes_i \left(\omega_{C_{j_1(i)/\tilde{B}}}(D_{C_{j_1(i)}}) \otimes \dots \otimes \omega_{C_{j_l(i)/\tilde{B}}}(D_{C_{j_l(i)}}) \right) \right) \\ &= r^* \pi_j^* \tau^* e^* \left(\omega_{C_1/\tilde{B}}^{\otimes l_1}(l_1 D_{C_1}) \otimes \dots \otimes \omega_{C_d/\tilde{B}}^{\otimes l_d}(l_d D_{C_d}) \right) \end{aligned}$$

where

$$D_{C_{j_1(i)}} = p_{j_1(i)}^* g_{j_1(i)}^* \Theta \quad , \dots , \quad D_{C_{j_l(i)}} = p_{j_l(i)}^* g_{j_l(i)}^* \Theta$$

and for $i = 1, \dots, d$,

$$D_{C_i} = p_i^* g_i^* \Theta.$$

The exponents $l_1, \dots, l_d$ are defined by the last line above, and each one is positive. The idea is that for each i we know that one of the $j_k(i)$'s is equal to i . Thus in the last line, when we tensor over all i , each $\omega_{C_1/\tilde{B}}(l_1 D_{C_1})$ appears at least once, and we let l_i indicate the number of times it appears.

Let M denote the line bundle $\omega_{C_1/\tilde{B}}^{\otimes l_1}(l_1 D_{C_1}) \otimes \dots \otimes \omega_{C_d/\tilde{B}}^{\otimes l_d}(l_d D_{C_d})$, and let M_W stand for $e^* M$, the pullback of M to W . By similar reasoning as in Lemma 3.5 of [P], we know that M is big on Z . Since W is the image of $G_{n+1} \setminus \Theta_{n+1}$ in $Z \setminus D_Z$, it follows that W surjects to each $C_i \setminus D_{C_i}$. By Lemma 3.5 in [P] this implies that M_W is big on W .

We now know that

$$r^* \pi_0^* \sigma_0^* \omega_{V_n}^{\otimes d}(dD_{V_n}) \otimes r^* \pi_1^* \tau^* M_W \otimes \dots \otimes r^* \pi_k^* \tau^* M_W \hookrightarrow \omega_{G_{n+k}}^{\otimes d}(\widetilde{d\Theta_{n+k}}).$$

Moreover, each term above is big. Recall that the image of the fixed points Φ_{n+1} in W is not equal to all of W ; hence the image forms a proper subvariety. This means that we obtain a non-zero ideal of sections of M_W which vanish on this subvariety. Call this ideal $\mathcal{I}$.

We borrow the following previously mentioned result from [A2], a result which generalizes Lemma 4.1 of [CHM]:

Lemma 2.4 (See [A2], Lemma 4). *Let $X_0 \subset X$ be an open inclusion of an irreducible variety X_0 in a smooth complex projective irreducible X of dimension n , such that the complement $D = X \setminus X_0$ is a divisor of normal crossings. Let G be a finite group acting on X, X_0, D compatibly. Let ω be a G -equivariant logarithmic k -canonical form on X_0 . If at any point x of X_0 which is fixed by some element in G , the form ω vanishes to order at least $C = k(|G| - 1)$, then ω descends to a regular logarithmic k -canonical form on any desingularization of X_0/G .*

We use this to show that log-differential forms on $\widetilde{G_{n+k} \setminus \Theta_{n+k}}$ will descend to smooth forms modulo the group action if they vanish to a prescribed order on the fixed point locus in $\widetilde{G_{n+k} \setminus \Theta_{n+k}}$.

By Lemma 4.4 of [P], we know that, for some sufficiently large integer s , the line bundle $M_W^{\otimes s} \otimes \mathcal{I}$ is big on W . Let $\mathcal{I}_{n+k}$ be the ideal consisting of the locus of fixed points in $G_{n+k} \setminus \Theta_{n+k}$. Then

$$\sum_{j=1}^k \pi_j^{-1} \tau^{-1} \mathcal{I} \subset \mathcal{I}_{n+k},$$

and

$$\prod_{j=1}^k \pi_j^{-1} \tau^{-1} \mathcal{I} \subset \mathcal{I}_{n+k}^k.$$

Hence,

$$\begin{aligned} r^* \pi_0^* \sigma_0^* \omega_{V_n}^{\otimes ds} (ds D_{V_n}) \otimes r^* \left(\bigotimes_{j=1}^k \pi_j^* \tau^* \left(M_W^{\otimes s} \otimes \prod_{j=1}^k \pi_j^{-1} \tau^{-1} \mathcal{I} \right) \right) \\ \hookrightarrow \omega_{G_{n+k}}^{\otimes ds} (ds \widetilde{\Theta_{n+k}}) \otimes \mathcal{I}_{n+k}^k. \end{aligned}$$

We know that $\omega_{V_n}^{\otimes ds} (ds D_{V_n})$ is big, and as in the proof of Lemma 4.4 in [P], for $k > s$, we have that $M_W^{\otimes s} \otimes \prod_{j=1}^k \pi_j^{-1} \tau^{-1} \mathcal{I}$ is big. Thus for $k > s$ the entire left-hand side is big, so there are a sufficiently high number of sections of $\omega_{G_{n+k}}^{\otimes ds} (ds \widetilde{\Theta_{n+k}}) \otimes \mathcal{I}_{n+k}^k$ vanishing to sufficiently high order on the fixed points.

This lets us conclude that any desingularization of the quotient $\widetilde{G_{n+k} \setminus \Theta_{n+k}}$ modulo the symmetric group action is of log-general type, which implies Proposition 1.15.

REFERENCES

- [A1] D. Abramovich. *Uniformité des points rationnels des courbes algébriques sur les extensions quadratiques et cubiques*, C.R. Acad. Sci. Paris, t. 321, Série I (1995), p. 755-758. MR **96g**:14017
- [A2] D. Abramovich. *Uniformity of stably integral points on elliptic curves*, Invent. Math. 127 (1997), p. 307-317. MR **98d**:14033
- [CHM] L. Caporaso, J. Harris, and B. Mazur. *Uniformity of rational points*, J. Amer. Math. Soc. 10 no. 1 (1997), p. 1-35. MR **97d**:14033
- [I] S. Iitaka. *Algebraic Geometry: an Introduction to Birational Geometry of Algebraic Varieties*, Springer-Verlag, (1982). MR **84j**:14001

- [KM] S. Kamienny and B. Mazur. *Rational torsion of prime order in elliptic curves over number fields*, Astrisque no. 228 (1995), p. 81-100. MR **96c**:11058
- [L] S. Lang. *Hyperbolic diophantine analysis*, Bull. A.M.S. 14 (1986), p. 159-205. MR **87h**:32051
- [M] L. Merel. *Bornes pour la torsion des courbes elliptiques sur les corps de nombres*, Invent. Math. 124 (1996), p. 437-449. MR **96i**:11057
- [P] P. Pacelli. *Uniform boundedness for rational points*, Duke Math. J. vol. 88 no. 1 (1997), p. 77-102. MR **98b**:14020
- [RS] K. Rubin and A. Silverberg. *Families of elliptic curves with constant mod p representations*, Elliptic curves, modular forms, and Fermat's last theorem (Hong Kong, 1993), Ser. Number Theory, I, Internat. Press, Cambridge, MA. (1995), p. 148-161. MR **96j**:11078
- [S] J. H. Silverman. *The Arithmetic of Elliptic Curves*, Springer-Verlag, 1986. MR **87g**:11070
- [V] P. Vojta. *A higher dimensional Mordell conjecture*, in Arithmetic Geometry, (G. Cornell and J. H. Silverman, eds.), Springer-Verlag, (1986). CMP 19:03

DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY, NEW YORK,
NEW YORK 10027-6598

E-mail address: `pacelli@math.columbia.edu`