ON RANDOM ALGEBRAIC POLYNOMIALS

K. FARAHMAND

(Communicated by Stanley Sawyer)

ABSTRACT. This paper provides asymptotic estimates for the expected number of real zeros and K -level crossings of a random algebraic polynomial of the form $a_0 \binom{n-1}{0}^{1/2} + a_1 \binom{n-1}{1}^{1/2} x + a_2 \binom{n-1}{2}^{1/2} x^2 + \cdots + a_{n-1} \binom{n-1}{n-1}^{1/2} x^{n-1}$, where $a_j (j = 0, 1, \dots, n-1)$ are independent standard normal random variables and K is a constant independent of x . It is shown that these asymptotic estimates are much greater than those for algebraic polynomials of the form $a_0 + a_1 x + a_2 x^2 + \cdots + a_{n-1} x^{n-1}$.

1. INTRODUCTION

Let $(\Omega, \mathcal{A}, \text{Pr})$ be a fixed probability space and let $\{a_j(\omega)\}_{j=0}^{n-1}$ be a sequence of independent random variables defined on Ω . The random algebraic polynomial was introduced in the pioneer work of Littlewood and Offord [9] and [10] as

$$Q(x) \equiv Q_n(x, \omega) = \sum_{j=0}^{n-1} a_j(\omega) x^j,$$

and since then has been greatly studied. Denote by $N_K(\alpha, \beta)$ the number of real roots of the equation $P(x) = K$ in the interval (α, β) and by $EN_K(\alpha, \beta)$ its expected value. In particular it is shown (for example see Kac [8] or Wilkins [12]) that if the coefficients are assumed to have a standard normal distribution and n is sufficiently large, $EN_0(-\infty, \infty) \sim (2/\pi) \log n$. Recently (see Farahmand [4]), it was shown that this asymptotic value remains valid for $EN_K(-\infty, \infty)$ as long as K is bounded. For K large such that $K^2/n \rightarrow 0$ as $n \rightarrow \infty$, $EN_K(-\infty, \infty)$ is asymptotically reduced to $(1/\pi) \log(n/K^2)$ in $(-1, 1)$ while it remains the same as for $K = 0$ in $(-\infty, -1) \cup (1, \infty)$ ([4]). In contrast, a random trigonometric polynomial

$$T(x) \equiv T_n(x, \omega) = \sum_{j=0}^{n-1} a_j(\omega) \cos j\theta$$

has more roots in $(0, 2\pi)$. In fact $EN_K(0, 2\pi) \sim 2n/\sqrt{3}$ for $K = o(\sqrt{n})$; see [5] and [6].

Motivated by the interesting results obtained in Littlewood and Offord [10] we considered the case when the coefficients $a_j(\omega)$ have variance $1/j!$. It is presumably

Received by the editors July 9, 1997 and, in revised form, December 17, 1997 and February 5, 1998.

1991 *Mathematics Subject Classification.* Primary 60H99; Secondary 42Bxx.

Key words and phrases. Number of real roots, real zeros, random algebraic polynomials, random trigonometric polynomials, Kac-Rice formula.

the case, possibly under some mild conditions for K and for n sufficiently large, that $EN_K(-\infty, \infty)$ is $o(\sqrt{n})$. This author, however, was unable to make any substantial progress towards this conjecture. Instead in this paper we study the polynomials

$$P(x) \equiv P_n(x, \omega) = \sum_{j=0}^{n-1} a_j(\omega) \binom{n-1}{j}^{1/2} x^j.$$

This is, indeed, the same as saying that the j th coefficient of $Q(x)$ has variance $\binom{n-1}{j}$. Besides the mathematical interest, as reported in Edelman and Kostlan [3, page 11], these polynomials have some relationship with physics [1]. We prove the following theorems. Theorem 1 was known to Edelman and Kostlan, however to be complete we give a proof here. Theorem 3, and its comparison with Theorem 1, shows that for $P(x)$ there are as many extrema as the number of zero crossings. Therefore, unlike $Q(x)$, all the oscillations of $P(x)$ are between two zero crossings, asymptotically in expectation.

Theorem 1. *If the coefficients a_j of $P(x)$ are independent standard normal random variables, then*

$$EN_0(-\infty, \infty) = \sqrt{n-1}.$$

Theorem 2. *Denote $\Upsilon(n) = (2n-1)!!/(2n)!!$ and assume $K^2\Upsilon(n) \rightarrow 0$. With the same assumptions as in Theorem 1 for the coefficients of $P(x)$, we have*

$$EN_K(-\infty, \infty) \sim \sqrt{n-1}.$$

Theorem 3. *Denote $M(-\infty, \infty)$ the number of extrema of $P(x)$ in $(-\infty, \infty)$. Then the expected value of $M(-\infty, \infty)$ satisfies*

$$EM(-\infty, \infty) \sim \sqrt{n-1}.$$

2. A FORMULA FOR THE EXPECTED NUMBER OF REAL ROOTS

Let

$$(2.1) \quad A^2 = \text{var}\{P(x) - K\},$$

$$(2.2) \quad B^2 = \text{var}\{P'(x)\},$$

$$(2.3) \quad C = \text{cov}[\{P(x) - K\}, P'(x)]$$

and

$$\eta = -CK/A\sqrt{A^2B^2 - C^2}.$$

Then by using the expected number of level crossings given by Cramer and Leadbetter [2, page 285] for our equation $P(x) - K = 0$, we can obtain

$$EN_K(\alpha, \beta) = \int_{\alpha}^{\beta} \frac{B\sqrt{1 - C^2/A^2B^2}}{A} \phi\left(-\frac{K}{A}\right) [2\phi(\eta) + \eta\{2\Phi(\eta) - 1\}] dx,$$

where as usual

$$\Phi(t) = (2\pi)^{-1/2} \int_{-\infty}^t \exp(-y^2/2) dy$$

and

$$\phi(t) = (2\pi)^{-1/2} \exp(-t^2/2).$$

Let $\Delta^2 = A^2 B^2 - C^2$ and $\operatorname{erf}(x) = \int_0^x \exp(-t^2) dt$; then we can write the extension of a formula obtained by Rice [11] for the case of $K = 0$ as

$$(2.4) \quad EN_K(\alpha, \beta) = I_1(\alpha, \beta) + I_2(\alpha, \beta),$$

where

$$(2.5) \quad I_1(\alpha, \beta) = \int_{\alpha}^{\beta} \frac{\Delta}{\pi A^2} \exp\left(-\frac{B^2 K^2}{2\Delta^2}\right) dx$$

and

$$(2.6) \quad I_2(\alpha, \beta) = \int_{\alpha}^{\beta} \frac{\sqrt{2}KC}{\pi A^3} \exp\left(-\frac{K^2}{2A^2}\right) \operatorname{erf}\left(\frac{KC}{\sqrt{2}A\Delta}\right) dx.$$

3. PROOF OF THEOREM 1

We need the following, where (3.2) and (3.3) are obtained by differentiation of (3.1),

$$(3.1) \quad A^2 = \sum_{j=0}^{n-1} \binom{n-1}{j} x^{2j} = (x^2 + 1)^{n-1},$$

$$(3.2) \quad B^2 = \sum_{j=0}^{n-1} j^2 \binom{n-1}{j} x^{2j-2} = (n-1)(x^2 + 1)^{n-3}(nx^2 - x^2 + 1),$$

and

$$(3.3) \quad C = \sum_{j=0}^{n-1} j \binom{n-1}{j} x^{2j-1} = (n-1)x(x^2 + 1)^{n-2}.$$

Therefore, since from (3.1)-(3.3)

$$(3.4) \quad \Delta^2 = A^2 B^2 - C^2 = (n-1)(x^2 + 1)^{2n-4},$$

from (2.4)-(2.6) we obtain

$$(3.5) \quad \begin{aligned} EN_0(0, \infty) &= \pi^{-1} \int_0^{\infty} \frac{\Delta}{A^2} dx \\ &= \frac{\sqrt{n-1}}{\pi} \int_0^{\infty} \frac{dx}{x^2 + 1} = \frac{\sqrt{n-1}}{2}. \end{aligned}$$

This gives the proof of Theorem 1. It is, indeed, interesting to note that from (3.5)

$$\frac{\sqrt{n-1}}{\pi(x^2 + 1)}$$

is the density function of the number of real zeros of $P(x)$; see also [3, page 12]. Obtaining a closed form of the above density function is uncommon. An asymptotic result, for most cases, is the best that can be achieved.

4. PROOF OF THEOREM 2

Because $|\operatorname{erf} u| < \sqrt{\pi}/2$, it follows that from (2.6), (3.1), and (3.3) that

$$\begin{aligned} 0 \leq I_2(-\infty, \infty) &\leq \frac{|K|}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{C}{A^3} \exp\left(-\frac{K^2}{2A^2}\right) dx \\ &= \frac{2}{\sqrt{\pi}} \int_0^{|K|/\sqrt{2}} e^{-\nu^2} d\nu, \end{aligned}$$

if $\nu = |K|/(A\sqrt{2})$. Therefore, because $\operatorname{erf} u \leq u$ when $u \geq 0$,

$$(4.1) \quad 0 \leq I_2(-\infty, \infty) \leq \frac{2}{\sqrt{\pi}} \operatorname{erf} \left(\frac{K}{\sqrt{2}} \right) \leq \min \left\{ 1, \frac{\sqrt{2}|K|}{\sqrt{\pi}} \right\}.$$

Moreover, it follows from (2.5), (3.1), (3.2) and (3.4) that

$$I_1(-\infty, \infty) = \frac{\sqrt{n-1}}{\pi} \int_{-\infty}^{\infty} \frac{e^{-s}}{x^2+1} dx,$$

in which $s = K^2(nx^2 - x^2 + 1)/\{2(x^2 + 1)^{n-1}\}$. If $x = \tan \theta$, we find that

$$I_1(-\infty, \infty) = \frac{2\sqrt{n-1}}{\pi} \int_0^{\pi/2} e^{-s} d\theta,$$

in which $s = (K^2/2) \{(n-1)\sin^2 \theta + \cos^2 \theta\} \cos^{2n-4} \theta$. Therefore,

$$(4.2) \quad I_1(-\infty, \infty) = \frac{2\sqrt{n-1}}{\pi} \int_0^{\pi/2} \{1 - (1 - e^{-s})\} d\theta = \sqrt{n-1} - R,$$

in which from [7, page 369],

$$\begin{aligned} 0 \leq R &= \frac{2\sqrt{n-1}}{\pi} \int_0^{\pi/2} (1 - e^{-s}) d\theta \leq \frac{2\sqrt{n-1}}{\pi} \int_0^{\pi/2} s d\theta = K^2 \beta_n, \\ \beta_n &= \frac{\sqrt{n-1}(3n-4)(2n-5)!!}{(2n-2)(2n-4)!!} = \sqrt{n-1} \Upsilon(n-2) \frac{3n-4}{2n-2}. \end{aligned}$$

A straight-forward algebraic calculation shows that $\beta_{n+1} < \beta_n$ when $n \geq 2$. We conclude that $0 \leq R \leq K^2 \beta_2 = K^2/2$ and then that $R = o(\sqrt{n})$ because $K^2 = o\{\Upsilon^{-1}(n)\}$ and $\Upsilon(n) \sim \sqrt{n\pi}$. When this last result is combined with (4.1), (4.2) and (2.4), it is clear that Theorem 2 is true. In fact, we have actually proved the better result that

$$-\frac{K^2}{2} \leq EN_k(-\infty, \infty) - \sqrt{n-1} \leq \min \left\{ 1, \frac{\sqrt{2}|K|}{\sqrt{\pi}} \right\},$$

from which we can also infer not only Theorem 2 but also that

$$EN_K(-\infty, \infty) = \sqrt{n-1} + O(1)$$

when K is bounded.

5. EXTREMA

The expected number of extrema of $P(x)$, denoted by $EM(-\infty, \infty)$, is simply the expected number of real zeros of $P'(x) = \sum_{j=1}^{n-1} a_j j \binom{n-1}{j}^{1/2} x^{j-1}$. Therefore we apply $EN_0(-\infty, \infty)$ for $P'(x)$. To this end, by successive differentiation of (2.2)

we obtain

$$\begin{aligned}
 A^2 &= \sum_{j=0}^{n-1} j^2 \binom{n-1}{j} x^{2j-2} \\
 (5.1) \quad &= (n-1)(x^2+1)^{n-3}(nx^2-x^2+1), \\
 C &= \sum_{j=1}^{n-1} j^2(j-1) \binom{n-1}{j} x^{2j-3} \\
 (5.2) \quad &= (n-1)(n-2)x(x^2+1)^{n-4} \{(n-1)x^2+2\}
 \end{aligned}$$

and

$$\begin{aligned}
 B^2 &= \sum_{j=1}^{n-1} j^2(j-1)^2 \binom{n-1}{j} x^{2j-4} \\
 (5.3) \quad &= (n-1)(n-2)(x^2+1)^{n-5} \{(n-1)(n-2)x^4+4(n-2)x^2+2\}.
 \end{aligned}$$

Therefore, from (5.1)-(5.3) we obtain

$$(5.4) \quad \frac{\Delta}{A^2} = \frac{\sqrt{(n-2)\{(n-1)(n-2)x^4+2(n-1)x^2+2\}}}{(x^2+1)\{(n-1)x^2+1\}}.$$

From (5.4) it then follows that, for all non-zero x

$$\lim_{n \rightarrow \infty} \frac{\Delta}{A^2 \sqrt{n-2}} = \frac{1}{x^2+1}.$$

Then since $\Delta/A^2 \sqrt{n-2} \leq \sqrt{2}/(x^2+1)$ for all real x , the dominated convergent theorem for Lebesgue integrals shows that

$$\int_{-\infty}^{\infty} \frac{\Delta}{A^2 \sqrt{n-2}} dx = \int_{-\infty}^{\infty} \frac{dx}{x^2+1} = \pi.$$

Theorem 3 is then an immediate consequence of this result.

ACKNOWLEDGMENT

The author wishes to thank the referee and the probability editor for their comments on the earlier version of this paper.

REFERENCES

- [1] E. Bogomolny, O. Bohigas, and P. Leboeuf. Distribution of roots of random polynomials. *Phys. Rev. Lett.*, 68:2726–2729, 1992. MR **92m**:81054
- [2] H. Cramér and M.R. Leadbetter. *Stationary and Related Stochastic Processes*. Wiley, N.Y., 1967. MR **36**:949
- [3] A. Edelman and E. Kostlan. How many zeros of a random polynomial are real? *Bull. Amer. Math. Soc.*, 32:1–37, 1995. MR **95m**:60082
- [4] K. Farahmand. On the average number of real roots of a random algebraic equation. *Ann. Probab.*, 14:702–709, 1986. MR **87k**:60140
- [5] K. Farahmand. On the average number of level crossings of a random trigonometric polynomial. *Ann. Probab.*, 18:1403–1409, 1990. MR **91i**:60140
- [6] K. Farahmand. Level crossings of a random trigonometric polynomial. *Proc. Amer. Math. Soc.*, 111:551–557, 1991. MR **91f**:60092
- [7] I.S. Gradshteyn and I.M. Ryzhik. *Table of Integrals, Series and Products*. Academic Press, London, 1980, xv+1160 pp. MR **81g**:33001
- [8] M. Kac. On the average number of real roots of a random algebraic equation. *Bull. Amer. Math. Soc.*, 49:314–320, 1943. MR 4,196d

- [9] J.E. Littlewood and A.C. Offord. On the number of real roots of a random algebraic equation. *J. London Math. Soc.*, 13:288–295, 1938.
- [10] J.E. Littlewood and A.C. Offord. On the number of real roots of a random algebraic equation II. *Proc. Camb. Phil. Soc.*, 35:133–148, 1939.
- [11] S.O. Rice. Mathematical theory of random noise. *Bell. System Tech. J.*, 25:46–156, 1945. Reprinted in: *Selected Papers on Noise And Stochastic Processes* (ed. N. Wax), Dover, New York, 1954, 133–294.
- [12] J.E. Wilkins. An asymptotic expansion for the expected number of real zeros of a random polynomial. *Proc. Amer. Math. Soc.*, 103:1249–1258, 1988. MR **90f**:60105

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ULSTER, JORDANSTOWN, CO. ANTRIM BT37 0QB, UNITED KINGDOM

E-mail address: `k.farahmand@ulst.ac.uk`