ON MODELS OF $U_q(sl(2))$ AND q -APPELL FUNCTIONS USING A q -INTEGRAL TRANSFORMATION

VIVEK SAHAI

(Communicated by Hal L. Smith)

ABSTRACT. We discuss few models of the quantum universal enveloping algebra of $sl(2)$ from the special function point of view. Two sets of such models are given, one acting on the space of ${}_1\phi_0$ functions while the other on the space of q -Appell functions. These models are closely related through a q -integral transformation. Some interesting identities are obtained.

1. INTRODUCTION

Quantum groups have generated considerable interest among mathematicians and physicists alike. The theory of q -special functions has found a group theoretic interpretation using the techniques of quantum groups and quantum algebras. In this direction major contributions have come from the works of Floreanini and Vinet [6]–[12], Kalnins and Miller [18], to name a few.

In a recent paper [29], the author has constructed models of irreps of classical $sl(2)$ in terms of difference-differential operators using an integral transformation which is inspired by the well known Euler integral representation for the hypergeometric functions ${}_2F_1$ with extension to some other functions as well. The models in [29] have culminated in various recurrence relations and identities involving ordinary hypergeometric series. To have an extension of this to q -series, we introduce a q -integral transformation, which is a generalization of and is motivated by Thomae's integral representation of ${}_2\phi_1$, given in [13], to construct models of $U_q(sl(2))$, the quantum universal enveloping algebra of $sl(2)$, in terms of difference-dilation operators. These models act on a space of basis functions involving q -Appell functions and give rise to identities involving certain generalized q -hypergeometric series in two and three variables. A q -difference equation satisfied by q -Appell functions is also obtained. Section-wise treatment is as follows.

In Section 2, we list various definitions and results needed for our discussion. In Section 3, we discuss models of $U_q(sl(2))$ and give its two variable models in terms of dilation operators in which ${}_1\phi_0$ functions appear as the basis functions. In Section 4, a q -integral transformation is defined and transforms of certain expressions are obtained. Next, we reproduce two theorems which help us in upgrading the models of Section 3 into new models of $U_q(sl(2))$ in terms of difference-dilation operators

Received by the editors January 15, 1998.

1991 *Mathematics Subject Classification.* Primary 33D80, 81R50.

The author is thankful to University Grants Commission, India, for the award of a Visiting Associateship.

in which the basis functions involve $\Phi^{(2)}$, the q -Appell functions. These models, in Section 5, are exploited for identities, which are believed to be new. Finally in Section 6, we outline the corresponding transformed models for a m -fold q -integral transformation.

2. PRELIMINARIES

The generalized basic (or q -) hypergeometric series is defined by [13]

$$(1) \quad {}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, x \right) = \sum_{n=0}^{\infty} \frac{(a_1, \dots, a_r; q)_n}{(b_1, \dots, b_s; q)_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+s-r} x^n,$$

where

$$(2) \quad (a_1, \dots, a_r; q)_n = \prod_{i=1}^r (a_i; q)_n$$

and q -shifted factorial $(a; q)_n$ is defined by

$$(3) \quad (a; q)_n = \begin{cases} (1-a)(1-aq) \cdots (1-aq^{n-1}), & n = 1, 2, \dots, \\ 1, & n = 0, \\ [(1-aq^{-1}) \cdots (1-aq^{-n})]^{-1}, & n = -1, -2, \dots, \end{cases}$$

$$(4) \quad (a; q)_{\infty} = \prod_{r=0}^{\infty} (1-aq^r).$$

The series ${}_r\phi_s$ terminates if one of the numerator parameter is of the form q^{-m} , $m = 0, 1, 2, \dots$ and $q \neq 0$. When $0 < |q| < 1$, the series ${}_r\phi_s$ converges absolutely for all z if $r \leq s$, and for $|z| < 1$ if $r = s + 1$. If $|q| > 1$ and $|z| < |b_1 \cdots b_s|/|a_1 \cdots a_r|$, then also ${}_r\phi_s$ converges absolutely. The series ${}_r\phi_s$ diverges for $z \neq 0$ when $0 < |q| < 1$ and $r > s + 1$, and when $|q| > 1$ and $|z| > |b_1 \cdots b_s|/|a_1 \cdots a_r|$, unless it terminates.

We now list q -analogues of some important functions which will be needed in the discussion. The q -analogues of the exponential function are

$$(5) \quad e_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{(q; q)_n} = \frac{1}{(x; q)_{\infty}}, \quad |x| < 1,$$

and

$$(6) \quad E_q(x) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} \frac{x^n}{(q; q)_n} = (-x; q)_{\infty}.$$

$E_q(x)$ converges for all x . Note that $e_q(x)E_q(-x) = 1$. The q -analogue of the binomial function is

$$(7) \quad {}_1\phi_0 \left(\begin{matrix} a \\ - \end{matrix}; x \right) = \frac{(ax; q)_{\infty}}{(x; q)_{\infty}}, \quad |x| < 1, |q| < 1.$$

The gamma function has the following q -analogue:

$$(8) \quad \Gamma_q(\alpha) = \frac{e_q(q^{\alpha})}{e_q(q)} (1-q)^{1-\alpha}, \quad \alpha \neq 0, -1, -2, \dots,$$

and satisfies the equation

$$(9) \quad \Gamma_q(\alpha + 1) = \frac{1-q^{\alpha}}{1-q} \Gamma_q(\alpha).$$

The q -derivative operator is defined as

$$(10) \quad \Delta_z f(z) = \frac{(1 - T_z) f(z)}{(1 - q)z}$$

where the q -dilation operator T_z is defined by $T_z f(z) = f(qz)$. Indeed, as $q \rightarrow 1$, the q -derivative operator becomes an ordinary differential operator, provided f is differentiable at z .

The q -integral is defined as

$$(11) \quad \int_0^1 f(t) d_q t = (1 - q) \sum_{n=0}^{\infty} f(q^n) q^n.$$

If f is continuous on $[0, 1]$, then the q -integral reduces to the ordinary integral.

Also we need the q -analogue of Appell function F_2 , [33], as

$$(12) \quad \Phi^{(2)} \left(\begin{matrix} a; b, b' \\ c, c' \end{matrix}; x, y \right) = \sum_{m,n=0}^{\infty} \frac{(a; q)_{m+n} (b; q)_m (b'; q)_n}{(c; q)_m (c'; q)_n} x^m y^n$$

and the q -analogue of generalized Kampé de Fériet function $F_{E;F;G;H}^{A:B;C;D}$, as

$$(13) \quad \begin{aligned} & \phi_{E;F;G;H}^{A:B;C;D} \left(\begin{matrix} (a_i); (b_j); (c_k); (d_l) \\ (e_m); (f_n); (g_u); (h_v) \end{matrix}; x, y, z \right) \\ &= \sum_{p,r,t=0}^{\infty} \frac{((a_i); q)_{p+r+t} ((b_j); q)_p ((c_k); q)_r ((d_l); q)_t}{((e_m); q)_{p+r+t} ((f_n); q)_p ((g_u); q)_r ((h_v); q)_t} x^p y^r z^t \end{aligned}$$

where (a_i) abbreviates parameters $a_1, \dots, a_A$, and where $((a_i); q)_{p+r+t}$ denotes $(a_1, \dots, a_A; q)_{p+r+t}$, etc. The arguments and the complex parameters in (12) and (13) are so constrained that the multiple series converges.

3. MODELS OF $U_q(sl(2))$

The commutation relations among the generators e, f, h

$$(14) \quad [h, e] = 2e, \quad [h, f] = -2f, \quad [e, f] = \frac{q^{h/2} - q^{-h/2}}{q^{1/2} - q^{-1/2}}$$

define an associative algebra [14], which may be regarded as the q -deformation of the universal enveloping algebra of $sl(2)$, referred to as $U_q(sl(2))$ in this paper. The Casimir element which belongs to the center of $U_q(sl(2))$ is given by

$$(15) \quad \mathbb{C} = \frac{k^2 q^{1/2} + k^{-2} q^{-1/2} - 2}{q^{1/2} - q^{-1/2}} + fe.$$

As the parameter $q \rightarrow 1^-$, these reduce to familiar commutator rules for $sl(2)$.

Let V_q be a complex vector space consisting of q -special functions with a basis $\{\phi_\lambda : \lambda \in S\}$ such that the functions $\{f_\lambda = \lim_{q \rightarrow 1} \phi_\lambda : \lambda \in S\}$ form a basis for the vector space, say V . Let $A(V_q)$ be the associative algebra of all linear operators on V_q over a complex field. Then a representation ρ_q of $sl(2)$ on V_q is defined as a mapping $\rho_q : sl(2) \rightarrow A(V_q)$ satisfying

(i) $\rho_q(ax + by) = a\rho_q(x) + b\rho_q(y)$.

(ii) There exists a Lie algebra representation ρ of $sl(2)$ on V such that

$$\lim_{q \rightarrow 1} \rho_q(x) \phi_\lambda = \rho(x) f_\lambda,$$

for all $x, y \in sl(2)$ and $a, b \in \mathbb{C}$.

This representation ρ_q of $sl(2)$ is said to be irreducible if there is no proper subspace W_q of V_q which is invariant under ρ_q . Define

$$(16) \quad J_q^+ = \rho_q(\mathcal{J}^+), \quad J_q^- = \rho_q(\mathcal{J}^-), \quad J_q^0 = \rho_q(\mathcal{J}^0)$$

where $J_q^+, J_q^-, J_q^0 \in A(V_q)$ and $\{\mathcal{J}^+, \mathcal{J}^-, \mathcal{J}^0\}$ form a basis of $sl(2)$. On his work on irreps of $sl(2)$, Manocha [25] has defined the following commutator rules satisfied by J_q -operators:

$$(17) \quad \begin{aligned} J_q^0 J_q^+ - q J_q^+ J_q^0 &= J_q^+, \\ q J_q^0 J_q^- - J_q^- J_q^0 &= -J_q^-, \\ q J_q^+ J_q^- - J_q^- J_q^+ &= 2q^{2u} J_q^0 - (1-q)q^{2u} J_q^0 J_q^0, \quad u \in C. \end{aligned}$$

If we define an operator C_q on V_q by

$$(18) \quad C_q = q J_q^+ J_q^- + q^{2u} J_q^0 J_q^0 - q^{2u} J_q^0,$$

then it can be verified that

$$(19) \quad \begin{aligned} q J_q^+ C_q &= C_q J_q^+, \\ J_q^- C_q &= q C_q J_q^-, \\ J_q^0 C_q &= C_q J_q^0. \end{aligned}$$

As will be seen, the operator C_q will be used to derive a q -difference equation satisfied by q -Appell function. Further, as $q \rightarrow 1$, the operators J_q^+, J_q^-, J_q^0 reduce to J^+, J^-, J^0 which satisfy the commutation relations obeyed by $sl(2)$ and the operator C_q reduces to the Casimir operator C and the relations (19) exhibit that C commutes with J^+, J^- and J^0 . Hence, as $q \rightarrow 1$, ρ_q reduces to a Lie algebra representation ρ of $sl(2)$.

Notice that the commutation relations given by eqs. (17) are equivalent to that given by (14) in the sense that if the operators e, f, h satisfy (14), then J_q -operators defined by

$$(20) \quad J_q^+ = e, \quad J_q^- = q^{2u+\frac{h-1}{2}} f, \quad J_q^0 = \frac{1-q^{h/2}}{1-q}$$

satisfy (17).

Following the analysis as in Manocha [25] and Miller [27], we have the following theorem:

Theorem 3.1. *Every representation ρ_q of $sl(2)$ is isomorphic to a representation in the following list:*

- (i) *The representation $D_q(\alpha, u)$, $\alpha \in C \setminus \{0\}$, $0 \leq \operatorname{Re} \alpha < 1$, such that $\alpha - 2u$ is not an integer. $S = \{\alpha + n : n = 0, \pm 1, \pm 2, \dots\}$.*
- (ii) *The representation $\uparrow_q(u)$, $u \in C$ and $2u$ is not a non-negative integer. $S = \{0, 1, 2, \dots\}$.*
- (iii) *The representation $D_q(2u)$ where $2u$ is a non-negative integer. $S = \{0, 1, 2, \dots, 2u\}$.*

For each of these representations there is a basis of V_q consisting of vectors $\{f_\lambda\}$, defined for each $\lambda \in S$ such that

$$\begin{aligned} J_q^0 f_\lambda &= \frac{1 - q^{\lambda-u}}{1 - q} f_\lambda, \\ J_q^+ f_\lambda &= \frac{q^{2u} - q^\lambda}{1 - q} f_{\lambda+1}, \\ J_q^- f_\lambda &= -\frac{1 - q^\lambda}{1 - q} f_{\lambda-1}. \end{aligned} \quad (21)$$

Guided by the above theorem, we give below two-variable models of representation $D_q(\alpha)$ in the particular case when $u = 0$.

Model I

$$\begin{aligned} J_q^+ &= t(zT_t\Delta_z + t\Delta_t), \\ J_q^- &= t^{-1}T_z^{-1}(z(1 - zT_t)\Delta_z - t\Delta_t), \\ J_q^0 &= t\Delta_t, \end{aligned} \quad (22)$$

$$f_\lambda = \frac{(q^\lambda z; q)_\infty}{(z; q)_\infty} t^\lambda = {}_1\phi_0\left(\begin{matrix} q^\lambda \\ - \end{matrix}; z\right) t^\lambda, \quad \lambda \in S. \quad (23)$$

Model II

$$\begin{aligned} J_q^+ &= t(z(1 - z)T_t\Delta_z + t\Delta_t), \\ J_q^- &= t^{-1}T_z^{-1}(z\Delta_z - t\Delta_t), \\ J_q^0 &= t\Delta_t, \end{aligned} \quad (24)$$

$$f_\lambda = \frac{(z; q)_\infty}{(q^\lambda z; q)_\infty} t^\lambda = {}_1\phi_0\left(\begin{matrix} q^{-\lambda} \\ - \end{matrix}; q^\lambda z\right) t^\lambda, \quad \lambda \in S, \quad (25)$$

where $S = \{\alpha + n : \alpha \in C \setminus \{0\}, 0 \leq \operatorname{Re} \alpha < 1, n = 0, \pm 1, \pm 2, \dots\}$. The dilation models given above satisfy (17) as well as (21) with $u = 0$.

A model of representation $\uparrow_q$ is same as Model II above with $S = \{0, 1, 2, \dots\}$. However, we concentrate on the model of representation $D_q(\alpha)$ and will return to the representation $\uparrow_q$ later in Section 5, where we obtain identities based on a model of representation $\uparrow_q$.

The models of irreducible representation of $U_q(sl(2))$ can be found in the literature too. For example, in [12], Floreanini and Vinet have considered two such models. Also, models of $U_q(su_2)$ are discussed in detail by Kalnins *et al.* [17, 19, 20, 21] and by Rideau and Winternitz [28]. The metaplectic representation of $su_q(1, 1)$ are presented in [8] and are connected with q -Gegenbauer polynomials. Models of q -harmonic oscillators given by Biedenharn [1] and Mcfarlane [26] are closely related to the representation of $U_q(sl(2))$. Work on these representations has been carried out by Drinfeld [3], Faddeev [5], Kulish and Reshetikhin [23], Sklyanin [31, 32] and Woronowich [34, 35] among others. Some models of q -representations of the Lie algebra $\mathcal{G}(a, b)$ are contained in Sahai [30]. For more details, see Chari and Pressley [2], Jantzen [15], Kassel [22] and Majid [24].

4. A q -INTEGRAL TRANSFORMATION AND TRANSFORMED MODELS

In this section, we introduce a two-fold q -integral transformation, based on Thomae's integral representation of ${}_2\phi_1$ [13, 4], viz.

$$(26) \quad {}_2\phi_1 \left(\begin{matrix} q^\alpha, q^\beta \\ q^\gamma \end{matrix}; q, z \right) = \Gamma_q \left(\begin{matrix} \gamma \\ \beta, \gamma - \beta \end{matrix} \right) \int_0^1 t^{\beta-1} \frac{(tq; q)_\infty}{(tq^{\gamma-\beta}; q)_\infty} {}_1\phi_0 \left(\begin{matrix} q^\alpha \\ - \end{matrix}; q, tz \right) d_q t$$

and then compute transforms of certain operator expressions. These transforms are later used to obtain new models from the ones given in Section 3 in which the basis functions will be in terms of q -Appell function.

Define

$$(27) \quad \begin{aligned} h(\beta, \beta', \gamma, \gamma') &= \mathcal{I}[f(u, v)] = \Gamma_q \left(\begin{matrix} \gamma, \gamma' \\ \beta, \beta', \gamma - \beta, \gamma' - \beta' \end{matrix} \right) \\ &\times \int_0^1 \int_0^1 u^{\beta-1} v^{\beta'-1} \frac{(qu, qv; q)_\infty}{(q^{\gamma-\beta}u, q^{\gamma'-\beta'}v; q)_\infty} f(u, v) d_q u d_q v, \\ \operatorname{Re} \gamma &> \operatorname{Re} \beta > 0, \quad \operatorname{Re} \gamma' > \operatorname{Re} \beta' > 0. \end{aligned}$$

Defining the operators

$$(28) \quad \begin{aligned} E_\beta(h(\beta, \beta', \gamma, \gamma')) &= h(\beta + 1, \beta', \gamma, \gamma'), \\ L_\beta(h(\beta, \beta', \gamma, \gamma')) &= h(\beta - 1, \beta', \gamma, \gamma'), \end{aligned}$$

and denoting $E_{\beta\gamma} = E_\beta E_\gamma$, etc. we have the following transforms of operator expressions needed for our discussion:

$$(29) \quad \begin{aligned} \mathcal{I}[u\Delta_u f] &= x\Delta_x h, \\ \mathcal{I}[v\Delta_v f] &= y\Delta_y h, \\ \mathcal{I}[u^2\Delta_u f] &= \frac{1-q^\beta}{1-q^\gamma} x\Delta_x E_{\beta\gamma} h, \\ \mathcal{I}[v^2\Delta_v f] &= \frac{1-q^{\beta'}}{1-q^{\gamma'}} y\Delta_y E_{\beta'\gamma'} h, \\ \mathcal{I}[\Delta_u f] &= \frac{1-q^{\gamma-1}}{1-q^{\beta-1}} x\Delta_x L_{\beta\gamma} h, \\ \mathcal{I}[\Delta_v f] &= \frac{1-q^{\gamma'-1}}{1-q^{\beta'-1}} y\Delta_y L_{\beta'\gamma'} h, \\ \mathcal{I}[uf] &= \frac{1-q^\beta}{1-q^\gamma} E_{\beta\gamma} h, \\ \mathcal{I}[vf] &= \frac{1-q^{\beta'}}{1-q^{\gamma'}} E_{\beta'\gamma'} h. \end{aligned}$$

To obtain models of $U_q(sl(2))$ in terms of difference-dilation operators (difference in $\beta, \beta', \gamma, \gamma'$ and dilation in x and y) in which the basis functions are in terms of q -Appell functions, we need the following theorems.

Theorem 4.1 ([25]). *Let ρ_q be a representation of $sl(2)$ in terms of $\{J_q^+, J_q^-, J_q^0\}$ with basis functions $\{f_\lambda : \lambda \in S\}$. Then ρ_q is also a representation of $sl(2)$ in terms of $\{K_q^+, K_q^-, K_q^0\}$ with basis functions $\{h_\lambda : \lambda \in S\}$ where*

$$(30) \quad K_q^+ = \mathcal{I}J_q^+ \mathcal{I}^{-1}, \quad K_q^- = \mathcal{I}J_q^- \mathcal{I}^{-1}, \quad K_q^0 = \mathcal{I}J_q^0 \mathcal{I}^{-1}, \quad h_\lambda = \mathcal{I}f_\lambda.$$

Theorem 4.2 ([13]). *If a and b are indeterminates such that $ab = qba$, q commutes with a and b and the associative law holds, then*

$$(31) \quad (a + b)^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q b^k a^{n-k},$$

where $\begin{bmatrix} n \\ k \end{bmatrix}_q$ is the q -binomial coefficient.

To upgrade the two-variable models to three-variable ones, we substitute $z = ux + vy$. This gives

$$(32) \quad z\Delta_z = u\Delta_u + v\Delta_v T_u,$$

where Δ_z is defined by (10) and

$$(33) \quad T_z f(z) = f(qz) = T_u T_v f(ux + vy).$$

Putting $z = ux + vy$, the Model I takes the following shape:

$$(34) \quad \begin{aligned} J_q^+ &= t[(u\Delta_u + v\Delta_v T_u)T_t + t\Delta_t], \\ J_q^- &= t^{-1}T_u^{-1}T_v^{-1}[(1 - T_t(ux + vy))(u\Delta_u + v\Delta_v T_u) - t\Delta_t], \\ J_q^0 &= t\Delta_t, \end{aligned}$$

with the basis functions as

$$(35) \quad f_\lambda = {}_1\phi_0 \left(\begin{matrix} q^\lambda \\ - \end{matrix}; ux + vy \right) t^\lambda, \quad \lambda \in S.$$

The J -model (34) along with (35) is now ready to be transformed into a K -model involving the q -Appell function $\Phi^{(2)}$ as the basis functions with the help of Theorems 4.1 and 4.2. We have

Model IA.

$$(36) \quad \begin{aligned} K_q^+ &= \mathcal{I}J_q^+ \mathcal{I}^{-1} = t[(x\Delta_x + y\Delta_y T_x)T_t + t\Delta_t], \\ K_q^- &= \mathcal{I}J_q^- \mathcal{I}^{-1} \\ &= t^{-1}T_x^{-1}T_y^{-1} \left[x\Delta_x + y\Delta_y T_x - xT_t \left(\frac{1 - q^\beta}{1 - q^\gamma} x\Delta_x E_{\beta\gamma} + \frac{1 - q^\beta}{1 - q^\gamma} y\Delta_y T_x E_{\beta\gamma} \right) \right. \\ &\quad \left. - yT_t \left(\frac{1 - q^{\beta'}}{1 - q^{\gamma'}} x\Delta_x E_{\beta'\gamma'} + \frac{1 - q^{\beta'}}{1 - q^{\gamma'}} y\Delta_y T_x E_{\beta'\gamma'} \right) - t\Delta_t \right], \\ K_q^0 &= \mathcal{I}J_q^0 \mathcal{I}^{-1} = t\Delta_t, \end{aligned}$$

with the basis functions as

$$(37) \quad h_\lambda(x, y, t) = \Phi^{(2)} \left(\begin{matrix} q^\lambda; b, b' \\ c, c' \end{matrix}; x, y \right) t^\lambda, \quad \lambda \in S,$$

where $b = q^\beta, b' = q^{\beta'}, c = q^\gamma, c' = q^{\gamma'}$. Note that Theorem 4.2 is utilized in obtaining $\mathcal{I}({}_1\phi_0) = \Phi^{(2)}$.

Likewise Model II is transformed to the following

Model IIA.

$$\begin{aligned}
(38) \quad K_q^+ &= t \left[(x\Delta_x + y\Delta_y T_x) T_t - xT_t \left(\frac{1-q^\beta}{1-q^\gamma} x\Delta_x E_{\beta\gamma} + \frac{1-q^\beta}{1-q^\gamma} y\Delta_y T_x E_{\beta\gamma} \right) \right. \\
&\quad \left. - yT_t \left(\frac{1-q^{\beta'}}{1-q^{\gamma'}} x\Delta_x E_{\beta'\gamma'} + \frac{1-q^{\beta'}}{1-q^{\gamma'}} y\Delta_y T_x E_{\beta'\gamma'} \right) + t\Delta_t \right], \\
K_q^- &= t^{-1} T_x^{-1} T_y^{-1} (x\Delta_x + y\Delta_y T_x - t\Delta_t), \\
K_q^0 &= t\Delta_t,
\end{aligned}$$

with the basis functions as

$$(39) \quad h_\lambda(x, y, t) = \Phi^{(2)} \left(q^{-\lambda}; b, b', c, c'; q^\lambda x, q^\lambda y \right) t^\lambda, \quad \lambda \in S.$$

The K -operators given above satisfy the following relations:

$$\begin{aligned}
(40) \quad K_q^0 h_\lambda &= \frac{1-q^\lambda}{1-q} h_\lambda, \\
K_q^+ h_\lambda &= \frac{1-q^\lambda}{1-q} h_{\lambda+1}, \\
K_q^- h_\lambda &= -\frac{1-q^\lambda}{1-q} h_{\lambda-1}, \\
C_q h_\lambda &= 0 \text{ where } C_q = qK_q^+ K_q^- + K_q^0 K_q^0 - K_q^0
\end{aligned}$$

as well as the commutation relations

$$\begin{aligned}
(41) \quad K_q^0 K_q^+ - qK_q^+ K_q^0 &= K_q^+, \\
qK_q^0 K_q^- - K_q^- K_q^0 &= -K_q^-, \\
qK_q^+ K_q^- - K_q^- K_q^+ &= 2K_q^0 - (1-q)K_q^0 K_q^0
\end{aligned}$$

and as such give difference-dilation models of $U_q(sl(2))$.

Note that the equation $C_q h_\lambda = 0$ in (40) can be used to obtain a q -difference equation satisfied by $\Phi^{(2)}$. For example, if h_λ is given by (37), then we get the q -difference equation

$$\begin{aligned}
(42) \quad & \left[\{ (1-aq^{-1}) + aq^{-1} (xD_x^+ + yD_y^+ T_x) \} \right. \\
& \times \left\{ a (xD_x^- + yD_y^- T_x) \left(\frac{1-b}{1-c} xT_{bc} + \frac{1-b'}{1-c'} yT_{b'c'} - 1 \right) - (1-a) \right\} \\
& \left. + q(1-a) (1-aq^{-1}) \right] \Phi^{(2)} \left(a; b, b', c, c'; x, y \right) = 0
\end{aligned}$$

where $D_x^+ = x^{-1}(1 - T_x)$ and $D_x^- = x^{-1}(1 - T_x^{-1})$ are q -difference operators [11]. Observe that $\frac{1}{1-q} D_x^+ \rightarrow \frac{\partial}{\partial x}$ and $\frac{1}{1-q^{-1}} D_x^- \rightarrow \frac{\partial}{\partial x}$ as $q \rightarrow 1$.

Further, we have

$$\begin{aligned}
(43) \quad qK_q^+ C_q &= C_q K_q^+, \\
K_q^- C_q &= qC_q K_q^-, \\
K_q^0 C_q &= C_q K_q^0.
\end{aligned}$$

The eqs. (43) will become instrumental in obtaining identities.

5. IDENTITIES

5.1. **Based on Model IA.** We have shown in eq. (40) that

$$(44) \quad h_\lambda(x, y, t) = \Phi^{(2)} \left(\begin{matrix} q^\lambda; b, b' \\ c, c' \end{matrix}; x, y \right) t^\lambda, \quad \lambda \in S,$$

is a solution of $C_q h_\lambda(x, y, t) = 0$ where $C_q = qK_q^+ K_q^- + K_q^0 K_q^0 - K_q^0$. This suggests that

$$(45) \quad \begin{aligned} u(x, y, t) &= \phi_{-1;1;1;1}^{1;1;1;2} \left(\begin{matrix} a : b; b'; b'', b''' \\ - : c; c'; c'' \end{matrix}; x, y, t \right) t^\alpha \\ &= \sum_{n=0}^{\infty} \frac{(a, b'', b'''; q)_n}{(c'', q; q)_n} \Phi^{(2)} \left(\begin{matrix} aq^n; b, b' \\ c, c' \end{matrix}; x, y \right) t^{\alpha+n} \end{aligned}$$

is a solution of $C_q u(x, y, t) = 0$. Using the fact that $qK_q^+ C_q = C_q K_q^+$, we have

$$(46) \quad C_q [e_q(sK_q^+)u](x, y, t) = 0,$$

where

$$(47) \quad \begin{aligned} [e_q(sK_q^+)u](x, y, t) &= \sum_{n=0}^{\infty} \frac{s^n}{(q; q)_n} K_q^{+n} u(x, y, t) \\ &= \frac{(\frac{ast}{1-q}; q)_\infty}{(\frac{st}{1-q}; q)_\infty} \phi_{1;1;1;1}^{1;1;1;2} \left(\begin{matrix} a : b; b'; b'', b''' \\ \frac{ast}{1-q} : c; c'; c'' \end{matrix}; x, y, t \right) t^\alpha. \end{aligned}$$

Now we have, using the Weisner's expansion, as explained in [25, 16]

$$(48) \quad [e_q(sK_q^+)u](x, y, t) = \sum_{n=0}^{\infty} A_n h_{\alpha+n}(x, y, t),$$

where A_n is found by putting $x = y = 0$. Rescaling suitably, we eventually arrive at

$$(49) \quad \begin{aligned} &\frac{(at; q)_\infty}{(t; q)_\infty} \phi_{1;1;1;1}^{1;1;1;2} \left(\begin{matrix} a : b; b'; b'', b''' \\ at : c; c'; c'' \end{matrix}; x, y, \omega t \right) \\ &= \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} \Phi^{(2)} \left(\begin{matrix} aq^n; b, b' \\ c, c' \end{matrix}; x, y \right) {}_3\phi_1 \left(\begin{matrix} q^{-n}, b'', b''' \\ c'' \end{matrix}; q, q^n \omega \right) t^n, \end{aligned}$$

where $\omega = \frac{s}{1-q}$.

5.2. **Based on Model IIA.** We follow exactly the same approach as given in section 5.1 with the change that we use the other q -exponential E_q instead of e_q .

We have shown that $C_q h_\lambda = 0$, where h_λ is given by eq.(39). It therefore follows that

$$(50) \quad u(x, y, t) = \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q, c''; q)_n} \Phi^{(2)} \left(\begin{matrix} a^{-1}q^{-n}; b, b' \\ c, c'; aq^n x, aq^n y \end{matrix} \right) t^{\alpha+n}$$

is a solution of $C_q u(x, y, t) = 0$. Considering that $K_q^- C_q = qC_q K_q^-$, we have

$$(51) \quad C_q [E_q(sK_q^-)u](x, y, t) = 0,$$

where

(52)

$$\begin{aligned} [E_q(sK_q^-)u](x, y, t) &= \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} s^n}{(q; q)_n} K_q^{-n} u(x, y, t) \\ &= \frac{(\frac{\omega}{t}; q)_{\infty}}{(\frac{a\omega}{t}; q)_{\infty}} \sum_{k,n=0}^{\infty} \frac{(a, \frac{a\omega}{t}; q)_n (a^{-1}q^{-n}, b; q)_k}{(c'', q; q)_n (\frac{q^{1-n}t}{a\omega}, c, q; q)_k} \\ &\quad \times {}_2\phi_2 \left(\begin{matrix} a^{-1}q^{-n+k}, b' \\ \frac{q^{1-n+k}t}{a\omega}, c' \end{matrix}; q, \frac{-q^{k+1}yt}{\omega} \right) q^{\binom{k}{2}} \left(\frac{-qxt}{\omega} \right)^k t^{\alpha+n}. \end{aligned}$$

By Weisner's expansion, we have

$$(53) \quad [E_q(sK_q^-)u](x, y, t) = \sum_{n=-\infty}^{\infty} B_n h_{\alpha+n}(x, y, t),$$

where B_n is obtained by putting $x = y = 0$. This leads to

$$\begin{aligned} (54) \quad &\frac{(\frac{\omega}{t}; q)_{\infty}}{(\frac{a\omega}{t}; q)_{\infty}} \sum_{k,n=0}^{\infty} \frac{(a, \frac{a\omega}{t}; q)_n (a^{-1}q^{-n}, b; q)_k}{(c'', q; q)_n (\frac{q^{1-n}t}{a\omega}, c, q; q)_k} \\ &\times {}_2\phi_2 \left(\begin{matrix} a^{-1}q^{-n+k}, b' \\ \frac{q^{1-n+k}t}{a\omega}, c' \end{matrix}; q, \frac{-q^{k+1}yt}{\omega} \right) \left(\frac{-q^{\frac{k+1}{2}}xt}{\omega} \right)^k t^n \\ &= \sum_{n=-\infty}^{\infty} \frac{\Gamma_q(\alpha+n)\Gamma_q(\gamma'')}{\Gamma_q(\alpha)\Gamma_q(\gamma''+n)\Gamma_q(q+1)} {}_2\phi_2 \left(\begin{matrix} aq^n, aq^{n+1} \\ q^{n+1}, c''q^n \end{matrix}; q, \omega \right) \\ &\times \Phi^{(2)} \left(\begin{matrix} a^{-1}q^{-n}; b, b' \\ c, c'; aq^n x, aq^n y \end{matrix} \right) t^n, \end{aligned}$$

where $\omega = \frac{s}{1-q}$.

We have discussed above identities corresponding to models of representation $D_q(\alpha)$. We now consider the Model IIA for the representation $\uparrow_q$ (i.e. eqs. (38) and (39) with $\lambda \in S = \{n = 0, 1, 2, \dots\}$). It can be verified that

$$(55) \quad u(x, y, t) = \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} \Phi^{(2)} \left(\begin{matrix} q^{-n}; b, b' \\ c, c'; q^n x, q^n y \end{matrix} \right) t^n$$

satisfies $C_q u(x, y, t) = 0$. Using the relation $\frac{1}{q}K_q^-C_q = C_qK_q^-$, we have

$$(56) \quad C_q[E_q(sK_q^-)u](x, y, t) = 0,$$

where

(57)

$$\begin{aligned} [E_q(sK_q^-)u](x, y, t) &= \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} s^n}{(q; q)_n} K_q^{-n} u(x, y, t) \\ &= \sum_{k,n=0}^{\infty} \frac{(a, \frac{\omega}{t}; q)_n (q^{-n}, b; q)_k}{(q; q)_n (\frac{q^{1-n}t}{\omega}, c, q; q)_k} \\ &\quad \times {}_2\phi_2 \left(\begin{matrix} q^{-n+k}, b' \\ \frac{q^{1-n+k}t}{\omega}, c' \end{matrix}; q, \frac{-q^{k+1}yt}{\omega} \right) q^{\binom{k}{2}} \left(\frac{-qxt}{\omega} \right)^k t^n. \end{aligned}$$

Using Weisner's expansion and proceeding as in the above two cases, we get the following identity:

$$(58) \quad \sum_{k,n=0}^{\infty} \frac{(a, \frac{\omega}{t}; q)_n (q^{-n}, b; q)_k}{(q; q)_n (\frac{q^{1-n}t}{\omega}, c, q; q)_k} {}_2\phi_2 \left(\frac{q^{-n+k}}{\omega}, b'; q, \frac{-q^{k+1}yt}{\omega} \right) \left(\frac{-q^{\frac{k+1}{2}}xt}{\omega} \right)^k t^n$$

$$= \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} \Phi^{(2)} \left(\begin{matrix} q^{-n}; b, b' \\ c, c'; q^n x, q^n y \end{matrix} \right) {}_1\phi_0 \left(\begin{matrix} aq^n \\ - \end{matrix}; \omega \right) t^n,$$

where $\omega = \frac{s}{1-q}$.

6. CONCLUSION

The models and identities obtained in above sections can be generalized. For this, we need the m -fold q -integral transformation, defined as

$$(59) \quad h(\beta_i, \gamma_i) = \mathcal{I}[f(u_1, \dots, u_m)]$$

$$= \Gamma_q \left(\begin{matrix} \gamma_1, \dots, \gamma_m \\ \beta_1, \dots, \beta_m, \gamma_1 - \beta_1, \dots, \gamma_m - \beta_m \end{matrix} \right)$$

$$\times \underbrace{\int_0^1 \dots \int_0^1}_{m\text{-fold}} \prod_{i=1}^m u_i^{\beta_i-1} \frac{(qu_1, \dots, qu_m; q)_{\infty}}{(q^{\gamma_1-\beta_1}u_1, \dots, q^{\gamma_m-\beta_m}u_m; q)_{\infty}}$$

$$\times f(u_1, \dots, u_m) d_q u_1 \dots d_q u_m,$$

$$\operatorname{Re} \gamma_i > \operatorname{Re} \beta_i > 0, \quad i = 1, 2, \dots, m.$$

Putting $z = \sum_{i=1}^m u_i x_i$ in Model I (*i.e.* eqs. (22) and (23)), and using

$$(60) \quad z \Delta_z = u_1 \Delta_{u_1} + u_2 \Delta_{u_2} T_{u_1} + \dots + u_m \Delta_{u_m} T_{u_1} \dots T_{u_{m-1}},$$

$$(61) \quad T_z f(z) = T_{u_1} \dots T_{u_m} f\left(\sum_{i=1}^m u_i x_i\right),$$

we get the transformed $(m+1)$ -variable model as

$$(62) \quad K_q^+ = t[(x_1 \Delta_{x_1} + \dots + x_m \Delta_{x_m} T_{x_1} \dots T_{x_{m-1}}) T_t + t \Delta_t],$$

$$K_q^- = t^{-1} T_{x_1}^{-1} \dots T_{x_m}^{-1}$$

$$\times \left[\left(1 - x_1 T_t \frac{1 - q^{\beta_1}}{1 - q^{\gamma_1}} E_{\beta_1 \gamma_1} - \dots - x_m T_t \frac{1 - q^{\beta_m}}{1 - q^{\gamma_m}} E_{\beta_m \gamma_m} \right) \right.$$

$$\left. \times (x_1 \Delta_{x_1} + \dots + x_m \Delta_{x_m} T_{x_1} \dots T_{x_{m-1}}) - t \Delta_t \right],$$

$$K_q^0 = t \Delta_t,$$

with the basis functions as

$$(63) \quad h_{\lambda} = \phi_{-1;1;\dots;1}^{1;1;\dots;1} \left(\begin{matrix} q^{\lambda} : b_1; \dots; b_m; x_1, \dots, x_m \\ - : c_1; \dots; c_m \end{matrix} \right) t^{\lambda}$$

$$= \sum_{k_1, \dots, k_m=0}^{\infty} \frac{(q^{\lambda}; q)_{k_1+\dots+k_m} (b_1; q)_{k_1} \dots (b_m; q)_{k_m}}{(c_1, q; q)_{k_1} \dots (c_m, q; q)_{k_m}} x_1^{k_1} \dots x_m^{k_m} t^{\lambda}, \quad \lambda \in S,$$

where $b_i = q^{\beta_i}$, $c_i = q^{\gamma_i}$.

Using the same techniques as given in section 5, the above model, and similar other $(m + 1)$ -variable models, can be exploited for obtaining identities.

ACKNOWLEDGEMENTS

The author wishes to thank the referee for his valuable suggestions.

REFERENCES

- [1] L. C. BIEDENHARN, *The quantum group $SU_q(2)$ and a q -analogue of the Boson operators*, J. Phys. A: Math. Gen. **22**(1989), L873-L878. MR **90k**:17027
- [2] V. CHARI and A. PRESSLEY, *A guide to quantum groups*, Cambridge Univ. Press, 1994. MR **95j**:17010
- [3] V. G. DRINFELD, *Quantum Groups*, Proc. ICM, Vol. 1 (1987), 798-820. MR **89f**:17017
- [4] H. EXTON, *q -Hypergeometric functions and Applications*, Ellis-Horwood, 1983. MR **85g**:33001
- [5] L. D. FADDEEV, *Quantum groups*, Bol. Soc. Brasil Mat. **20**(1989) 47-54. MR **92f**:17016
- [6] R. FLOREANINI and L. VINET, *$U_q(sl(2))$ and q -Special functions*, Contemporary Math. **160**(1994), 85-100. MR **95b**:33054
- [7] —, *q -orthogonal polynomials and the orthogonal quantum group*, Lett. Math. Phys. **22**(1991) 45-54. MR **93g**:33015
- [8] —, *The metaplectic representation of $su_q(1,1)$ and the q -Gegenbauer polynomials*, J. Math. Phys. **33**(1992), 1358-1363. MR **93f**:33020
- [9] —, *Quantum algebras and q -special functions*, Ann. Phys. **221**(1993) 53-70. MR **94d**:33014
- [10] —, *On the quantum group and quantum algebras approach to q -special functions*, Lett. Math. Phys. **27**(1993) 179-190. MR **94e**:33040
- [11] —, *Using quantum algebras in q -special function theory*, Phys. Lett. A **170**(1992) 21-28. MR **94f**:33032
- [12] —, *A model for the continuous q -ultraspherical polynomials*, J. Math. Phys. **36**(1995), 3800-3813. MR **96d**:33012
- [13] G. GASPER and M. RAHMAN, *Basic hypergeometric series*, Cambridge Univ. Press, 1990. MR **91d**:33034
- [14] M. JIMBO, *A q -difference analogue of $U(g)$ and the Yang-Baxter equation*, Lett. Math. Phys. **10**(1985), 63-69. MR **86k**:17008
- [15] J. C. JANTZEN, *Lectures on quantum groups*, Graduate Series in Maths 6, AMS, 1996. MR **96m**:17029
- [16] E. G. KALNINS, H. L. MANOCHA and W. MILLER, *Harmonic analysis and expansion formulas for the two variable hypergeometric functions*, Stud. Appl. Math. **66**(1982), 69-89. MR **84a**:58091
- [17] —, *Models of q -algebra representations: Tensor products of special unitary and oscillator algebras*, J. Math. Phys. **33**(1992), 2365-2383. MR **93d**:17021
- [18] E. G. KALNINS and W. MILLER, *Models of q -algebra representations: q -integral transforms and addition theorems*, J. Math. Phys. **35**(1994), 1951-1975. MR **95c**:33030
- [19] E. G. KALNINS, W. MILLER and S. MUKHERJEE, *Models of q -algebra representations: The group of plane motions*, SIAM J. Math. Anal. **25**(1994), 513-527. MR **95c**:33029
- [20] —, *Models of q -algebra representations: Matrix elements of the q -oscillator algebra*, J. Math. Phys. **34**(1993), 5333-5356. MR **94j**:17012
- [21] —, *Models of q -algebra representations: Matrix elements of $U_q(su_2)$* , Contemporary Math., AMS **160**(1994), 191-208. MR **95b**:33055
- [22] C. KASSEL, *Quantum groups*, Springer Verlag, 1995. MR **96e**:17041
- [23] P. P. KULISH and N. YU RESHETIKHIN, *Quantum linear problem for the Sine-Gordon equation and higher representations*, J. Sov. Math. **23**(1983) 2435-2441.
- [24] S. MAJID, *Foundations of quantum group theory*, Cambridge Univ. Press, 1995. MR **97g**:17016
- [25] H. L. MANOCHA, *On models of irreducible q -representations of $sl(2, C)$* , Appl. Anal. **37** (1990), 19-47. MR **92h**:33042
- [26] A. J. MACFARLANE, *On the q -analogues of the quantum harmonic oscillator and the quantum group $SU_q(2)$* , J. Phys. A: Math. Gen. **22**(1989), 4581-4588. MR **91f**:81067
- [27] W. MILLER, *Lie theory and Special functions*, Academic Press, 1968. MR **41**:8736

- [28] G. RIDEAU and P. WINTERNITZ, *Representation of the quantum algebra $su_q(2)$ on a real two dimensional sphere*, J. Math. Phys. **34**(1993), 6030-6044. MR **94j**:33034
- [29] V. SAHAI, *Lie theory of hypergeometric functions ${}_3F_2$ by means of Euler transformation*, J. Indian Math. Soc. **61**(1995), 135-152. CMP 97:15
- [30] —, *q -Representations of the Lie algebras $\mathcal{G}(a, b)$* , J. Indian Math. Soc. **63**(1997), 83-98. CMP 98:12
- [31] E. K. SKLYANIN, *Some algebraic structures connected with the Yang-Baxter equation*, Funct. Anal. Appl. **16**(1982) 27-34, 96. MR **84c**:82004
- [32] —, *Some algebraic structures connected with the Yang-Baxter equation. Representations of quantum algebras*, Funct. Anal. Appl. **17**(1983) 34-48. MR **85k**:82011
- [33] H. M. SRIVASTAVA and H. L. MANOCHA, *A treatise on generating functions*, Ellis-Horwood, 1984. MR **85m**:33016
- [34] S. L. WORONOWICZ, *Compact matrix pseudogroups*, Comm. Math. Phys. **111**(1987) 613-665. MR **88m**:46079
- [35] —, *Twisted $SU(2)$ group. An example of noncommutative differential calculus*, Publ. RIMS Kyoto Univ. **23**(1987) 117-181. MR **88h**:46130

DEPARTMENT OF MATHEMATICS & ASTRONOMY, LUCKNOW UNIVERSITY, LUCKNOW – 226 007,
INDIA

E-mail address: vsahai@hotmail.com