THE ROGERS-RAMANUJAN IDENTITIES, THE FINITE GENERAL LINEAR GROUPS, AND THE HALL-LITTLEWOOD POLYNOMIALS

JASON FULMAN

(Communicated by Ronald M. Solomon)

ABSTRACT. We connect Gordon's generalization of the Rogers-Ramanujan identities with the Hall-Littlewood polynomials and with generating functions which arise in a probabilistic setting in the finite general linear groups. This yields a Rogers-Ramanujan type product formula for the $n \rightarrow \infty$ probability that an element of $GL(n, q)$ or $Mat(n, q)$ is semisimple.

1. BACKGROUND AND NOTATION

The Rogers-Ramanujan identities are among the most remarkable partition identities in number theory and combinatorics. This paper will be concerned with the following generalization of the Rogers-Ramanujan identities, due to Gordon. Let $(x)_n$ denote $(1-x)(1-x^2)\cdots(1-x^n)$.

Theorem 1 ([A, page 111]). *For $1 \leq i \leq k, k \geq 2$, and complex x with $|x| < 1$,*

$$\sum_{n_1, \dots, n_{k-1} \geq 0} \frac{x^{N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}}}{(x)_{n_1} \cdots (x)_{n_{k-1}}} = \prod_{\substack{r=1 \\ r \neq 0, \pm i \pmod{2k+1}}}^{\infty} \frac{1}{1-x^r}$$

where $N_j = n_j + \dots + n_{k-1}$.

Gordon's generalization of the Rogers-Ramanujan identities has been widely studied and appears in many places in mathematics and physics. Andrews [A] discusses combinatorial aspects of these identities. In an important series of papers, Lepowsky and Wilson [LW1], [LW2], [LW3] connect the Gordon identities with affine Lie algebras and structures that they called Z -algebras (later interpreted as parafermion algebras in conformal field theory). Meurman and Primc [MP] solve a problem left open in [LW3], proving the independence of a Z -algebra basis and obtaining a Z -algebra proof of the Gordon identities. Feigin and Frenkel [FF] interpret the Gordon identities as a character formula for the Virasoro algebra. Andrews, Baxter, and Forrester [ABF] and Warnaar [W] relate the Gordon identities with statistical mechanics. For some number theoretic connections see the conference proceedings [AABRR].

We use the following standard notation from the theory of partitions. Call $\lambda = (\lambda_1, \lambda_2, \dots)$ a partition of $n = |\lambda|$ if $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ where the λ_i are

Received by the editors March 6, 1998.

1991 *Mathematics Subject Classification*. Primary 20P05, 05E05.

a sequence of positive integers stabilizing to 0 such that $\sum_i \lambda_i = n$. The λ_i are referred to as the parts of λ . Let $m_i(\lambda)$ be the number of parts of λ of size i , and set $\lambda'_i = m_i(\lambda) + m_{i+1}(\lambda) + \dots$. Define $n(\lambda)$ by $\sum_{i \geq 1} (i-1)\lambda_i$. Let $[u^n]f(u)$ denote the coefficient of u^n in a power series $f(u)$.

2. MAIN RESULTS

To begin we recall the Hall-Littlewood polynomials associated to a partition λ (page 208 of [Mac]). Let n be any integer such that $n \geq \lambda_1$. The permutation $w \in S_n$ acts on the variables $x_1, \dots, x_n$ by sending x_i to $x_{w(i)}$. Letting t be a complex number, the Hall-Littlewood polynomials are defined as

$$P_\lambda(x_1, \dots, x_n; t) = \frac{1}{\prod_{i \geq 0} \prod_{r=1}^{m_i(\lambda)} \frac{1-t^r}{1-t}} \sum_{w \in S_n} w(x_1^{\lambda_1} \dots x_n^{\lambda_n} \prod_{i < j} \frac{x_i - tx_j}{x_i - x_j}).$$

At first glance it is not obvious that these are polynomials, but the denominators cancel out after the symmetrization. The Hall-Littlewood polynomials interpolate between the Schur functions ($t = 0$) and the monomial symmetric functions ($t = 1$).

Theorem 2. For $q > 1$ and an integer $k \geq 2$,

$$\begin{aligned} \sum_{\lambda: \lambda_1 < k} \frac{P_\lambda(\frac{1}{q}, \frac{1}{q^2}, \frac{1}{q^3}, \frac{1}{q^4}, \dots; \frac{1}{q})}{q^{n(\lambda)}} &= \prod_{\substack{r=1 \\ r \neq 0, \pm k \pmod{2k+1}}}^{\infty} \left(\frac{1}{1 - \frac{1}{q^r}} \right), \\ \sum_{\lambda: \lambda_1 < k} \frac{P_\lambda(\frac{1}{q^2}, \frac{1}{q^3}, \frac{1}{q^4}, \frac{1}{q^5}, \dots; \frac{1}{q})}{q^{n(\lambda)}} &= \prod_{\substack{r=1 \\ r \neq 0, \pm 1 \pmod{2k+1}}}^{\infty} \left(\frac{1}{1 - \frac{1}{q^r}} \right). \end{aligned}$$

Proof. Macdonald's principal specialization formula (page 337 of [Mac]) states that

$$P_\lambda(\frac{1}{q}, \frac{1}{q^2}, \frac{1}{q^3}, \frac{1}{q^4}, \dots; \frac{1}{q}) = \frac{1}{q^{|\lambda|+n(\lambda)}} \prod_i \frac{1}{(\frac{1}{q})_{m_i(\lambda)}}.$$

Combining this with the elementary fact that $n(\lambda) = \sum_i \binom{\lambda'_i}{2}$ shows that

$$\frac{P_\lambda(\frac{1}{q}, \frac{1}{q^2}, \frac{1}{q^3}, \frac{1}{q^4}, \dots; \frac{1}{q})}{q^{n(\lambda)}} = \frac{1}{q^{\sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q})_{m_i(\lambda)}}.$$

The first equality of the theorem follows by applying Theorem 1 with $i = k, x = \frac{1}{q}, n_j = \lambda_j$ and $N_j = \lambda'_j$. For the second equality, observe that for u complex,

$$\begin{aligned} \frac{P_\lambda(\frac{u}{q}, \frac{u}{q^2}, \frac{u}{q^3}, \frac{u}{q^4}, \dots; \frac{1}{q})}{q^{n(\lambda)}} &= u^{|\lambda|} \frac{P_\lambda(\frac{1}{q}, \frac{1}{q^2}, \frac{1}{q^3}, \frac{1}{q^4}, \dots; \frac{1}{q})}{q^{n(\lambda)}} \\ &= \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q})_{m_i(\lambda)}}. \end{aligned}$$

Now set $u = \frac{1}{q}$ and apply Theorem 1 with $i = 1, x = \frac{1}{q}, n_j = \lambda_j$ and $N_j = \lambda'_j$. $\square$

Remark. Stembridge [Ste] used properties of the Hall-Littlewood polynomials as a tool in giving proofs of the Rogers-Ramanujan identities. The statement of Theorem 2 gives a direct connection.

Recall that the conjugacy classes of $GL(n, q)$ are parameterized by rational canonical form (Chapter 6 of Herstein [H]). This form corresponds to the following combinatorial data. To each monic non-constant irreducible polynomial ϕ over a field of q elements, associate a partition λ_ϕ of some non-negative integer $|\lambda_\phi|$. Let $\deg(\phi)$ denote the degree of ϕ . This data represents a conjugacy class of $GL(n, q)$ if and only if $|\lambda_z| = 0$ and $\sum_\phi |\lambda_\phi| \deg(\phi) = n$.

Definition. For $\alpha \in GL(n, q)$ and ϕ a monic, irreducible polynomial over F_q , a field of q elements, define $\lambda_\phi(\alpha)$ to be the partition corresponding to the polynomial ϕ in the rational canonical form of α .

The following elementary lemmas will be of use in studying the partitions λ_ϕ .

Lemma 1. *If the Taylor series of $f(u)$ around 0 converges at $u = 1$, then*

$$\lim_{n \rightarrow \infty} [u^n] \frac{f(u)}{1-u} = f(1).$$

Proof. Write the Taylor expansion $f(u) = \sum_{n=0}^{\infty} a_n u^n$. Then observe that $[u^n] \frac{f(u)}{1-u} = \sum_{i=0}^n a_i$. $\square$

Lemma 2. *For $t \geq 1$, q a prime power, and u a formal variable,*

$$\prod_{\phi \text{ irred.}} \left(1 - \frac{u^{\deg(\phi)}}{q^{t \cdot \deg(\phi)}}\right) = 1 - \frac{u}{q^{t-1}}.$$

Proof. Assume that $t = 1$, the general case following by replacing u with $\frac{u}{q^{t-1}}$. Expanding $\frac{1}{1 - \frac{u^{\deg(\phi)}}{q^{\deg(\phi)}}}$ as a geometric series, unique factorization in $F_q[x]$ implies that the coefficient of u^d in the reciprocal of the left hand side is $\frac{1}{q^d}$ times the number of monic polynomials of degree d , hence 1. Comparing with the reciprocal of the right hand side proves the lemma. $\square$

Lemma 3. *For $t \geq 1$, q a prime power, and u a formal variable,*

$$1 - u = \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}}\right).$$

Proof. By Lemma 2,

$$\prod_{r=1}^{\infty} \left(1 - \frac{u}{q^{r-1}}\right) = \prod_{\phi \text{ irred.}} \prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}}\right).$$

The lemma follows by cancelling the terms corresponding to $\phi = z$. $\square$

Lemma 4. *For q a prime power and u complex with $|u| \leq 1$,*

$$\sum_{\lambda} \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)}} = \prod_{r=1}^{\infty} \left(\frac{1}{1 - \frac{u}{q^r}}\right).$$

Proof. On page 225 of [Mac] it is proved that

$$\prod_{r=1}^{\infty} (1 - x_i) \sum_{\lambda} t^{n(\lambda)} P_{\lambda}(x_1, x_2, x_3, \dots; t) = 1.$$

Applying Macdonald's principal specialization formula with $t = \frac{1}{q}$, $x_i = \frac{u}{q^i}$ as in Theorem 2 proves the result. $\square$

Lemma 5 (Euler). *For $|q| > 1$ and u complex with $|u| \leq 1$,*

1. $\prod_{r=1}^{\infty} (1 - \frac{u}{q^r}) = \sum_{n=0}^{\infty} \frac{(-u)^n}{(q^n-1)\cdots(q-1)},$
2. $\prod_{r=1}^{\infty} (\frac{1}{1-\frac{u}{q^r}}) = \sum_{n=0}^{\infty} \frac{u^n q^{\binom{n}{2}}}{(q^n-1)\cdots(q-1)}.$

Furthermore, these Taylor series converge at $u = 1$.

Theorem 3 relates the Gordon identities with probability in the finite general linear groups.

Theorem 3. *Let ϕ be a monic, irreducible polynomial over F_q . Let $k \geq 2$ be an integer. Then the $n \rightarrow \infty$ limit of the chance that a uniformly chosen element α of $GL(n, q)$ has the largest part of the partition $\lambda_\phi(\alpha)$ less than k is equal to*

$$\prod_{\substack{r=1 \\ r=0, \pm k \pmod{2k+1}}}^{\infty} (1 - \frac{1}{q^{r \cdot \deg(\phi)}}).$$

Proof. Assume for simplicity that $\phi = z - 1$. From the proof it will be clear that the general case follows. Stong [Sto] used Kung's [Ku] formula for the sizes of the conjugacy classes of $GL(n, q)$ to find a “cycle index” for the general linear groups. Using the notation

$$d_i(\lambda) = 1m_1(\lambda) + 2m_2(\lambda) + \cdots + (i-1)m_{i-1}(\lambda) + i(m_i(\lambda) + m_{i+1}(\lambda) + \cdots + m_j(\lambda)),$$

he obtained the equality

$$\begin{aligned} 1 + \sum_{n=1}^{\infty} \frac{u^n}{|GL(n, q)|} \sum_{\alpha \in GL(n, q)} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} x_{\phi, \lambda_\phi(\alpha)} \\ = \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \left[\sum_{\lambda} x_{\phi, \lambda} \frac{u^{|\lambda| \deg(\phi)}}{\prod_i \prod_{k=1}^{m_i(\lambda)} (q^{\deg(\phi)d_i} - q^{\deg(\phi)(d_i-k)})} \right]. \end{aligned}$$

Observe that

$$\begin{aligned} \prod_i \prod_{k=1}^{m_i} (q^{d_i} - q^{(d_i-k)}) &= q^{\sum_i m_i(\lambda)d_i(\lambda)} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)} \\ &= q^{\sum_i m_i(\lambda)[(\sum_{h < i} h m_h(\lambda)) + i m_i(\lambda) + \sum_{i < k} i m_k(\lambda)]} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)} \\ &= q^{\sum_i [i m_i(\lambda)^2 + 2 m_i(\lambda) \sum_{h < i} h m_h(\lambda)]} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)} \\ &= q^{\sum_i [\sum_{h < i} m_h(\lambda)]^2} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)} \\ &= q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)}. \end{aligned}$$

Combining this observation with Lemma 3 shows that

$$\begin{aligned} & 1 + \sum_{n=1}^{\infty} \frac{u^n}{|GL(n, q)|} \sum_{\alpha \in GL(n, q)} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} x_{\phi, \lambda_{\phi}(\alpha)} \\ &= \frac{1}{1-u} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \left[\prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}} \right) \right] \left[\sum_{\lambda} x_{\phi, \lambda} \frac{u^{|\lambda| \deg(\phi)}}{q^{\deg(\phi) \sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q^{\deg(\phi)}} \right)_{m_i(\lambda)}} \right]. \end{aligned}$$

Setting $x_{z-1, \lambda} = 0$ if the largest part of λ is greater than or equal to k , and all $x_{\phi, \lambda} = 1$ otherwise shows by Lemma 4 that the sought probability is

$$\lim_{n \rightarrow \infty} [u^n] \frac{1}{1-u} \left(\prod_{r=1}^{\infty} \left(1 - \frac{u}{q^r} \right) \right) \left(\sum_{\lambda: \lambda_1 < k} \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q} \right)_{m_i(\lambda)}} \right).$$

By Lemmas 4 and 5, Lemma 1 applies and

$$\begin{aligned} & \lim_{n \rightarrow \infty} [u^n] \frac{1}{1-u} \left(\prod_{r=1}^{\infty} \left(1 - \frac{u}{q^r} \right) \right) \left(\sum_{\lambda: \lambda_1 < k} \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q} \right)_{m_i(\lambda)}} \right) \\ &= \left(\prod_{r=1}^{\infty} \left(1 - \frac{1}{q^r} \right) \right) \left(\sum_{\lambda: \lambda_1 < k} \frac{1}{q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q} \right)_{m_i(\lambda)}} \right). \end{aligned}$$

The theorem follows from the Gordon identities with $i = k, x = \frac{1}{q}, n_j = m_j(\lambda)$, and $N_j = \lambda'_j$. $\square$

Let $Mat(n, q)$ denote the set of $n \times n$ matrices with entries in the finite field F_q . Recall that $\alpha \in Mat(n, q)$ is said to be semisimple if it is diagonalizable over the algebraic closure $\bar{F}_q$.

Theorem 4. *The $n \rightarrow \infty$ limiting probability that an element of $GL(n, q)$ is semisimple is*

$$\prod_{\substack{r=1 \\ r \equiv \pm 2 \pmod{5}}}^{\infty} \frac{\left(1 - \frac{1}{q^{r-1}} \right)}{\left(1 - \frac{1}{q^r} \right)}.$$

Proof. From the theory of Jordan canonical forms, an element α of $GL(n, q)$ is semisimple if and only if the largest part of $\lambda_{\phi}(\alpha)$ is less than two for all ϕ . Thus by the cycle index for $GL(n, q)$ (see the proof of Theorem 3) and Lemma 3, the sought probability is

$$\begin{aligned} & \lim_{n \rightarrow \infty} [u^n] \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \sum_{\lambda: \lambda_1 < 2} \frac{u^{\deg(\phi) |\lambda|}}{q^{\deg(\phi) \sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q^{\deg(\phi)}} \right)_{m_i(\lambda)}} \\ &= \lim_{n \rightarrow \infty} [u^n] \frac{1}{1-u} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \left(\prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}} \right) \right) \\ & \quad \cdot \left(\sum_{\lambda: \lambda_1 < 2} \frac{u^{\deg(\phi) |\lambda|}}{q^{\deg(\phi) \sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q^{\deg(\phi)}} \right)_{m_i(\lambda)}} \right). \end{aligned}$$

By Theorem 1 with $i = 1, k = 2, x = \frac{1}{q^{\deg(\phi)}}, n_j = \lambda_j, N_j = \lambda'_j$ and Lemma 2, the desired probability becomes

$$\begin{aligned}
& \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} \left(1 - \frac{1}{q^{r \cdot \deg(\phi)}}\right) \\
&= \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} \left(\frac{1}{1 - \frac{1}{q^r}}\right) \prod_{\phi \text{ irred.}} \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} \left(1 - \frac{1}{q^{r \cdot \deg(\phi)}}\right) \\
&= \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} \frac{\left(1 - \frac{1}{q^{r-1}}\right)}{\left(1 - \frac{1}{q^r}\right)}.
\end{aligned}$$

□

Theorem 5. *The $n \rightarrow \infty$ limiting probability that an element of $\text{Mat}(n, q)$ is semisimple is*

$$\prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} \left(1 - \frac{1}{q^{r-1}}\right).$$

Proof. The orbits of $GL(n, q)$ on $\text{Mat}(n, q)$ under conjugation are also parameterized by the data λ_ϕ . However the polynomial z may appear with non-zero multiplicity, so the restriction that $|\lambda_z| = 0$ does not apply. Stong [Sto] obtained the generating function equality

$$\begin{aligned}
& 1 + \sum_{n=1}^{\infty} \frac{u^n}{|\text{GL}(n, q)|} \sum_{\alpha \in \text{Mat}(n, q)} \prod_{\phi \text{ irred.}} x_{\phi, \lambda_\phi(\alpha)} \\
&= \prod_{\phi \text{ irred.}} \left[\sum_{\lambda} x_{\phi, \lambda} \frac{u^{|\lambda| \deg(\phi)}}{\prod_i \prod_{k=1}^{m_i(\lambda)} (q^{\deg(\phi) d_i} - q^{\deg(\phi)(d_i - k)})} \right],
\end{aligned}$$

where $d_i(\lambda)$ is defined by

$$\begin{aligned}
d_i(\lambda) &= 1m_1(\lambda) + 2m_2(\lambda) + \cdots + (i-1)m_{i-1}(\lambda) \\
&\quad + i(m_i(\lambda) + m_{i+1}(\lambda) + \cdots + m_j(\lambda)).
\end{aligned}$$

Manipulations identical to those in Theorem 3 show that

$$\begin{aligned}
& 1 + \sum_{n=1}^{\infty} \frac{u^n}{|\text{GL}(n, q)|} \sum_{\alpha \in \text{Mat}(n, q)} \prod_{\phi \text{ irred.}} x_{\phi, \lambda_\phi(\alpha)} \\
&= \frac{1}{1-u} \prod_{\phi \text{ irred.}} \left[\prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}}\right) \right] \left[\sum_{\lambda} x_{\phi, \lambda} \frac{u^{|\lambda| \deg(\phi)}}{q^{\deg(\phi) \sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q^{\deg(\phi)}}\right)^{m_i(\lambda)}} \right].
\end{aligned}$$

Arguing as in Theorem 4 shows that the $n \rightarrow \infty$ limit of the chance that an element of $Mat(n, q)$ is semisimple is

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \frac{|GL(n, q)|}{q^{n^2}} [u^n] \prod_{\phi \text{ irred.}} \sum_{\lambda: \lambda_1 < 2} \frac{u^{deg(\phi)|\lambda|}}{q^{deg(\phi) \sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q^{deg(\phi)}})_{m_i(\lambda)}} \\
&= \lim_{n \rightarrow \infty} \frac{|GL(n, q)|}{q^{n^2}} [u^n] \frac{1}{1-u} \frac{1}{\prod_{r=1}^{\infty} (1 - \frac{u}{q^r})} \prod_{\phi \text{ irred.}} \\
&\quad \left[\prod_{r=1}^{\infty} (1 - \frac{u^{deg(\phi)}}{q^{r \cdot deg(\phi)}}) \sum_{\lambda: \lambda_1 < 2} \frac{u^{deg(\phi)|\lambda|}}{q^{deg(\phi) \sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q^{deg(\phi)}})_{m_i(\lambda)}} \right] \\
&= \prod_{\phi \text{ irred.}} \left[\prod_{r=1}^{\infty} (1 - \frac{1}{q^{r \cdot deg(\phi)}}) \right] \left[\sum_{\lambda: \lambda_1 < 2} \frac{1}{q^{deg(\phi) \sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q^{deg(\phi)}})_{m_i(\lambda)}} \right] \\
&= \prod_{\phi \text{ irred.}} \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} (1 - \frac{1}{q^{r \cdot deg(\phi)}}) = \prod_{\substack{r=1 \\ r=0, \pm 2 \pmod{5}}}^{\infty} (1 - \frac{1}{q^{r-1}}). \quad \square
\end{aligned}$$

Theorems 2 and 3 suggest a connection between the Hall-Littlewood polynomials and the finite general linear groups. Theorem 6 makes this connection precise.

Theorem 6. *For q a prime power and u a formal variable,*

$$\begin{aligned}
& (1-u) \left[1 + \sum_{n=1}^{\infty} \frac{u^n}{|GL(n, q)|} \sum_{\alpha \in GL(n, q)} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} x_{\phi, \lambda_{\phi}(\alpha)} \right] \\
&= \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \prod_{r=1}^{\infty} (1 - \frac{u^{deg(\phi)}}{q^{r \cdot deg(\phi)}}) \left[\frac{P_{\lambda} \left((\frac{u}{q})^{deg(\phi)}, (\frac{u}{q})^{2deg(\phi)}, (\frac{u}{q})^{3deg(\phi)}, \dots, (\frac{1}{q})^{deg(\phi)} \right)}{q^{n(\lambda) \cdot deg(\phi)}} \right].
\end{aligned}$$

Proof. The proof of Theorem 3 contained the equality

$$\begin{aligned}
& (1-u) \left[1 + \sum_{n=1}^{\infty} \frac{u^n}{|GL(n, q)|} \sum_{\alpha \in GL(n, q)} \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} x_{\phi, \lambda_{\phi}(\alpha)} \right] \\
&= \prod_{\substack{\phi \neq z \\ \phi \text{ irred.}}} \left[\prod_{r=1}^{\infty} (1 - \frac{u^{deg(\phi)}}{q^{r \cdot deg(\phi)}}) \right] \left[\sum_{\lambda} x_{\phi, \lambda} \frac{u^{|\lambda|deg(\phi)}}{q^{deg(\phi) \sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q^{deg(\phi)}})_{m_i(\lambda)}} \right].
\end{aligned}$$

The theorem now follows from the use of Macdonald's principal specialization formula (as in Theorem 2) to conclude that

$$\frac{P_{\lambda}(\frac{u}{q}, \frac{u}{q^2}, \frac{u}{q^3}, \frac{u}{q^4}, \dots, \frac{1}{q})}{q^{n(\lambda)}} = \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i (\frac{1}{q})_{m_i(\lambda)}}. \quad \square$$

Remark. Although Theorem 6 followed easily from techniques in Theorems 2 and 3, its formulation admits an attractive probabilistic interpretation. Fix u such that $0 < u < 1$ and fix an irreducible monic polynomial $\phi \neq z$. Then pick a natural number with probability of getting n equal to $(1-u)u^n$. Finally, choose α uniformly at random in $GL(n, q)$ and let $\lambda_{\phi}(\alpha)$ be the random partition so defined. This procedure induces a probability measure on the set of all partitions of all integers.

Theorem 6 implies that measures obtained for different ϕ are independent, and that the mass the measure for a given ϕ assigns to a partition λ is equal to

$$\prod_{r=1}^{\infty} \left(1 - \frac{u^{\deg(\phi)}}{q^{r \cdot \deg(\phi)}}\right) \left[\frac{P_{\lambda} \left(\left(\frac{u}{q}\right)^{\deg(\phi)}, \left(\frac{u}{q}\right)^{2\deg(\phi)}, \left(\frac{u}{q}\right)^{3\deg(\phi)}, \dots; \left(\frac{1}{q}\right)^{\deg(\phi)} \right)}{q^{n(\lambda) \cdot \deg(\phi)}} \right].$$

This connection with symmetric functions leads to a probabilistic algorithm for growing the random partitions λ_{ϕ} . This viewpoint is used in [Fu1] to give probabilistic proofs of some group theoretic results of Steinberg, Rudvalis/Shinoda, and Lusztig. Cycle indices for the finite unitary, symplectic, and orthogonal groups appear in [Fu2].

3. CONCLUSION

This paper has offered connections between Gordon's generalization of the Rogers-Ramanujan identities, the Hall-Littlewood polynomials, and generating functions which arise in the study of the finite general linear groups. The following questions seem natural.

- Are there analogs of the Gordon identities for the finite unitary, symplectic, and orthogonal groups? The Gordon identities do play a role in Misra's work on affine symplectic Lie algebras [Mi1] and in Mandia's work [Man] on the affine Lie algebras $B_l^{(1)}$, $F_4^{(1)}$, and $G_2^{(1)}$. Misra has also used Z -algebra methods to relate the Gordon identities to $\widetilde{A}_n$.
- Ian Macdonald has suggested that the results of this paper should carry over to the affine finite general linear groups. The Hall-Littlewood polynomials have affine analogs [EK].
- It is known (e.g. Kac [Ka]) that the product side of the Rogers-Ramanujan identities has interesting modular properties. Can this phenomenon be understood group theoretically in terms of the conjugacy classes of the general linear groups; i.e. is there "general linear group moonshine"? In this regard note that Jing [J1, J2] has connected the Hall-Littlewood polynomials with vertex operators.

ACKNOWLEDGEMENTS

This work is taken from the author's Ph.D. thesis, done under the supervision of Persi Diaconis. His idea of studying the random partitions λ_{ϕ} led to this work. We thank Ed Frenkel, Jim Lepowsky, and the referee for providing pointers to the Rogers-Ramanujan literature. This research was done under the generous 3-year support of the National Defense Science and Engineering Graduate Fellowship and the support of the Alfred P. Sloan Foundation Dissertation Fellowship.

REFERENCES

- [A] Andrews, G., *The theory of partitions. Encyclopedia of Mathematics and its Applications, Vol. 2.* Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976. MR **58**:27738
- [AABRR] Andrews, G., Askey, R., Berndt, B., Ramanathan, K., and Rankin, R., *Ramanujan revisited.* Proceedings of the Ramanujan Centenary Conference, Academic Press, Inc., Boston, 1988. MR **89c**:11001

- [ABF] *Andrews, G., Baxter, R. J., and Forrester, P. J.*: Eight-vertex SOS model and generalized Rogers-Ramanujan type identities, *J. Stat. Phys.* **35** (1984), 193-266. MR **86a**:82001
- [EK] *Etingof, P. and Kirillov, A.*, On the affine analogue of Jack and Macdonald polynomials, *Duke Math. J.* **78** (1995), 229-256. MR **97k**:17035
- [FF] *Feigin, B. and Frenkel, E.*: Coinvariants of nilpotent subalgebras of the Virasoro algebra and partition identities. I. M. Gelfand Seminar, *Adv. Soviet Math.* **16**, Part 1, 139-148, Amer. Math. Soc., Providence, RI, 1993. MR **94g**:17054
- [Fu1] *Fulman, J.*, A probabilistic approach toward conjugacy classes in the finite general linear and unitary groups, *J. Algebra* **212** (1999), 557-590. CMP 99:09
- [Fu2] *Fulman, J.*, Cycle indices for the finite classical groups, *J. Group Theory* **2** (1999), 251-289.
- [H] *Herstein, I.N.*, *Topics in algebra*. Second edition. Xerox College Publishing, Lexington, Mass. - Toronto, Ont., 1975. MR **50**:9456
- [J1] *Jing, N.*, Vertex operators and Hall-Littlewood symmetric functions, *Advances in Math.* **87** (1991), 226-248. MR **93c**:17039
- [J2] *Jing, N.* Boson-fermion correspondence for Hall-Littlewood polynomials, *J. Math. Physics* **36** (1995), 7073-7080. MR **96m**:17049
- [Ka] *Kac, V.G.*, *Modular invariance in mathematics and physics*. American Mathematical Society centennial publications, Vol. II (Providence, RI, 1988), 337-350, Amer. Math. Soc., Providence, RI, 1992. MR **93h**:17061
- [Ku] *Kung, J.*, The cycle structure of a linear transformation over a finite field, *Lin. Alg. Appl.* **36** (1981), 141-155. MR **82d**:15012
- [LW1] *Lepowsky, J. and Wilson, R. L.*, A new family of algebras underlying the Rogers-Ramanujan identities and generalizations, *Proc. Nat. Acad. Sci. USA* **78** (1981), 7254-7258. MR **82k**:10016
- [LW2] *Lepowsky, J. and Wilson, R.L.*, The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities, *Invent. Math.* **77** (1984), 199-290. MR **85m**:17008
- [LW3] *Lepowsky, J. and Wilson, R.L.*, The structure of standard modules, II: The case $A_1^{(1)}$ principal gradation, *Invent. Math.* **79** (1985), 417-442. MR **86g**:17014
- [Mac] *Macdonald, I.G.*, *Symmetric functions and Hall polynomials*. Second Edition. Clarendon Press, Oxford, 1995. MR **96h**:05207
- [Man] *Mandia, M.*, Structure of the level one standard modules for the affine Lie algebras $B_l^{(1)}$, $F_4^{(1)}$, and $G_2^{(1)}$, *Memoirs American Math. Soc.* **362**, 1987. MR **88h**:17023
- [Mi1] *Misra, K.*, Level one standard modules for affine symplectic Lie algebras, *Math. Ann.* **287** (1990), 287-302. MR **91j**:17036
- [Mi2] *Misra, K.*, Level two standard $\widetilde{A}_n$ -modules, *J. Algebra* **137** (1991), 56-76. MR **92a**:17040
- [MP] *Meurman, A. and Primc, M.*, Annihilating ideals of standard modules of $sl(2, C)^\sim$ and combinatorial identities, *Advances in Math.* **64** (1987), 177-240. MR **89c**:17031
- [Ste] *Stembridge, J.*: Hall-Littlewood functions, plane partitions, and the Rogers-Ramanujan identities, *Trans. Amer. Math. Soc.* **319** (1990), 469-498. MR **90j**:05021
- [Sto] *Stong, R.*, Some asymptotic results on finite vector spaces, *Adv. Appl. Math.* **9** (1988), 167-199. MR **89c**:05007
- [W] *Warnaar, S. Ole*, The Andrews-Gordon identities and q-multinomial coefficients, *Commun. Math. Phys.* **184** (1997), 203-232. MR **98m**:11108

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NEW HAMPSHIRE 03755-3551

E-mail address: Fulman@Dartmouth.Edu